

MS02.07.06 XAS STUDIES OF THE Cu_A CENTERS OF CYTOCHROME *c* OXIDASE: A UNIQUE BINUCLEAR COPPER CLUSTER. Ninian J. Blackburn[†], Simon de Vries[§], Robert A. Scott[‡], James Fee[§], Yi Lu[#], Chris Dennison[¶] and Gerard Canters[¶]. [†]Oregon Graduate Institute of Science and Technology; [§]Delft University of Technology; [‡]University of Georgia; [¶]University of California, San Diego; [#]University of Illinois, Urbana-Champaign; [¶]Leiden Institute of Chemistry

The Cu_A centers of cytochrome *c* oxidases are unique examples of a new type of binuclear copper cluster. X-ray crystallography of enzymes from beef heart, *Paracoccus*, and the engineered cyoA fragment of the quinol oxidase of *E. Coli* have provided a structural description of the site. The coppers are bridged by two cysteine ligands, and have an extremely short Cu-Cu distance of ~ 2.4 Å. X-ray absorption spectroscopy, which had previously predicted the short Cu-Cu distance, has been used to further refine the structural details of the site, in both the oxidized and reduced forms. Subtle changes are detected in the metrical parameters of the oxidized versus reduced proteins which suggest that the short distance may be the result, in part, of a weak metal-metal bond. These studies have been extended to include Cu_A derivatives of the blue proteins azurin and amicyanin produced by "loop-directed mutagenesis", in which the Cu_A -binding sequence has been introduced into the blue copper proteins.

MS02.07.07 STRUCTURE OF SUBUNIT II OF A QUINOL OXIDASE WITH REENGINEERED Cu_A SITE. Matthias Wilmanns, Kristina Djinovic, Pekka Lappalainen, Mark Kelly, Elisabeth Sauer-Deriksson & Matti Saraste, EMBL Heidelberg, Postfach 102209, D-69012 Heidelberg, Germany

The crystal structures of the periplasmic fragment from the wild-type CyoA subunit II of the *Escherichia coli* quinol oxidase and of a mutant with a reengineered dinuclear copper centre ("purple CyoA") have been solved at 2.3 and 2.5 Å, respectively. Quinol oxidases belong to the superfamily of cytochrome oxidases. This enzyme is a member of the protein complex that catalyses reduction of molecular oxygen to water and utilizes the free energy of this reaction to generate a transmembrane proton gradient during respiration. The electron entry site in subunit II is a mixed-valence dinuclear copper in the enzymes which oxidize cytochrome *c*. This centre has been lost during the evolution of the quinol-oxidizing branch of cytochrome oxidases.

CyoA is folded as a 11-stranded, mostly antiparallel β -sandwich followed by three α -helices. The dinuclear copper centre is located at the loops between strands $\beta 5$ - $\beta 6$ and $\beta 9$ - $\beta 10$. The two coppers are at 2.5 Å distance and symmetrically coordinated to the main ligands which are two bridging cysteines and two terminal histidines. The residues that are distinct in cytochrome *c* and quinol oxidases are around the dinuclear copper centre. A recent structure of CyoA with reduced dinuclear copper centre shows a virtual identical arrangement of the two coppers except for increased distances between the two terminal histidines and the copper ions. Structural comparison suggests a common ancestry for subunit II of cytochrome oxidase and blue copper proteins.

PS.02.07.08 CRYSTAL STRUCTURE OF CUCUMBER STELLACYANIN AT 1.7 Å RESOLUTION. P. John Hart¹, Aram N. Nersissian², Joan Selverstone Valentine², and David Eisenberg¹. ¹UCLA-DOE Laboratory of Structural Biology and Molecular Medicine, Box 951570, UCLA, Los Angeles, CA 90095. ²Department of Chemistry and Biochemistry, University of California, Los Angeles, CA 90024.

Stellacyanins are blue (type I) copper glycoproteins that differ from other cupredoxins (such as plastocyanin and azurin) in many

of their properties. They have an unusual copper ligand (Gln instead of Met found in other mononuclear blue copper proteins), they perform more rapid long-range electron transfer, and they exhibit pH-dependent, reversible EPR and electronic absorption spectra. Until now, stellacyanins have eluded structure determination. Here we report the refined three-dimensional crystal structure at 1.7 Å resolution of stellacyanin from cucumber peelings.

The overall fold of the cucumber stellacyanin copper-binding domain is organized in two β -sheets, one of three β -strands and one of four. Two α -helices are found in loop regions between β -strands. One side of the molecule is predominantly negatively-charged, and provides a possible interaction site for redox partners. The characteristic spectroscopic properties and electron transfer reactivity of stellacyanin, relative to other well characterized blue copper proteins, may be explained by a copper binding site that is solvent exposed, and the fact that the copper is held in a nearly tetrahedral geometry by a strong interaction with the Gln ligand.

PS02.07.09 XAFS AND CRYSTALLOGRAPHIC STUDIES OF AN AZURIN AND A BLUE COPPER NITRITE REDUCTASE FROM A DENITRIFYING BACTERIUM. S. Samar Hasnain^{1,2}, Fraser Dodd^{1,2}, Richard Strange¹, Gunter Grossmann¹, Lorreta Murphy¹, Zeldi Abraham³, Robert Eady³, Bary Smith³. ¹Molecular Biophysics Group, Daresbury Laboratory, Warrington WA4 4AD, ²School of Applied Sciences, De Montfort University, Leicester LE1 9bH, ³John Innes Centre, Norwich Science Park, Norwich, UK

Results from a high resolution (~ 1.7 Å) crystallographic studies of oxidised and reduced azurin crystals are compared to the XAFS studies of this new azurin.

Crystal structure study of the first blue copper nitrite reductase will be reported. Results from substrate and ligand binding studies will be presented and the chemical (structural) changes associated with such binding will be discussed in terms of reaction mechanism. Background references:

1. 'X-ray scattering provides direct evidence for a trimeric structure in solution of Nitrite Reductase from *Alcaligenes xylosoxidans*', J. G. Grossmann et al., *Biochemistry*, 32, 7360-7366 (1993)
2. 'Evidence of two distinct azurins in *Alcaligenes xylosoxidans*' (NCIMB 11015): potential electron donors to nitrite reductase', F. E. Dodd et al *Biochemistry*, 34, 10180-10186 (1995)
3. 'The substrate binding site in copper nitrite reductase and its similarity to Zn carbonic anhydrase', R. W. Strange et al. *Nature-Structural Biology* 4, 287-292 (1995)
4. 'Structure of a new azurin from *Alcaligenes xylosoxidans* at 1.9 Å resolution', F. E. Dodd, S. S. Hasnain et al *Acta Cryst D* 51 1052(1995)

PS02.07.10 CRYSTALLIZATION AND PRELIMINARY X-RAY STUDIES OF AZURIN-I AND AZURIN-II FROM DENITRIFYING BACTERIUM *ALCALIGENS XYLOSOXIDANS* GIFU 1051. Chunmin Li*, Tsuyoshi Inoue*, Masaharu Gotowda*, Kazuyuki Hamada*, Nobuya Nishio*, Shinichiro Suzuki**, Kazuya Yamaguchi**, and Yasushi Kai*. *Department of Applied Chemistry, Faculty of Engineering, Osaka University, Suita 565, **Department of Chemistry, Faculty of Science, Osaka University, Toyonaka 560, Japan.

Azurins are the small copper-containing proteins that function as electron transfer agents in the redox systems of some bacteria. It has been known for a long time that only one azurin is obtained from one species of bacteria, except for the case of *Methylobacterium J*. Recently, two azurins were found instead of the single previously identified one in both *Alcaligenes xylosoxidans* NCIB 11015 and GIFU 1051. Here we present our recent work on the crystallization and preliminary X-ray studies of Azurin-I and Azurin-II from Denitrifying Bacterium *Alcaligenes Xylosoxidans* GIFU 1051. Both azurins were crystallized by the hanging drop

vapor diffusion method at 20°C. The crystals of azurin-I belong to the monoclinic crystal system and have space group *C2* with unit cell parameters of $a=130.6\text{\AA}$, $b=54.4\text{\AA}$, $c=74.7\text{\AA}$, and $\beta=96.1^\circ$. Four molecules are in the asymmetric unit. The crystals of azurin-II belong to the tetragonal crystal system and have space group *P4122* with unit cell parameters $a=b=52.6\text{\AA}$, $c=100.7\text{\AA}$. Only one molecule is in the asymmetric unit. They diffract up to 2.8Å and 2.0Å resolutions, respectively. Their crystal structures were solved by molecular replacement method using AMoRe in the CCP4. Molecular structure of Azurin II from *Alcaligenes xylooxidans* NCIB 11015 was chosen as a starting model for both azurins. The refinement for both azurins is being carried out by XPLOR.

PS02.07.11 A NOVEL IRON CENTER IN DESULFOFERRODOXIN FROM *D. DESULFURICANS* ATCC 27774: CRYSTAL STRUCTURE AT 1.8 Å RESOLUTION. Pedro M. Matias¹, Ana Coelho^{1,2}, Maria A. Carrondo^{1,3}, Vilmos Fülöp⁴, Ana Gonzalez⁵ and Andy Thompson⁶. ¹ITQB, Universidade Nova de Lisboa, 2780 OEIRAS, Portugal; ²Universidade de Évora, 7000 ÉVORA, Portugal; ³IST, Universidade Técnica de Lisboa, 1000 LISBOA, Portugal; ⁴LMB and OCMS, University of Oxford, Oxford OX1 3QU, UK ⁵ESRF, BP-220, 38043 GRENOBLE CEDEX France; ⁶EMBL Grenoble Outstation, BP-156, 38042 GRENOBLE CEDEX France

A novel iron centre in desulfoferrodoxin from sulfate reducing bacteria *D. desulfuricans* ATCC 27774 (DFX) has been structurally characterized: this center contains a single Fe atom octahedrally coordinated to four equatorial histidines, one axial cysteine and one axial water molecule, and unlike most other iron centres in proteins, it is exposed to the solvent, rather than buried within the bulk of the polypeptide chain. The three dimensional structure of DFX contains two domains linked by a short stretch of amino acid residues. The larger domain is associated with the novel Fe centre. The smaller domain contains a desulfoferrodoxin-like Fe-S₄ centre (DX). This presentation will focus on the structural details that can be perceived from the current model and will include a comparison between DX and the DX-like domain of DFX. The crystal structure of DFX has been determined to 2.5 Å by the MAD method using data measured at the ESRF. The current R-factor value is 26.0 % (R_{free} 34.0 %). Refinement against 1.8 Å data measured at SRS-Daresbury is in progress.

PS02.07.12 CRYSTAL STRUCTURES OF CYTOCHROME c' AND ITS n-BUTYL-ISOCYANIDE-BOUND FORM FROM PURPLE PHOTOTROPHIC BACTERIUM. Tahir H. Tahirov, Shintaro Misaki, Yoshiki Higuchi and Noritake Yasuoka, Department of Life Science, Faculty of Science, Himeji Institute of Technology, Hyogo 678-12, Japan and Terry E. Meyer, and Michael A. Cusanovich, Department of Biochemistry, University of Arizona, Tucson, AZ 85721, U.S.A.

The structures of cytochrome c' from *Rhodobacter capsulatus* (RCCP) strain M110 have been determined by the molecular replacement method. Iron anomalous scattering data were also used to confirm the molecular replacement solution. The structures are refined at 1.72 Å and 2.0 Å resolutions to R values of 15.0% and 16.3% respectively. The RCCP molecule is a dimer and each of the identical 129 residue subunits folds as a four-helical bundle with a covalently bound heme group in the center. This structural motif resembles that of cytochromes c' reported from *Rhodospirillum rubrum* (RMCP), *Rhodospirillum rubrum* (RRCP), and *Chromatium vinosum* (CVCP). However, the architecture of the RCCP dimer, that is, the mode of association of subunits, differs substantially from that of the other cytochromes c'.

In the RCCP dimer, the subunits are roughly parallel to each other and only helix B of each subunit participates in formation of the dimer interface. In RMCP, CVCP and RRCP the subunits cross each other to form an X shape, and two helices, A and B, of each subunit are involved in the interactions across the dimer interface. Structural comparison of four cytochromes c' reveal that they can be divided into two groups. In group 1 cytochromes c', CVCP and RRCP, the amino acid sequences and the folding of subunits are arranged in such a way to allow the formation of a large groove between helices B and C with direct solvent accessibility to the heme sixth ligand position. There is no such groove in group 2 cytochromes c', RMCP and RRCP. This may account in part for the differences in carbon monoxide binding.

An X-ray structure analysis of n-butylisocyanide-bound *Rhodobacter capsulatus* cytochrome c' was also carried out at 2.4 Å resolution. A comparison of the ligand bound structure with that of the native protein reveals that a significant conformational change of amino acid residues occurs in the heme vicinity, accompanied by a rearrangement of the hydrogen bonding pattern. The results suggest that heme puckering in cytochrome c' is a unique mechanical mechanism which may control the dissociation of the dimer to monomers upon ligand binding.

PS02.07.13 CRYSTAL STRUCTURES OF CLOSTRIDIUM PASTEURIANUM RUBREDOXIN MUTANTS V8A AND V8N. Tom J. Brett*, John J. Stezowski*, Kathy M. Selbo*, Charles R. Ross II*, Qiandong Zeng+, Donald M. Kurtz+, and Robert A. Scott+, *Department of Chemistry, University of NebraskaLincoln, Lincoln, NE 68588-0304 +Center for Metalloenzyme Studies, University of Georgia, Athens, GA 30602-2556

Crystal structures of V8A and V8N mutants of *Clostridium pasteurianum* rubredoxin (Rd) have been determined at 1.68Å and 1.82Å resolution data giving final R-values of 14.10% and 14.40%, respectively. The structure determinations were pursued to enhance observations from recent studies in which redox potentials of several Rd mutants were determined¹. In these studies, site-directed mutagenesis was used to study the effects of mutations resulting in surface charge changes near the Fe(Cys)₄ site of Rd. As predicted by simple electrostatic considerations, Rd mutants with positively charged arginine residues in place of neutral surface residues (V8R and L41R) exhibit significant increases in the Fe(II/III) reduction potential. However, significant increases in reduction potential were also seen in mutants where the local negative charge was increased (V8D and L41D), and also when neutral residues were replaced by other neutral residues (V8A and V8N). These findings suggest that reduction potential changes are not dominated by electrostatic effects. Rather, it has been proposed that these reduction potential changes are due to increases in the local dielectric at the Fe(Cys)₄ center. Under this hypothesis, increasing the local dielectric, e.g., by increasing the solvent accessibility of the Fe(Cys)₄ site, would increase the reduction potential. In comparison with the Rd wild-type structure, the V8A and V8N mutants show increased solvent accessibility at one of the cysteine sulfurs (S-Cys42).

1 Q. Zeng, E.T. Smith, D.M. Kurtz, Jr., R.A. Scott. *Inorg. Chim. Acta*, in press.