

MD, USA. ³Johns Hopkins School of Medicine, Baltimore, MD, USA.
E-mail: eviss@jhu.edu

Bacteriophage λ replication initiator, known as λ O protein, is one of the two replication proteins encoded by the bacteriophage's own genome. Its role in initiating bacteriophage DNA replication has been firmly established. Crystals of λ O protein N-terminal fragment, belonging to space group $p2_12_12$ and diffracting to 2.5 Å, were obtained and solved by using anomalous dispersion of selenium atoms. There are two λ O dimers in the asymmetric unit, related to each other by a two-fold non-crystallographic symmetry operator. Genetic data and molecular surface area considerations seem to indicate that the interface(s) between these dimers could be physiologically important and not just a crystallization artifact.

Despite very low homology at the amino-acid sequence level (11% for the structurally aligned regions), the DNA-binding region of λ O displays great similarity to the CAP DNA-binding domain. This structural homology, in combination with genetic data, indicates that DNA-binding mode for λ O protein is similar to the one present in CAP.

Keywords: macromolecular crystallography, replication, protein-DNA interaction

P.04.06.5

Acta Cryst. (2005). A61, C221

Structural Study of the Nucleolin-G-quartet Complex

Christophe Crézé^a, Bruno Rinaldi^b, Philippe Bouvet^b, Richard Haser^a, Patrice Gouet^a, ^aLaboratoire de BioCristallographie, IBCP IFR128, Lyon, France. ^bLaboratoire de Biologie Moléculaire de la Cellule, ENS, Lyon, France. E-mail: c.creze@ibcp.fr

Nucleolin is a very abundant protein of molecular mass 70 kDa that localizes to the nucleolus. It is often used as a cell proliferation marker. Nucleolin interacts with oligonucleotides such as ribosomal RNA [1]. A 40 kDa-fragment of this protein named P40 has been purified to homogeneity. Nucleolin full-length as well as P40 interact with short DNA segments structured in G-quartet [2]. G-quartets are known as potential therapeutic agent against carcinogenesis.

Various G-rich sequences have been structured and tested in interaction with P40 by "Band-shift". Crystallisation trials have been performed for the most stable complexes. Thus, a crystallization condition has been obtained for an expected P40-4(TG4T) complex. Crystals of dimensions 200 x 100 x 20 μm^3 diffract up to 2 Å in house. They belong to space group P1 with unit-cell parameters $a=29\text{Å}$ $b=32\text{Å}$ $c=35\text{Å}$ $\alpha=66^\circ$ $\beta=72^\circ$ $\gamma=83^\circ$. Such small cell volume corresponds to G-quartet only. X-ray data have been collected and the structure has been solved by molecular replacement, using as model the structure of the same G-quartet obtained in another crystal form [3]. Crystallisation trials on other complexes are under way.

[1] Allain F.H., Gilbert D.E., Bouvet P., Feigon J., *J Mol Biol.*, 2000, **303**, 227-41. [2] Hanakahi L.A., Sun H., Maizels N., *J Biol Chem.*, 1999, **274**, 15908-12. [3] Laughlan G., Murchie A.I., Norman D.G., Moore M.H., Moody P.C., Lilley D.M., Luisi B., *Science*, 1994, **265**, 520-4.

Keywords: human proteins, DNA and protein crystallography, anticancer drug structural study

P.04.06.6

Acta Cryst. (2005). A61, C221

Crystal Structure of DNA Polymerase from Bacteriophage M2

Hiroshi Hashimoto^a, Toshiyuki Shimizu^a, Mamoru Sato^a, Hidekazu Koda^b, Yoshiteru Hashimoto^b, Hiroki Higashibata^b, Michihiko Kobayashi^b, ^aGraduate School of Integrated Science, Yokohama City University. ^bGraduate School of Life and Environmental Sciences, The University of Tsukuba, Japan. E-mail: hash@tsurumi.yokohama-cu.ac.jp

DNA polymerases are a group of enzymes that use single-stranded DNA as a template for the synthesis of the complementary DNA strand and are a central player in DNA repair and replication. The many known DNA polymerases can be classified into six families based on phylogenetic relationships: families A, B, C, D, X and Y.

DNA polymerase from bacteriophage M2 (M2 DNA polymerase) is classified into B-family DNA polymerase. M2 DNA polymerase replicates the genome of bacteriophage M2. The enzyme can start to synthesize DNA using a primer-protein as a primer and does not require DNA/RNA primers. We are involved in structural analysis of M2 DNA polymerase to clarify the relationship of its structure and function. Crystals of M2 DNA polymerase were obtained by hanging drop vapor diffusion technique. The crystal belongs to the space group of $P6_122$ and the unit cell parameters are $a = b = 97.0$, $c = 292.1$ Å and $\gamma = 120^\circ$. All X-ray data was collected at SPring-8 BL41XU. Crystal structure of M2 DNA polymerase was determined by multiple isomorphous replacement method with anomalous scatterings using Hg and Pt derivatives at 3.0 Å resolution. Phase calculation was performed with the programs *SOLVE* and *RESOLVE*. The structure was refined with the programs *CNS* and *REFMAC*.

Keywords: protein structures, DNA polymerases, DNA replication

P.04.06.7

Acta Cryst. (2005). A61, C221

The Role of Water in Protein-DNA Complexes from high Resolution X-ray Crystallography

Chin-Yu Chen^{a,c}, Tzu-Ping Ko^a, Andrew H.-J. Wang^{a,b}, ^aInstitute of Biological Chemistry, ^bCore Facility X-ray Crystallography, Academia Sinica, ^cDepartment of Chemistry, Taipei, Taiwan. E-mail: cutecoca@gate.sinica.edu.tw

Sac7d is a small, abundant, non-specific DNA-binding protein of the hyperthermophilic archaeon *Sulfolobus acidocaldarius*. Crystal structures of different Sac7d mutants in complex with DNA have been analyzed at high resolution (1.45 Å – 2.25Å). The high resolution structures of Sac7d-DNA complexes afford good opportunity to study protein-DNA-water interactions in detail. Four recurring well-ordered water molecules are observed in the buried cavity located between protein and DNA surfaces near the intercalation site. They fill up the cavity and enable close packing of protein-DNA interface. These four water molecules are always present at the interface of Sac7d and DNA, although with varying arrangements in different complexes. The buried cavity between Sac7d V26F/M29F and DNA becomes narrower than that in wt-Sac7d due to the phenylalanine/base stacking interaction between Phe26 and G3 base. The four bridging water molecules rearrange in the longitudinal direction along helix axis to match the shapes of the protein-DNA interface.

An important question is how do Sac7d/Sso7d bind to DNA in a sequence-general manner. This water-filled cavity is important in that it allows G-C base pairs to be bound without steric problems due to the additional N2 amino group, thus permitting the proteins its sequence-general binding to DNA.

It is interesting to note that bridging water molecules play an important role in modulating the non-sequence-specific binding of Sac7d by acting as filler, whereas they play an entirely different role as specific linkers between protein and DNA in defining the sequence specificity in the *Trp* repressor-DNA recognition.

Keywords: water, hydrogen bond, protein-DNA interaction

P.04.06.8

Acta Cryst. (2005). A61, C221-C222

Crystal Structure of PriB – a Primosomal DNA Replication Protein of *E. coli*

Chwan-Deng Hsiao, Institute of Molecular Biology, Academia Sinica, Taipei, Taiwan. E-mail: hsiao@gate.sinica.edu.tw

PriB is one of the *Escherichia coli* *phiX*-type primosome proteins which are required for assembly of the primosome, a mobile multi-enzyme complex responsible for the initiation of DNA replication. Here we report the crystal structure of the *Escherichia coli* PriB at 2.1 Å resolution by multi-wavelength anomalous diffraction using a mercury derivative [1]. The polypeptide chain of PriB is structurally similar to that of single-stranded DNA-binding protein (SSB). However, the biological unit of PriB is a dimer, not a homo-tetramer like SSB. Electrophoretic mobility shift assays demonstrated that PriB binds single-stranded DNA and single-stranded RNA with comparable affinity. We also show that PriB binds single-stranded DNA with