

m30.p13**New 1D, 2D and 3D chiral MOF's based on cyclohexane-1,2-diamine**

Olha Sereda, Helen Stoeckli-Evans

*Institut de Chimie, Université de Neuchâtel, Rue Emile-Argand 11, CH-2009 Neuchâtel, Switzerland. E-mail: olha.sereda@unine.ch***Keywords: crystal structure, cyano compounds, paramagnetism**

Cyanide-bridged bimetallic systems, prepared from assembling cyanometallates and transition metal complexes, have been shown to exhibit fascinating structures with interesting magnetic, electrochemical, magneto-optical and zeolite-like properties [1]. The design and synthesis of nanoporous metal-organic frameworks (MOF's), mimicking zeolites, is drawing a great deal of attention due to the application of these materials as catalysts, adsorbents and ion exchangers. Chiral structures are a new target for the lattice architecture in the field of magnetic materials. There are only a very few examples where cyclohexane-1,2-diamine has been incorporated into cyanide-containing complexes resulting in the formation of ferromagnets [2], [3], [4]. We will present some new one-, two- and three-dimensional chiral cyanide-bridged bimetallic MOF's, $[M_1(\text{chxn})_n]_n[M_2(\text{CN})_6]_m$ (where $M_1 = \text{Cu, Ni}$; $M_2 = \text{Ru, Cr, Co}$; $n = 2, 3$; $m = 1, 2$; $\text{chxn} = \text{trans-1S,2S}$ or $1R,2R$ -cyclohexane-1,2-diamine), and their structural features, magnetic and zeolite-like properties will be discussed.

- [1] M. Ohba and K. Okawa, *Coord. Chem. Rev.*, 2000, 198, 313.
 [2] E. Coronado, C.J. Gómez-García, A. Nuez, F.M. Romero, E. Rusanov, H. Stoeckli-Evans, *Inorg. Chem.* 2002, 41, 4615.
 [3] E. Coronado, C. Giménez-Saiz, J.M. Martínez-Agudo, A. Nuez, F.M. Romero, H. Stoeckli-Evans, *Polyhedron* 2003, 22, 2435.
 [4] E. Coronado, C. Giménez-Saiz, A. Nuez, V. Sanchez, F.M. Romero, *Eur.J.Inorg.Chem.* 2003, 4289.

m30.p14**Atomic and domain structure of the low temperature phase of barium metagermanate**Daniel Maria Töbrens,^a Volker Kahlenberg,^a Christian Gspan^b, Gerald Kothleitner^b*^aInstitute of Mineralogy and Petrography, University of Innsbruck, Austria, and ^bResearch Institute for Electron Microscopy, Graz, Austria. E-mail: daniel.toebbens@uibk.ac.at***Keywords: domain structure, layered structures, electron microscopy**

The crystal structure of the low temperature form of barium metagermanate (BaGeO_3) has been determined from laboratory X-ray powder diffraction data collected at 25.5(5) °C. The compound is monoclinic with space group $C2/c$ and unit cell parameters of $a = 13.1877(2) \text{ \AA}$, $b = 7.6222(1) \text{ \AA}$, $c = 11.7190(1) \text{ \AA}$, $\beta = 112.295(1)^\circ$, $V = 1089.9(1) \text{ \AA}^3$ and $Z = 12$. The structure was found to consist of alternating layers of barium atoms and three-membered $[\text{GeO}_3]_3$ -rings showing a twofold positional disorder. Apart from the disorder the structure is very similar to the one of SrGeO_3 [1]; both are six-layer structures build from the same type of layers. The same structure type also appears in pseudo-wollastonite, forming four- or six-layer polytypes [2]. Careful studies of possible structural variations using simulated annealing [3] did not reveal any deviation from symmetry $C2/c$.

The disorder of the ternary rings can be summarized as follows: ternary $[\text{GeO}_3]_3$ -rings appear in only one out of two possible orientations and occupy only 2/3 of the positions compatible with this orientation. Following [2], for six-layered structures of this type there is only one distinct polytype with the inherent symmetry $C2/c$. There are, however, two non-congruent enantiomorphs of this structure. The disorder observed in BaGeO_3 can be described as the simultaneous appearance of both possibilities.

Electron microscopy reveals in SAED patterns the existence of twins; the twin law can be described as a mirror plane normal to \mathbf{a} . Diffuse scattering within the lattice rods parallel \mathbf{c}^* indicates disorder within the layers of the structure. HRTEM shows fully ordered domains, which are separated by narrow disordered regions; the domain borders observed are normal to \mathbf{c}^* . Larger disordered regions are observed as well.

Based on these observations the disorder in the structure can be addressed as stacking fault disorder, a behaviour also found in polytypes of pseudo-wollastonite [2]. The stacking, however, is not random but can be rationalized by a twinning mechanism mapping the enantiomorphic polytypes onto each other.

- [1] Nishi, F. (1997). *Acta Cryst.* C53, 399-401.
 [2] Yamanaka, T., Mori, H. (1981). *Acta Cryst.* B37, 1010-1017.
 [3] Favre-Nicolin, V., Černý, R. (2004). *Z. Krist.* 219, 847-856.