

of crystals. As important as the space groups themselves are their subgroups. Here, we review how they have improved our understanding of the space groups themselves, how they have proved useful in solving various problems of crystallography and how they have been used to define new groups. Furthermore we give an outlook of what still has to be done. One of the most important subgroups is the group T of lattice translations, which we can use to construct the quotient group $P=G/T$, which is better known under the name point group. We start our presentation by reviewing the work of E. Ascher and A. Janner, who have shown how to view space groups as extensions of P by T. We pass on to more general subgroups and discuss coset and double coset decompositions and how they have been applied to phase transitions and twins in crystals. Closely connected to coset decompositions are colour groups, which have been studied among others by J.J. Burckhardt. A special kind of colorings are Bravais colorings, which leads us to similar sublattices and the group of similarity rotations, which is a supergroup of the point group. In between these two groups is the group of coincidence rotations, thus we arrive at coincidence site lattices (CSLs) and grain boundaries and return to twins. Last but not least we discuss the connections between colorings and CSLs, thus arriving at current research topics.

Keywords: space groups, colourings, coincidence site lattices

MS.89.4

Acta Cryst. (2008). A64, C150

J.J. Burckhardt's contributions to crystallography

Bernd Souvignier

Radboud University Nijmegen, Institute for Mathematics, Astrophysics and Particle Physics, Postbus 9010, Nijmegen, Gelderland, 6500 GL, The Netherlands, E-mail: souvi@math.ru.nl

Johan Jakob Burckhardt, who died in 2006 at the age of 103, was deeply interested in crystallography all of his life and made invaluable contributions to the mathematical foundations of space group theory. After first studying mathematics in Basel and Munich, he was so impressed by Andreas Speiser's *Die Theorie der Gruppen von endlicher Ordnung* that he moved to Zurich to study with Speiser the year after its publication. There he also attended lectures of Paul Niggli and (explicitly recommended by Speiser) Leonhard Weber. On the basis of his acquired mathematical background, Burckhardt developed a new derivation of the 230 space group types which placed the earlier work of Schoenflies and Fedorov in the modern algebraic context of cohomology theory. His approach via Frobenius congruences in particular freed space group theory from dimensional constraints. For example, he determined the space groups for cyclic, symmetric and alternating permutation groups (in their natural permutation representation) for arbitrary degree. Burckhardt's work on space groups culminated in the publication of *Die Bewegungsgruppen der Kristallographie* in 1947 which is still a standard reference in mathematical crystallography. Together with Bartel Leenert van der Waerden he published a short but crystal clear article about colour symmetry which clarified the somewhat obscured ideas around that topic and influenced a generation of researchers. Burckhardt also became a historian of mathematical crystallography, and helped to illuminate its long and multi-lingual, multi-disciplinary path. Perhaps most important of all, Burckhardt corresponded widely with mathematicians and crystallographers all over the world, helping to create the community that this symposium implicitly celebrates.

Keywords: J.J. Burckhardt, space group theory, colour symmetry

MS.89.5

Acta Cryst. (2008). A64, C150

Heterogeneous cylinder packing: Space group on periodic structures with $\langle 110 \rangle$ six directions

Yoshinori Teshima^{1,5}, Takeo Matsumoto², Yoshinori Watanabe³, Tohru Ogawa^{4,5}

¹National Institute of Advanced Industrial Science and Technology (AIST), 1-2 Namiki, Tsukuba, Ibaraki, 305-8564, Japan, ²Kanazawa University (Professor emeritus), 2-27 Tsuchishimizu, Kanazawa, 920-0955, Japan, ³University of Tsukuba, Tennodai, Tsukuba, Ibaraki, 305-8577, Japan, ⁴University of Tsukuba (Professor emeritus), 3-197 Edogawadai-higashi, Nagareyama, 270-0111, Japan, ⁵Interdisciplinary Institute of Science and Technology and Art (ISTA), 2-5-28 Kitahara, Asaka, Saitama, 351-0036, Japan, E-mail: yoshinori.teshima@aist.go.jp

Several cylinder packing structures were found by authors around 1995. We gave equations of structures and calculated packing density. But we have left their space group undetermined for many years. In this talk, we report space group on periodic structures of cylinder packing with $\langle 110 \rangle$ six directions. All the cylinders are congruent and the length of the cylinders is infinite and their directions are restricted to only six directions of $\langle 110 \rangle$. Each cylinder is fixed by cylinders of other directions, so that the whole structure sustains mechanical stability. The space group tell us each structure is not homogeneous. But they have ingenious feature: parallel cylinders in six directions form equivalent two-dimensional rhombic lattice respectively.



Keywords: cylinder packing, rod packing, space group

MS.90.1

Acta Cryst. (2008). A64, C150-151

Life sciences at Diamond Light Source and prospects with new light sources

Louise N. Johnson

Diamond Light Source, Harwell Science and Innovation Campus, Didcot, Oxon, OX11 0DE, UK, E-mail: louise.johnson@diamond.ac.uk

Diamond Light Source was commissioned for users in 2007. The machine consists of a 100 MeV linear accelerator, a 3 GeV booster synchrotron and a 3 GeV storage ring, together with interconnecting beam transfer lines. The storage ring (circumference 562 m) based on 24 cells is run at 300 mA. The electron beam emittance is 2.7 nm rad with bunch length (FWHM) 25.6 ps. The accelerating voltage is provided by two superconducting cavities based on inductive output tubes (IOTs) each capable of delivering a maximum total output power of 300 kW. In Phase I, 7 insertion devices were installed: 5 in-vacuum undulators, one variable polarisation device and one superconducting wiggler. At the end of Phase II (2011) Diamond will have 22 experimental stations for research in both the life sciences and physical sciences, 7 beam lines in Phase I (completed 2007) and 15 beam lines in Phase II. This talk will consider the impact of synchrotron radiation on life sciences research and will attempt to assess the future demand for new X-ray sources. Diamond will have 8 beam lines dedicated to life sciences research: 5 macromolecular crystallography beam lines, a non-crystalline diffraction and solution scattering beam line, a circular dichroism beam line and an infrared microspectroscopy beam line. These will provide for exciting and

innovative biological research. However, there is a gap in the life sciences between the detailed molecular information that can be obtained with the above methods and the precise location of such complexes in functioning cells. There is scope for new imaging technologies. At Diamond we are exploring new beam lines that exploit coherent diffraction imaging and soft X-ray microscopy and the possibilities of new light sources.

Keywords: diamond light source, life sciences research, macromolecular complexes

MS.90.2

Acta Cryst. (2008). A64, C151

A compact X-ray free electron laser at SPring-8

Tetsuya Ishikawa

RIKEN, SPring-8 Center, Kouto 1-1-1, Sayo, Hyogo, 679-5148, Japan, E-mail: ishikawa@spring8.or.jp

Japan launched a construction project of x-ray free electron laser (XFEL) in 2006, as one of five Key Technologies of National Importance, to complete in 2010 fiscal year. The XFEL emits coherent hard x-rays which will be extremely useful for exploring the nano-world. The operation principle of the Japanese XFEL at the initial phase will be self-amplified spontaneous emission (SASE), as the precedent two projects in the U.S. and Europe. The XFEL will be composed of a linear accelerator and a long undulator in which electrons interact with photons to form micro-bunches with the period of emitted radiation. The micro-bunched electrons radiate coherent photons when they run through the undulator. The unique feature of the Japanese XFEL is the use of in-vacuum undulator. This makes the magnetic period of the undulator shorter than that of conventional out-of-vacuum undulators, leading to the reduction of the electron energy to produce certain energy of x-ray photons. Therefore, the length of the linear accelerator will be shorter, and the whole facility will be more compact. However, lower electron energy requires higher beam quality to produce lasing than is realized by a conventional laser-RF gun system. We have developed a new injector system, instead, which was fully proven in the operation of a prototype ultraviolet FEL. The present status of the project will be reported together with scientific programs using the XFEL, synergetic use of XFEL and SPring-8, and the second phase program for seeded XFEL. Some recent result from the prototype FEL will be introduced as well.

Keywords: X-ray free electron laser, coherent radiation, accelerator based x-ray source

MS.90.3

Acta Cryst. (2008). A64, C151


Status of the Energy Recovery Linac (ERL) project at Cornell University

Sol M Gruner

Cornell University, Physics Dept. & CHESS, 162 Clark Hall, Ithaca, NY, 14853-2501, USA, E-mail: smg26@cornell.edu

The status of the Energy Recovery Linac (ERL) project at Cornell University is summarized. ERLs are being developed as next generation synchrotron light sources. The fundamental x-ray beam properties from storage ring sources, such as the source size, spectral brightness, and pulse duration are limited by the dynamic equilibrium characteristic of the magnetic lattice that is the storage ring. Advances

in laser-driven photoelectron sources and superconducting linacs allow the acceleration of electron bunches with superior properties for synchrotron radiation, and superconducting linac energy recovery allows the development of an x-ray ring without the need for electron storage. Relevant properties include x-ray beams of extraordinary spectral brightness and small source size, with concomitant high transverse coherence, sub-picosecond x-ray pulse durations, and flexibility of operation. ERLs are capable of hosting practically all experiments now being carried out at storage rings while also enabling new types of experiments. Progress is reported on development of the required high-spectral brightness photo-injector and superconducting linac. X-ray applications are discussed.



Keywords: X-ray synchrotron radiation, technology, instrument development

MS.90.4

Acta Cryst. (2008). A64, C151-152

Electron storage ring based tabletop light source named MIRRORCLE for protein crystallography

Hironari Yamada^{1,2}, Masaki Morita², Takayasu Hanashima², Ahsa Moon¹, Tuyoshi Shirai³

¹Ritsumeikan University, Synchrotron Light Life Science Center, 1-1-1 Nojihigashi, Kusatsu, Shiga, 525-8577, Japan, ²Photon Production Laboratory Co. Ltd., ³Nagahama Institute of Bio Science and Technology, E-mail: hironari@se.ritsume.ac.jp

Light source based on a 20 MeV storage ring has been commissioned. The average stored beam current was more than 2A at 400Hz beam injection. We have observed an extremely short 15 ms damping time, a small 3 mm diameter beam size, and a long 1 min lifetime to this low energy and 15cm orbit radius electron storage ring. We generate X-rays by a 10 micron size target placed in this storage ring. The observed X-ray density is more than that of 1kW X-ray tube, but the spectrum is polychromatic. The X-ray energy extends up to 20 MeV and the total power is 100W when an accelerator power is applied. We have commissioned a crystal monochromator beam line. A special configuration enabled extracting tunable monochromatic X-rays onto the fixed sample position. The 10 to 30 keV range x-rays is extracted. We have carried out EXAFS and found that the energy resolution is satisfactory for the protein crystallography (see fig.). Due to a background radiation around the beam line and a poor performance of CCD camera, diffraction pattern is

