

## MS5-P2 Electron Nanocrystallography for Organic and Macromolecular Structure Determination

Tim Gruene<sup>1</sup>, Eric van Genderen<sup>1</sup>

1. Paul Scherrer Institute

email: tim.gruene@psi.ch

Electron crystallography has a long tradition in material science and also in 2D crystallography. Electrons interact much more strongly with matter and they cause about three orders of magnitude less radiation damage per diffracted quantum than X-rays. Crystals with a few to a few hundred nanometre thickness can be used for structure determination. These features have important applications for organic and (also macromolecular) structure determination: Often compounds, that represent themselves as powders to X-ray sources, consist of single nanocrystals suitable for structure determination with electron diffraction. In addition, crystal defects like disorder increase with crystal size so that data from nanocrystals can produce better quality models.

We recently exploited a quantum hybrid electron detector, Timepix, to determine the crystal structure of two organic compounds from single crystals at room temperature [1]. Here I will present our progress since this publication. We have improved the experimental procedure and developed methods to reduce systematic errors to such an extent that the data quality allows structure resolution with the default options of shelxt. Both structure solution and refinement can be carried out simply with well established programs like shelxl/shelxle. My presentation will include the effects of dynamic scattering that is often used as argument why electron diffraction was not practical to solve the structures of organic or macromolecular compounds.

Currently electron nanocrystallography is at a stage comparable with X-ray crystallography 3-4 decades ago. As we exploit a vast pool of experience from X-ray crystallography, electron nanocrystallography will soon become a new standard tool for biomolecular structure determination covering those systems that fail X-ray diffraction.

[1] van Genderen et al., *Acta Cryst* (2016), A72, 235-24

**Keywords:** Electron Nanocrystallography, Dynamic Scattering

## MS5-P3 Targeting Epigenetic Vulnerabilities of Cancer Cells by Exploiting Chromatin Structure and Chemistry

Zenita Adhiksan<sup>1</sup>, Gabriela E. Davey<sup>1</sup>, Zhujun Ma<sup>1</sup>, Giulia Palermo<sup>2</sup>, Tina Riedel<sup>2</sup>, Benjamin S. Murray<sup>3</sup>, Alexey A. Nazarov<sup>4</sup>, Christian G. Hartinger<sup>5</sup>, Ursula Röthlisberger<sup>2</sup>, Paul J. Dyson<sup>2</sup>, Curt A. Davey<sup>1</sup>

1. School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive, Singapore 637551

2. Institut des Sciences et Ingénierie Chimiques, Ecole Polytechnique Fédérale de Lausanne (EPFL), CH-1015 Lausanne, Switzerland

3. Department of Chemistry, University of Hull, Cottingham Road, Hull, HU6 7RX, United Kingdom

4. Moscow State University, Department of Chemistry, Leninskoye, 119991 Moscow, Russia

5. School of Chemical Sciences, University of Auckland, Private Bag 92019, Auckland 1142, New Zealand

email: zenita@ntu.edu.sg

Pronounced differences in the gene expression profiles between healthy and cancerous cells coincide with extensive epigenetic distinctions in chromatin, and yet the therapeutic landscape of chromatin is largely unexplored. We thus hypothesize that compounds capable of recognizing specific features of chromatin could enable targeting weak points of cancer cells [1-3]. Recent years have witnessed a resurgence of interest in developing new metal-based anticancer agents that are more effective and have fewer side effects, with focus shifting towards metals that are alternatives to the traditional platinum. We have so far characterized the nucleosome binding activity of over 50 different non-Pt drugs and therapeutic candidates, including various Ru-, Os-, Rh- and Au-based compounds. We observe that only one specific class of Ru agent preferentially forms DNA adducts, while the rest have an apparent preference for reaction at histone protein sites. In conjunction with studies on Pt compounds, this has revealed principles for selective targeting of different protein and DNA sites within chromatin and corresponding relationships to cytotoxicity and impact on cancer cell function. Moreover, newly designed binuclear antitumor compounds target a key regulatory site on the nucleosome, which alters chromatin structure in vitro and appears to kill cancer cells by interfering with chromatin dynamics. Beyond this, we have also discovered that two unrelated histone-targeting agents, in combination, display synergistic antitumor activity, and we can link this to a novel mechanical mechanism involving allosteric modulation in the nucleosome.

[1] C.A. Davey. 2015. Exposure to Metals Can Be Therapeutic. *Chimia*. **69**: 125-130.

[2] B. Wu\*, M.S. Ong\*, M. Groessl, Z. Adhiksan, C.G. Hartinger, P.J. Dyson & C.A. Davey. 2011. A Ruthenium Antimetastasis Agent Forms Specific Histone Protein Adducts in the Nucleosome Core. *Chem. Eur. J.* **17**: 3562-3566.

[3] Z. Adhiksan\*, G.E. Davey\*, P. Campomanes\*, M. Groessl, C.M. Clavel, H. Yu, A.A. Nazarov, H.F. Yeo, W.H. Ang, P. Dröge, U. Röthlisberger, P.J. Dyson & C.A. Davey. 2014. Ligand Substitutions between Ruthenium-Cymene Compounds Can Control Protein versus DNA Targeting and Anticancer Activity. *Nat. Commun.* **5**: 3462.

**Funding:** Ministry of Health National Medical Research Council (grant NMRC/1312/2011); Ministry of Education Academic Research Fund Tier 3 Programme (grant MOE2012-T3-1-001).

E-mail for correspondence: davey@ntu.edu.sg

**Keywords:** chromatin, nucleosome, metal-based compound, anticancer drug

**MS5-P4** Structure analysis of arylpiperazine derivatives displaying affinity towards 5-HT<sub>1A</sub> and 5-HT<sub>7</sub> receptors

Mateusz Jabłoński<sup>1</sup>, Katarzyna Rzęsikowska<sup>1</sup>, Justyna Kalinowska-Thluścik<sup>1</sup>

<sup>1</sup>. Faculty of Chemistry, Jagiellonian University, Ingardena 3, 30 - 060 Kraków, Poland

email: jablonskikrystalografia@gmail.com

Serotonin (5-HT, 5-hydroxytryptamine) is one of the evolutionary oldest biogenic monoamines, which regulates various physiological functions in the human body. The serotonin receptor family was divided into 7 main classes among which, depending on the classification, at least 14 distinct classes may be distinguished. Six out of seven main classes belong to the G protein-coupled receptors (GPCRs) family. The member of the last main class, 5-HT<sub>7</sub> receptor, was found to be involved in regulation of temperature, circadian rhythm and proper functioning of memory. The lack of an accurate experimental structural data for most 5-HT receptor family members is a challenge in the process of designing selective and safe agonists and antagonists for individual classes. The selectivity problem is particularly important and visible in designing selective 5-HT<sub>1A</sub> and 5-HT<sub>7</sub> ligands, because the 5-HT<sub>7</sub> receptor shares a high sequence similarity to 5-HT<sub>1A</sub> receptor, especially in the binding pocket region.

Arylpiperazine is one of the most universal fragments found in ligands acting on various central nervous system receptors. Arylpiperazines are also a noteworthy scaffold component in 5-HT receptor ligands. Depending on the aryl part and its potential substituents, they can demonstrate diverse intrinsic activity and manifest different selectivity among the distinct 5-HT receptor classes. Substituents may variously affect the electronic features of the aromatic ring, which may result in different preferential mutual orientation of the aryl ring in regards with the piperazine moiety.

In this work, structures of selected long chain arylpiperazine derivatives are presented. Each compound contains a theophylline moiety connected to the piperazine with an alkyl linker. The crystal structure analysis is accompanied by molecular docking and structure-affinity relationship investigation.

**Keywords:** long chain arylpiperazines, serotonin receptors, structure-activity relationship, molecular docking