

Acknowledgments:

Polish National Science Centre MAESTRO grant decision number DEC2012/04/A/ST5/00609, DSM grant No. 501-D112-86-DSM-115 100 are acknowledged.

References:

Coppens, P. (1997). X-Ray Charge Densities and Chemical Bonding; International Union of Crystallography.
 Dittrich, B.; Weber, M.; Kalinowski, R.; Grabowsky, S.; Hübschle, C. B.; Luger, P. (2009). *Acta Crystallogr.* 65 (6), 749–756.
 Jarzemska, K. N.; Dominiak, P. M. (2012). *Acta Crystallogr. A*. 68 (Pt 1), 139–147.

Keywords: 9-aminoacridine hemihydrate, charge density, TAAM refinement

MS21-P02

Preparation of non-covalent organic frameworks using dodecasubstituted porphyrins


Keith Flanagan¹, Mathias O. Senge¹

1. School of Chemistry, Trinity College Dublin the University of Dublin, Dublin, Ireland

email: kflanaga@tcd.ie

Nowadays the use of porphyrins to create highly porous, sponge-like frameworks is a well-studied field, established by the seminal work of Byrn *et al.* coining the phrase “porphyrin sponge” in 1991.^[1] This initial study used a porphyrin host, trapping a variety of guests within the crystal lattice. Since then, many contributions have focused on the manipulation of planar porphyrins containing a variety of functional groups (namely carboxylic acids or halogens) to obtain directly linked porphyrin frameworks.^[2] However, as of yet, there are no specific examples of the use of highly distorted porphyrins^[3] being used to obtain such highly porous materials. In this work, we have synthesized a series of 5,10,15,20-tetraaryl-2,3,7,8,12,13,17,18-octaethylporphyrins with alternating halogen, alkyl, and aryl substitution patterns to probe the effects of non-planarity on the formation of non-covalent organic frameworks.

The target porphyrins were synthesized according to a modified Lindsey condensation reaction followed by metal(II) insertion. To investigate substituent interactions, modified aldehydes were used. For halogen effects, a family of derivatives with F, Cl, Br or I in either the *ortho*- or *para*-position of the meso-aryl residues were selected. Hydrogen bonding interactions were examined by incorporating a nitrile group. Steric effects were examined through the incorporation of a benzyloxy group at either the *meta*- or *para*-positions. Finally, investigations were conducted into the use of arm-extended porphyrins using a triazole linker group to further extend their functionality. Solid-state structures were determined *via* small molecule X-ray crystallography to elucidate the major distortion modes and to characterize any non-covalent interactions. These were then compared to their planar derivatives to determine effects the substitution patterns may have on the formation of non-covalent organic frameworks.



References:

- [1] Byrn, M. P. *et al.* (1991). *J. Am. Chem. Soc.*, 113, 6549–6557.
 [2] Goldberg, I. (2008). *Cryst. Eng. Comm.*, 10, 637–645.
 [3] Senge, M. O. *et al.* (2015). *Chem. Commun.*, 51, 17031–17063.

Keywords: Porphyrin, Halogens, Macrocycles