

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

catena-Poly[[diaqua[(4-tolylsulfanyl)-acetato-κO]cadmium(II)]-μ-4,4'-bipyridine-κ²N:N']

Wen-Xuan Cai,‡ Xiao-Yong Zheng,‡ Xiao-Hong Geng and Yun-Long Feng*

Zhejiang Key Laboratory for Reactive Chemistry on Solid Surfaces, Institute of Physical Chemistry, Zhejiang Normal University, Jinhua, Zhejiang 321004, People's Republic of China


Correspondence e-mail: sky37@zjnu.cn

Received 11 July 2008; accepted 1 August 2008

 Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.032; wR factor = 0.113; data-to-parameter ratio = 17.1.

The title complex, $[\text{Cd}(\text{C}_9\text{H}_9\text{O}_2\text{S})_2(\text{C}_{10}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_2]_n$, has a linear chain structure. The central Cd^{II} ion is in a slightly distorted octahedral environment, coordinated by two aqua ligands, two (4-tolylsulfanyl)acetate ligands and two bridging 4,4'-bipyridine ligands. The Cd^{II} ion lies on a twofold rotation axis. Intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds connect adjacent chains, forming a layer structure. An intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond is also present.

Related literature

 For related literature, see: Lin *et al.* (2006); Zheng *et al.* (2006).


Experimental

Crystal data

$[\text{Cd}(\text{C}_9\text{H}_9\text{O}_2\text{S})_2(\text{C}_{10}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_2]$
 $M_r = 667.09$
 Monoclinic, $C2/c$
 $a = 21.659$ (4) Å
 $b = 11.590$ (2) Å
 $c = 11.137$ (2) Å
 $\beta = 93.88$ (3)°

$V = 2789.3$ (9) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.98$ mm⁻¹
 $T = 296$ (2) K
 $0.40 \times 0.35 \times 0.17$ mm

Data collection

Bruker APEXII area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.68$, $T_{\text{max}} = 0.85$

12185 measured reflections
 3154 independent reflections
 2937 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.017$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.112$
 $S = 1.12$
 3154 reflections
 184 parameters
 3 restraints

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\text{max}} = 0.60$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.37$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1W}-\text{H1WA}\cdots\text{O2}$	0.82	1.94	2.667 (4)	148
$\text{O1W}-\text{H1WB}\cdots\text{O1}^{\dagger}$	0.805 (17)	2.04 (2)	2.782 (3)	154 (4)

 Symmetry code: (i) $x, -y, z + \frac{1}{2}$.

Data collection: APEX2 (Bruker, 2006); cell refinement: SAINT (Bruker, 2006); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: AT2595).

References

- Bruker (2006). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Lin, H., Su, H. & Feng, Y. (2006). *Acta Cryst.* **E62**, m747–m749.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Zheng, X.-Y., Su, H. & Feng, Y.-L. (2006). *Acta Cryst.* **E62**, m1393–m1394.

‡ Current address: Wenzhou Medical College Pharmacy School, Wenzhou Medical College, Wenzhou, Zhejiang 325035, People's Republic of China.

supporting information

Acta Cryst. (2008). E64, m1137 [doi:10.1107/S1600536808024720]

catena-Poly[[diaqua[(4-tolylsulfanyl)acetato- κ O]cadmium(II)]- μ -4,4'-bipyridine- κ^2 N:N']

Wen-Xuan Cai, Xiao-Yong Zheng, Xiao-Hong Geng and Yun-Long Feng

S1. Comment

The title compound, (I), is isostructural with the Ni^{II} (Lin *et al.*, 2006) and Co^{II} analogues (Zheng *et al.*, 2006). The structure of (I) (Fig. 1) consists of linear chains formed through 4,4'-bipy ligands linking six-coordinated Cd^{II} ions which lie on twofold rotation axes. Intermolecular O—H \cdots O hydrogen bonds link neighboring chains to form a two-dimensional network. It is notable that these chains are arranged alternately and the 4-tolylsulfanyl groups are almost coplanar. There is no significant π - π interactions between the planes of adjacent chains with centroid-centroid distance of 6.19 (1) Å and plane-to-plane distance of 3.64 (1) Å.

S2. Experimental

CdSO₄·8/3H₂O (0.128 g, 0.5 mmol), (4-tolylsulfanyl)acetic acid (0.091 g, 0.5 mmol), 4,4'-bipy (0.039 g, 0.25 mmol) and H₂O (18 ml) were sealed in a 25 ml stainless-steel reactor with a Teflon-lined stainless steel reactor and the solution was heated at 433 K for 72 h and then cooled to room temperature over a period of 72 h. Colourless crystals suitable for X-ray analysis were obtained.

S3. Refinement

The methyl groups were allowed to rotate to fit the electron density [C—H = 0.96 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$]; the other H atoms were positioned geometrically [aromatic C—H = 0.93 Å and aliphatic C—H = 0.97 Å, $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$]. Water H atom H1WA was positioned geometrically, with O—H = 0.82 Å, and the other water H atoms H1WB was located from a difference Fourier map, and they were refined with distance restraints of O—H = 0.85 (2) Å and H \cdots H = 1.30 (2) Å; their displacement parameters were set to $1.5U_{\text{eq}}(\text{O})$.




Figure 1

A view of part of the title structure, showing 30% probability displacement ellipsoids. [Symmetry codes: (a) $-x, y, -z+1/2$; (b) $x, y+1, z$; (c) $x, y-1, z$.]




Figure 2

The chain structure of the title compound. All H atoms have been omitted for clarity.

catena-Poly[[[diaqua[(4-tolylsulfonyl)acetato- κ O]cadmium(II)]- μ -4,4'-bipyridine- κ^2 N:N']

Crystal data

$[\text{Cd}(\text{C}_9\text{H}_9\text{O}_2\text{S})_2(\text{C}_{10}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_2]$

$M_r = 667.09$

Monoclinic, $C2/c$

Hall symbol: $-C 2yc$

$a = 21.659 (4) \text{ \AA}$

$b = 11.590 (2) \text{ \AA}$

$c = 11.137 (2) \text{ \AA}$

$\beta = 93.88 (3)^\circ$

$V = 2789.3 (9) \text{ \AA}^3$

$Z = 4$

$F(000) = 1360$

$D_x = 1.589 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 7745 reflections

$\theta = 2.7\text{--}27.5^\circ$

$\mu = 0.98 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Block, colourless

$0.40 \times 0.35 \times 0.17 \text{ mm}$

Data collection

Bruker APEXII area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.68, T_{\max} = 0.85$

12185 measured reflections

3154 independent reflections

2937 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.017$

$\theta_{\max} = 27.5^\circ, \theta_{\min} = 2.7^\circ$

$h = -27 \rightarrow 27$

$k = -14 \rightarrow 14$

$l = -14 \rightarrow 14$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.112$
 $S = 1.12$
 3154 reflections
 184 parameters
 3 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0658P)^2 + 5.8321P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.60 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.37 \text{ e } \text{Å}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cd1	0.0000	0.05131 (2)	0.2500	0.03198 (12)
S1	0.14880 (4)	0.21253 (9)	-0.05125 (7)	0.0514 (2)
O1	0.08144 (10)	0.05635 (16)	0.1325 (2)	0.0353 (5)
O1W	0.06445 (12)	0.0644 (2)	0.4169 (2)	0.0415 (5)
H1WA	0.0990	0.0836	0.3982	0.062*
H1WB	0.0700 (14)	0.012 (3)	0.464 (3)	0.047 (11)*
O2	0.15198 (12)	0.1218 (3)	0.2705 (2)	0.0632 (8)
N1	0.0000	-0.1422 (3)	0.2500	0.0290 (6)
N2	0.0000	-0.7537 (3)	0.2500	0.0303 (7)
C1	0.16190 (13)	0.3534 (3)	0.0059 (3)	0.0422 (7)
C2	0.19048 (16)	0.3817 (4)	0.1177 (3)	0.0506 (9)
H2A	0.2025	0.3234	0.1717	0.061*
C3	0.20102 (17)	0.4945 (4)	0.1488 (3)	0.0544 (9)
H3A	0.2210	0.5109	0.2234	0.065*
C4	0.18302 (16)	0.5854 (4)	0.0730 (3)	0.0520 (8)
C5	0.15375 (18)	0.5566 (3)	-0.0382 (3)	0.0508 (9)
H5A	0.1407	0.6152	-0.0910	0.061*
C6	0.14372 (16)	0.4440 (3)	-0.0715 (3)	0.0469 (9)
H6A	0.1245	0.4275	-0.1468	0.056*
C7	0.1952 (2)	0.7075 (5)	0.1109 (4)	0.0714 (12)
H7A	0.2389	0.7186	0.1271	0.107*
H7B	0.1742	0.7236	0.1822	0.107*
H7C	0.1803	0.7586	0.0476	0.107*
C8	0.17850 (15)	0.1165 (4)	0.0670 (3)	0.0495 (8)

H8A	0.1879	0.0426	0.0316	0.059*
H8B	0.2169	0.1481	0.1029	0.059*
C9	0.13410 (13)	0.0967 (3)	0.1664 (3)	0.0385 (6)
C10	-0.02550 (18)	-0.2021 (3)	0.3350 (3)	0.0470 (8)
H10A	-0.0435	-0.1619	0.3959	0.056*
C11	-0.02676 (18)	-0.3207 (3)	0.3379 (3)	0.0441 (8)
H11A	-0.0457	-0.3584	0.3992	0.053*
C12	0.0000	-0.3838 (3)	0.2500	0.0259 (7)
C13	0.0000	-0.5111 (3)	0.2500	0.0250 (7)
C14	-0.02533 (15)	-0.5744 (3)	0.3404 (3)	0.0357 (6)
H14A	-0.0432	-0.5363	0.4029	0.043*
C15	-0.02423 (15)	-0.6931 (3)	0.3382 (3)	0.0365 (6)
H15A	-0.0410	-0.7331	0.4006	0.044*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cd1	0.03971 (19)	0.02551 (18)	0.03160 (18)	0.000	0.00893 (12)	0.000
S1	0.0526 (5)	0.0682 (6)	0.0343 (4)	-0.0107 (4)	0.0090 (3)	0.0052 (4)
O1	0.0353 (11)	0.0355 (11)	0.0362 (11)	-0.0019 (7)	0.0104 (9)	-0.0040 (8)
O1W	0.0525 (13)	0.0448 (12)	0.0281 (10)	0.0011 (10)	0.0085 (9)	0.0047 (9)
O2	0.0444 (13)	0.107 (2)	0.0385 (13)	-0.0222 (15)	0.0024 (10)	0.0066 (14)
N1	0.0354 (16)	0.0245 (15)	0.0280 (15)	0.000	0.0076 (13)	0.000
N2	0.0364 (16)	0.0283 (16)	0.0272 (15)	0.000	0.0089 (13)	0.000
C1	0.0259 (13)	0.071 (2)	0.0310 (14)	-0.0051 (13)	0.0104 (11)	0.0042 (14)
C2	0.0409 (16)	0.080 (3)	0.0311 (15)	-0.0024 (17)	0.0043 (12)	0.0064 (16)
C3	0.0416 (17)	0.088 (3)	0.0336 (16)	-0.0069 (19)	0.0047 (13)	-0.0062 (18)
C4	0.0392 (16)	0.074 (2)	0.0447 (18)	-0.0076 (17)	0.0169 (14)	-0.0093 (18)
C5	0.0455 (19)	0.068 (3)	0.0399 (18)	0.0016 (15)	0.0091 (15)	0.0059 (15)
C6	0.0368 (16)	0.074 (3)	0.0302 (15)	-0.0021 (14)	0.0025 (12)	0.0062 (14)
C7	0.070 (3)	0.080 (3)	0.066 (3)	-0.010 (2)	0.023 (2)	-0.014 (2)
C8	0.0342 (15)	0.067 (2)	0.0492 (18)	0.0037 (15)	0.0146 (13)	0.0072 (16)
C9	0.0321 (13)	0.0426 (16)	0.0417 (16)	0.0027 (12)	0.0084 (12)	0.0068 (13)
C10	0.073 (2)	0.0307 (15)	0.0411 (16)	0.0028 (15)	0.0315 (16)	-0.0034 (13)
C11	0.069 (2)	0.0292 (14)	0.0374 (15)	0.0012 (14)	0.0306 (15)	0.0021 (12)
C12	0.0284 (16)	0.0253 (17)	0.0245 (16)	0.000	0.0048 (13)	0.000
C13	0.0260 (16)	0.0236 (17)	0.0259 (16)	0.000	0.0047 (13)	0.000
C14	0.0510 (17)	0.0298 (13)	0.0289 (13)	-0.0056 (12)	0.0207 (12)	-0.0053 (11)
C15	0.0506 (16)	0.0293 (14)	0.0320 (13)	-0.0068 (12)	0.0198 (12)	-0.0030 (11)

Geometric parameters (Å, °)

Cd1—N1	2.243 (3)	C4—C5	1.393 (5)
Cd1—O1W	2.253 (2)	C4—C7	1.495 (6)
Cd1—O1W ⁱ	2.253 (2)	C5—C6	1.371 (5)
Cd1—N2 ⁱⁱ	2.259 (3)	C5—H5A	0.9300
Cd1—O1	2.267 (2)	C6—H6A	0.9300
Cd1—O1 ⁱ	2.267 (2)	C7—H7A	0.9600

S1—C1	1.769 (4)	C7—H7B	0.9600
S1—C8	1.809 (4)	C7—H7C	0.9600
O1—C9	1.266 (4)	C8—C9	1.532 (4)
O1W—H1WA	0.8200	C8—H8A	0.9700
O1W—H1WB	0.805 (17)	C8—H8B	0.9700
O2—C9	1.232 (4)	C10—C11	1.375 (4)
N1—C10 ⁱ	1.325 (3)	C10—H10A	0.9300
N1—C10	1.325 (3)	C11—C12	1.381 (3)
N2—C15	1.343 (3)	C11—H11A	0.9300
N2—C15 ⁱ	1.343 (3)	C12—C11 ⁱ	1.381 (3)
N2—Cd1 ⁱⁱⁱ	2.259 (3)	C12—C13	1.475 (5)
C1—C2	1.391 (5)	C13—C14 ⁱ	1.389 (3)
C1—C6	1.398 (5)	C13—C14	1.389 (3)
C2—C3	1.367 (7)	C14—C15	1.376 (4)
C2—H2A	0.9300	C14—H14A	0.9300
C3—C4	1.389 (6)	C15—H15A	0.9300
C3—H3A	0.9300		
N1—Cd1—O1W	93.87 (6)	C6—C5—H5A	119.2
N1—Cd1—O1W ⁱ	93.87 (6)	C4—C5—H5A	119.2
O1W—Cd1—O1W ⁱ	172.26 (12)	C5—C6—C1	121.0 (3)
N1—Cd1—N2 ⁱⁱ	180.0	C5—C6—H6A	119.5
O1W—Cd1—N2 ⁱⁱ	86.13 (6)	C1—C6—H6A	119.5
O1W ⁱ —Cd1—N2 ⁱⁱ	86.13 (6)	C4—C7—H7A	109.5
N1—Cd1—O1	91.48 (5)	C4—C7—H7B	109.5
O1W—Cd1—O1	90.67 (9)	H7A—C7—H7B	109.5
O1W ⁱ —Cd1—O1	89.13 (9)	C4—C7—H7C	109.5
N2 ⁱⁱ —Cd1—O1	88.52 (5)	H7A—C7—H7C	109.5
N1—Cd1—O1 ⁱ	91.48 (5)	H7B—C7—H7C	109.5
O1W—Cd1—O1 ⁱ	89.13 (9)	C9—C8—S1	114.1 (2)
O1W ⁱ —Cd1—O1 ⁱ	90.67 (9)	C9—C8—H8A	108.7
N2 ⁱⁱ —Cd1—O1 ⁱ	88.52 (5)	S1—C8—H8A	108.7
O1—Cd1—O1 ⁱ	177.04 (10)	C9—C8—H8B	108.7
C1—S1—C8	105.40 (18)	S1—C8—H8B	108.7
C9—O1—Cd1	123.9 (2)	H8A—C8—H8B	107.6
Cd1—O1W—H1WA	109.5	O2—C9—O1	126.0 (3)
Cd1—O1W—H1WB	123 (3)	O2—C9—C8	118.2 (3)
H1WA—O1W—H1WB	105.7	O1—C9—C8	115.9 (3)
C10 ⁱ —N1—C10	116.8 (4)	N1—C10—C11	123.4 (3)
C10 ⁱ —N1—Cd1	121.62 (18)	N1—C10—H10A	118.3
C10—N1—Cd1	121.62 (18)	C11—C10—H10A	118.3
C15—N2—C15 ⁱ	116.8 (4)	C10—C11—C12	120.2 (3)
C15—N2—Cd1 ⁱⁱⁱ	121.58 (18)	C10—C11—H11A	119.9
C15 ⁱ —N2—Cd1 ⁱⁱⁱ	121.58 (18)	C12—C11—H11A	119.9
C2—C1—C6	117.7 (4)	C11 ⁱ —C12—C11	116.0 (4)
C2—C1—S1	126.2 (3)	C11 ⁱ —C12—C13	122.02 (18)
C6—C1—S1	116.1 (3)	C11—C12—C13	122.02 (18)
C3—C2—C1	120.6 (4)	C14 ⁱ —C13—C14	116.2 (3)

C3—C2—H2A	119.7	C14 ⁱ —C13—C12	121.89 (17)
C1—C2—H2A	119.7	C14—C13—C12	121.89 (17)
C2—C3—C4	122.4 (3)	C15—C14—C13	120.4 (3)
C2—C3—H3A	118.8	C15—C14—H14A	119.8
C4—C3—H3A	118.8	C13—C14—H14A	119.8
C3—C4—C5	116.8 (4)	N2—C15—C14	123.0 (3)
C3—C4—C7	120.6 (4)	N2—C15—H15A	118.5
C5—C4—C7	122.6 (4)	C14—C15—H15A	118.5
C6—C5—C4	121.5 (4)		

Symmetry codes: (i) $-x, y, -z+1/2$; (ii) $x, y+1, z$; (iii) $x, y-1, z$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O1 <i>W</i> —H1 <i>WA</i> ...O2	0.82	1.94	2.667 (4)	148
O1 <i>W</i> —H1 <i>WB</i> ...O1 ^{iv}	0.81 (2)	2.04 (2)	2.782 (3)	154 (4)

Symmetry code: (iv) $x, -y, z+1/2$.