

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## *N*-(2,4-Dichlorophenyl)-2,4-dimethylbenzenesulfonamide

P. G. Nirmala,<sup>a</sup> Sabine Foro,<sup>b</sup> B. Thimme Gowda<sup>a\*</sup> and Hartmut Fuess<sup>b</sup>

<sup>a</sup>Department of Chemistry, Mangalore University, Mangalagangothri 574 199, Mangalore, India, and <sup>b</sup>Institute of Materials Science, Darmstadt University of Technology, Petersenstrasse 23, D-64287 Darmstadt, Germany  
Correspondence e-mail: gowdabt@yahoo.com


Received 5 November 2010; accepted 6 November 2010

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.004$  Å;  $R$  factor = 0.056;  $wR$  factor = 0.159; data-to-parameter ratio = 14.1.

In the title compound,  $\text{C}_{14}\text{H}_{13}\text{Cl}_2\text{NO}_2\text{S}$ , the molecule is bent at the S atom with an  $\text{C}-\text{SO}_2-\text{NH}-\text{C}$  torsion angle of  $-69.9$  (2)°. The dihedral angle between the sulfonyl and aniline benzene rings is  $44.0$  (1)°. The crystal structure features inversion dimers linked by pairs of  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds. An intramolecular  $\text{N}-\text{H}\cdots\text{Cl}$  hydrogen bond is also observed.

### Related literature

For the preparation of the compound, see: Savitha & Gowda (2006). For our study of the effect of substituents on the structures of *N*-(aryl)arylsulfonamides, see: Gowda *et al.* (2009); Nirmala *et al.* (2010*a,b*). For related structures, see: Gelbrich *et al.* (2007); Perlovich *et al.* (2006).



### Experimental

#### Crystal data

$\text{C}_{14}\text{H}_{13}\text{Cl}_2\text{NO}_2\text{S}$   
 $M_r = 330.21$   
Triclinic,  $P\bar{1}$   
 $a = 8.2407$  (9) Å

$b = 8.3418$  (9) Å  
 $c = 10.849$  (1) Å  
 $\alpha = 84.59$  (1)°  
 $\beta = 89.06$  (1)°

$\gamma = 85.07$  (1)°  
 $V = 739.69$  (13) Å<sup>3</sup>  
 $Z = 2$   
Cu  $K\alpha$  radiation

$\mu = 5.27$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.45 \times 0.40 \times 0.13$  mm

#### Data collection

Enraf–Nonius CAD-4 diffractometer  
Absorption correction:  $\psi$  scan (North *et al.*, 1968)  
 $T_{\min} = 0.200$ ,  $T_{\max} = 0.547$   
5173 measured reflections

2629 independent reflections  
2496 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.033$   
3 standard reflections every 120 min  
intensity decay: 1.0%

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.056$   
 $wR(F^2) = 0.159$   
 $S = 1.05$   
2629 reflections  
187 parameters  
1 restraint

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\text{max}} = 0.41$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.80$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1N}\cdots\text{O2}^i$	0.85 (2)	2.24 (2)	3.056 (3)	159 (3)
$\text{N1}-\text{H1N}\cdots\text{Cl1}$	0.85 (2)	2.68 (3)	3.020 (2)	105 (2)

Symmetry code: (i)  $-x, -y + 1, -z$ .

Data collection: *CAD-4-PC* (Enraf–Nonius, 1996); cell refinement: *CAD-4-PC*; data reduction: *REDU4* (Stoe & Cie, 1987); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2251).

### References

- Enraf–Nonius (1996). *CAD-4-PC*. Enraf–Nonius, Delft, The Netherlands.  
Gelbrich, T., Hursthouse, M. B. & Threlfall, T. L. (2007). *Acta Cryst.* **B63**, 621–632.  
Gowda, B. T., Foro, S., Nirmala, P. G., Babitha, K. S. & Fuess, H. (2009). *Acta Cryst.* **E65**, o576.  
Nirmala, P. G., Gowda, B. T., Foro, S. & Fuess, H. (2010*a*). *Acta Cryst.* **E66**, o1059.  
Nirmala, P. G., Gowda, B. T., Foro, S. & Fuess, H. (2010*b*). *Acta Cryst.* **E66**, o1090.  
North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.  
Perlovich, G. L., Tkachev, V. V., Schaper, K.-J. & Raevsky, O. A. (2006). *Acta Cryst.* **E62**, o780–o782.  
Savitha, M. B. & Gowda, B. T. (2006). *Z. Naturforsch. Teil A*, **60**, 600–606.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.  
Stoe & Cie (1987). *REDU4*. Stoe & Cie GmbH, Darmstadt, Germany.

## supporting information

*Acta Cryst.* (2010). E66, o3144 [https://doi.org/10.1107/S1600536810045563]

***N*-(2,4-Dichlorophenyl)-2,4-dimethylbenzenesulfonamide**

**P. G. Nirmala, Sabine Foro, B. Thimme Gowda and Hartmut Fuess**

**S1. Comment**

In the present work, as part of a study of the substituent effects on the structures of *N*-(aryl)arylsulfonamides (Gowda *et al.*, 2009; Nirmala *et al.*, 2010*a,b*), the structure of 2,4-dimethyl-*N*-(2,4-dichlorophenyl)benzenesulfonamide (I) has been determined (Fig. 1). The molecule is bent at the *N* atom with the C1—SO<sub>2</sub>—NH—C7 torsion angle of -69.9 (2)°, compared to the values of 46.1 (3)° (glide image of molecule 1) and 47.7 (3)° (molecule 2) in the two independent molecules of 2,4-dimethyl-*N*-(phenyl)benzenesulfonamide (II) (Gowda *et al.*, 2009), -54.9 (3)° in 2,4-dimethyl-*N*-(3,5-dichlorophenyl)benzenesulfonamide (III) (Nirmala *et al.*, 2010*b*) and 66.5 (2)° in 2,4-dimethyl-*N*-(2,4-dimethylphenyl)benzenesulfonamide (IV) (Nirmala *et al.*, 2010*a*)

The two benzene rings in (I) are tilted relative to each other by 44.0 (1)°, compared to the values of 67.5 (1)° (molecule 1) and 72.9 (1)° (molecule 2) in the two independent molecules of (II), 82.3 (1)° in (III) and 41.0 (1)° in (IV). The other bond parameters in (I) are similar to those observed in (II), (III), (IV) and other aryl sulfonamides (Perlovich *et al.*, 2006; Gelbrich *et al.*, 2007).

The structure shows simultaneous N—H···Cl intramolecular and N—H···O intermolecular H-bonding (Table 1). The crystal packing of molecules in (I) *via* N—H···O(S) hydrogen bonds is shown in Fig.2.


**S2. Experimental**

The solution of 1,3-xylene (1,3-dimethylbenzene) (10 ml) in chloroform (40 ml) was treated dropwise with chlorosulfonic acid (25 ml) at 0 ° C. After the initial evolution of hydrogen chloride subsided, the reaction mixture was brought to room temperature and poured into crushed ice in a beaker. The chloroform layer was separated, washed with cold water and allowed to evaporate slowly. The residual 2,4-dimethylbenzenesulfonylchloride was treated with 2,4-dichloroaniline in the stoichiometric ratio and boiled for ten minutes. The reaction mixture was then cooled to room temperature and added to ice cold water (100 ml). The resultant solid 2,4-dimethyl-*N*-(2,4-dichlorophenyl)benzenesulfonamide was filtered under suction and washed thoroughly with cold water. It was then recrystallized to constant melting point from dilute ethanol. The purity of the compound was checked and characterized by recording its infrared and NMR spectra (Savitha & Gowda, 2006).


The plate like colorless single crystals used in X-ray diffraction studies were grown in ethanolic solution by a slow evaporation at room temperature.

**S3. Refinement**

The H atoms of the NH groups were located in a difference map and later restrained to N—H = 0.86 (2) Å. The other H atoms were positioned with idealized geometry using a riding model with C—H = 0.93–0.96 Å. All H atoms were refined with isotropic displacement parameters (set to 1.2 times of the  $U_{eq}$  of the parent atom).

**Figure 1**

Molecular structure of (I), showing the atom labeling scheme and displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

Molecular packing of (I) with hydrogen bonding shown as dashed lines.

*N*-(2,4-Dichlorophenyl)-2,4-dimethylbenzenesulfonamide

## Crystal data

$C_{14}H_{13}Cl_2NO_2S$   
 $M_r = 330.21$   
 Triclinic,  $P\bar{1}$   
 Hall symbol:  $-P\ 1$   
 $a = 8.2407$  (9) Å  
 $b = 8.3418$  (9) Å  
 $c = 10.849$  (1) Å  
 $\alpha = 84.59$  (1)°  
 $\beta = 89.06$  (1)°  
 $\gamma = 85.07$  (1)°  
 $V = 739.69$  (13) Å<sup>3</sup>

$Z = 2$   
 $F(000) = 340$   
 $D_x = 1.483$  Mg m<sup>-3</sup>  
 Cu  $K\alpha$  radiation,  $\lambda = 1.54180$  Å  
 Cell parameters from 25 reflections  
 $\theta = 6.7$ – $18.3$ °  
 $\mu = 5.27$  mm<sup>-1</sup>  
 $T = 293$  K  
 Plate, colorless  
 $0.45 \times 0.40 \times 0.13$  mm

## Data collection

Enraf–Nonius CAD-4  
 diffractometer  
 Radiation source: fine-focus sealed tube  
 Graphite monochromator  
 $\omega/2\theta$  scans  
 Absorption correction:  $\psi$  scan  
 (North *et al.*, 1968)  
 $T_{\min} = 0.200$ ,  $T_{\max} = 0.547$   
 5173 measured reflections

2629 independent reflections  
 2496 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.033$   
 $\theta_{\text{max}} = 66.9$ °,  $\theta_{\text{min}} = 4.1$ °  
 $h = -9 \rightarrow 9$   
 $k = -9 \rightarrow 9$   
 $l = -12 \rightarrow 12$   
 3 standard reflections every 120 min  
 intensity decay: 1.0%

## Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.056$   
 $wR(F^2) = 0.159$   
 $S = 1.05$   
 2629 reflections  
 187 parameters  
 1 restraint  
 Primary atom site location: structure-invariant  
 direct methods  
 Secondary atom site location: difference Fourier  
 map

Hydrogen site location: inferred from  
 neighbouring sites  
 H atoms treated by a mixture of independent  
 and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.1122P)^2 + 0.3056P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} = 0.003$   
 $\Delta\rho_{\text{max}} = 0.41$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.80$  e Å<sup>-3</sup>  
 Extinction correction: *SHELXL97* (Sheldrick,  
 2008),  $F_c^* = kFc[1 + 0.001xFe^2\lambda^3/\sin(2\theta)]^{-1/4}$   
 Extinction coefficient: 0.0131 (18)

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å<sup>2</sup>)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.0771 (3)	0.4275 (3)	0.2785 (2)	0.0417 (5)

C2	0.1504 (3)	0.3433 (3)	0.3839 (2)	0.0494 (6)
C3	0.0454 (4)	0.2704 (4)	0.4696 (3)	0.0593 (7)
H3	0.0903	0.2133	0.5405	0.071*
C4	-0.1211 (4)	0.2779 (4)	0.4556 (3)	0.0586 (7)
C5	-0.1883 (4)	0.3659 (4)	0.3526 (3)	0.0571 (7)
H5	-0.3007	0.3764	0.3428	0.069*
C6	-0.0905 (3)	0.4382 (3)	0.2642 (2)	0.0497 (6)
H6	-0.1372	0.4952	0.1939	0.060*
C7	0.3576 (3)	0.2472 (3)	0.0982 (2)	0.0385 (5)
C8	0.3155 (3)	0.0907 (3)	0.1315 (2)	0.0394 (5)
C9	0.4329 (3)	-0.0337 (3)	0.1657 (2)	0.0440 (6)
H9	0.4039	-0.1381	0.1872	0.053*
C10	0.5934 (3)	0.0004 (3)	0.1671 (2)	0.0448 (6)
C11	0.6398 (3)	0.1535 (3)	0.1328 (2)	0.0496 (6)
H11	0.7490	0.1741	0.1331	0.060*
C12	0.5214 (3)	0.2753 (3)	0.0980 (2)	0.0459 (6)
H12	0.5516	0.3785	0.0739	0.055*
C13	0.3296 (4)	0.3251 (5)	0.4112 (3)	0.0705 (9)
H13A	0.3685	0.4300	0.4134	0.085*
H13B	0.3869	0.2698	0.3477	0.085*
H13C	0.3478	0.2639	0.4899	0.085*
C14	-0.2268 (5)	0.1923 (5)	0.5518 (4)	0.0833 (11)
H14A	-0.1703	0.1745	0.6292	0.100*
H14B	-0.2503	0.0904	0.5250	0.100*
H14C	-0.3268	0.2579	0.5621	0.100*
N1	0.2394 (3)	0.3750 (2)	0.06051 (18)	0.0425 (5)
H1N	0.156 (3)	0.347 (4)	0.026 (3)	0.051*
O1	0.3348 (2)	0.5697 (2)	0.1953 (2)	0.0564 (5)
O2	0.0776 (3)	0.6299 (2)	0.08078 (18)	0.0559 (5)
Cl1	0.11490 (8)	0.04794 (8)	0.12875 (7)	0.0581 (3)
Cl2	0.74198 (9)	-0.15459 (9)	0.20963 (7)	0.0672 (3)
S1	0.18758 (7)	0.51721 (7)	0.15244 (5)	0.0428 (3)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0409 (12)	0.0415 (13)	0.0434 (12)	-0.0039 (9)	0.0029 (10)	-0.0073 (10)
C2	0.0487 (15)	0.0545 (16)	0.0446 (13)	0.0003 (11)	-0.0017 (11)	-0.0077 (11)
C3	0.0670 (18)	0.0663 (18)	0.0424 (13)	0.0009 (14)	-0.0003 (12)	0.0003 (12)
C4	0.0614 (17)	0.0637 (18)	0.0516 (15)	-0.0101 (14)	0.0112 (13)	-0.0078 (13)
C5	0.0440 (14)	0.0716 (19)	0.0575 (15)	-0.0091 (13)	0.0056 (12)	-0.0119 (13)
C6	0.0450 (14)	0.0559 (15)	0.0475 (13)	-0.0012 (11)	-0.0019 (10)	-0.0034 (11)
C7	0.0397 (12)	0.0382 (12)	0.0374 (11)	-0.0034 (9)	0.0025 (9)	-0.0027 (9)
C8	0.0398 (12)	0.0406 (13)	0.0384 (11)	-0.0086 (9)	0.0006 (9)	-0.0029 (9)
C9	0.0503 (14)	0.0380 (12)	0.0429 (12)	-0.0041 (10)	-0.0009 (10)	0.0008 (9)
C10	0.0440 (13)	0.0455 (13)	0.0439 (12)	0.0053 (10)	-0.0032 (10)	-0.0071 (10)
C11	0.0381 (13)	0.0524 (15)	0.0599 (15)	-0.0058 (11)	0.0026 (11)	-0.0122 (12)
C12	0.0413 (13)	0.0404 (13)	0.0568 (14)	-0.0082 (10)	0.0079 (10)	-0.0058 (10)

C13	0.0512 (17)	0.097 (3)	0.0608 (17)	0.0054 (16)	-0.0101 (13)	-0.0045 (16)
C14	0.084 (3)	0.092 (3)	0.073 (2)	-0.018 (2)	0.0244 (19)	0.0047 (19)
N1	0.0435 (11)	0.0382 (11)	0.0450 (11)	-0.0008 (8)	0.0010 (8)	-0.0015 (8)
O1	0.0508 (11)	0.0462 (11)	0.0749 (12)	-0.0120 (8)	0.0031 (9)	-0.0134 (9)
O2	0.0590 (12)	0.0403 (10)	0.0646 (11)	0.0061 (8)	0.0026 (9)	0.0059 (8)
Cl1	0.0405 (4)	0.0519 (4)	0.0825 (5)	-0.0131 (3)	-0.0042 (3)	-0.0001 (3)
Cl2	0.0586 (5)	0.0630 (5)	0.0758 (5)	0.0180 (3)	-0.0144 (4)	-0.0036 (4)
S1	0.0434 (4)	0.0333 (4)	0.0512 (4)	-0.0023 (2)	0.0033 (3)	-0.0024 (3)

*Geometric parameters (Å, °)*

C1—C6	1.387 (4)	C9—C10	1.378 (4)
C1—C2	1.402 (4)	C9—H9	0.9300
C1—S1	1.773 (2)	C10—C11	1.381 (4)
C2—C3	1.393 (4)	C10—Cl2	1.737 (2)
C2—C13	1.502 (4)	C11—C12	1.377 (4)
C3—C4	1.378 (5)	C11—H11	0.9300
C3—H3	0.9300	C12—H12	0.9300
C4—C5	1.374 (4)	C13—H13A	0.9600
C4—C14	1.515 (4)	C13—H13B	0.9600
C5—C6	1.372 (4)	C13—H13C	0.9600
C5—H5	0.9300	C14—H14A	0.9600
C6—H6	0.9300	C14—H14B	0.9600
C7—C12	1.389 (3)	C14—H14C	0.9600
C7—C8	1.393 (3)	N1—S1	1.645 (2)
C7—N1	1.416 (3)	N1—H1N	0.853 (18)
C8—C9	1.384 (3)	O1—S1	1.424 (2)
C8—C11	1.722 (2)	O2—S1	1.4311 (19)
C6—C1—C2	120.8 (2)	C11—C10—Cl2	119.1 (2)
C6—C1—S1	115.43 (19)	C12—C11—C10	118.7 (2)
C2—C1—S1	123.7 (2)	C12—C11—H11	120.6
C3—C2—C1	115.9 (3)	C10—C11—H11	120.6
C3—C2—C13	118.1 (3)	C11—C12—C7	121.4 (2)
C1—C2—C13	125.9 (3)	C11—C12—H12	119.3
C4—C3—C2	123.9 (3)	C7—C12—H12	119.3
C4—C3—H3	118.1	C2—C13—H13A	109.5
C2—C3—H3	118.1	C2—C13—H13B	109.5
C5—C4—C3	118.2 (3)	H13A—C13—H13B	109.5
C5—C4—C14	121.1 (3)	C2—C13—H13C	109.5
C3—C4—C14	120.7 (3)	H13A—C13—H13C	109.5
C6—C5—C4	120.4 (3)	H13B—C13—H13C	109.5
C6—C5—H5	119.8	C4—C14—H14A	109.5
C4—C5—H5	119.8	C4—C14—H14B	109.5
C5—C6—C1	120.8 (3)	H14A—C14—H14B	109.5
C5—C6—H6	119.6	C4—C14—H14C	109.5
C1—C6—H6	119.6	H14A—C14—H14C	109.5
C12—C7—C8	118.3 (2)	H14B—C14—H14C	109.5

C12—C7—N1	119.8 (2)	C7—N1—S1	120.33 (16)
C8—C7—N1	121.9 (2)	C7—N1—H1N	115 (2)
C9—C8—C7	121.2 (2)	S1—N1—H1N	110 (2)
C9—C8—C11	118.61 (18)	O1—S1—O2	119.34 (13)
C7—C8—C11	120.21 (19)	O1—S1—N1	106.94 (12)
C10—C9—C8	118.6 (2)	O2—S1—N1	104.56 (11)
C10—C9—H9	120.7	O1—S1—C1	110.04 (12)
C8—C9—H9	120.7	O2—S1—C1	108.11 (12)
C9—C10—C11	121.8 (2)	N1—S1—C1	107.11 (11)
C9—C10—C12	119.1 (2)		
C6—C1—C2—C3	1.3 (4)	C8—C9—C10—C11	-1.5 (4)
S1—C1—C2—C3	-175.4 (2)	C8—C9—C10—C12	-179.91 (18)
C6—C1—C2—C13	-179.0 (3)	C9—C10—C11—C12	1.0 (4)
S1—C1—C2—C13	4.4 (4)	C12—C10—C11—C12	179.3 (2)
C1—C2—C3—C4	-0.2 (4)	C10—C11—C12—C7	0.7 (4)
C13—C2—C3—C4	-180.0 (3)	C8—C7—C12—C11	-1.7 (4)
C2—C3—C4—C5	-1.8 (5)	N1—C7—C12—C11	-179.3 (2)
C2—C3—C4—C14	178.7 (3)	C12—C7—N1—S1	-74.9 (3)
C3—C4—C5—C6	2.6 (5)	C8—C7—N1—S1	107.6 (2)
C14—C4—C5—C6	-177.8 (3)	C7—N1—S1—O1	48.1 (2)
C4—C5—C6—C1	-1.6 (4)	C7—N1—S1—O2	175.56 (18)
C2—C1—C6—C5	-0.4 (4)	C7—N1—S1—C1	-69.8 (2)
S1—C1—C6—C5	176.5 (2)	C6—C1—S1—O1	151.8 (2)
C12—C7—C8—C9	1.1 (4)	C2—C1—S1—O1	-31.4 (3)
N1—C7—C8—C9	178.6 (2)	C6—C1—S1—O2	19.9 (2)
C12—C7—C8—C11	-178.02 (18)	C2—C1—S1—O2	-163.3 (2)
N1—C7—C8—C11	-0.5 (3)	C6—C1—S1—N1	-92.3 (2)
C7—C8—C9—C10	0.5 (4)	C2—C1—S1—N1	84.5 (2)
C11—C8—C9—C10	179.62 (18)		

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1N...O2 <sup>i</sup>	0.85 (2)	2.24 (2)	3.056 (3)	159 (3)
N1—H1N...C11	0.85 (2)	2.68 (3)	3.020 (2)	105 (2)

Symmetry code: (i)  $-x, -y+1, -z$ .