

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

2-(5,6-Dibromo-7-methyl-3*H*-imidazo-  
[4,5-*b*]pyridin-2-yl)phenol

Haixia Wang

Department of Chemistry and Environmental Science, Henan Normal University,  
Xinxiang 453007, People's Republic of China  
Correspondence e-mail: xxhxiawang@126.com


Received 25 October 2010; accepted 4 November 2010

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(\text{C}-\text{C}) = 0.008$  Å;  
 $R$  factor = 0.043;  $wR$  factor = 0.138; data-to-parameter ratio = 12.9.

In the title compound,  $\text{C}_{13}\text{H}_9\text{Br}_2\text{N}_3\text{O}$ , the molecular skeleton, influenced by an intramolecular  $\text{O}-\text{H}\cdots\text{N}$  hydrogen bond, is roughly planar, with a mean deviation of 0.033 Å. In the crystal, intermolecular  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds link the molecules into chains propagating in [100]. Weak intermolecular  $\pi-\pi$  interactions [centroid-centroid distances = 3.760 (3) and 3.723 (3) Å] further consolidate the packing.

## Related literature

For background to the use of imidazole and its derivatives in transition metal complexes, see: Huang *et al.* (2004). For related structures, see: Eltayeb *et al.* (2009); Xiao *et al.* (2009); Elerman & Kabak (1997).



## Experimental

## Crystal data

$\text{C}_{13}\text{H}_9\text{Br}_2\text{N}_3\text{O}$   
 $M_r = 383.05$

Orthorhombic,  $Pbca$   
 $a = 13.181$  (5) Å

$b = 8.494$  (3) Å  
 $c = 22.692$  (8) Å  
 $V = 2540.5$  (16) Å<sup>3</sup>  
 $Z = 8$

Mo  $K\alpha$  radiation  
 $\mu = 6.38$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.31 \times 0.28 \times 0.24$  mm

## Data collection

Bruker APEXII CCD area-detector diffractometer  
Absorption correction: multi-scan (SADABS; Sheldrick, 2008a)  
 $T_{\min} = 0.243$ ,  $T_{\max} = 0.310$

11656 measured reflections  
2234 independent reflections  
1706 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.041$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$   
 $wR(F^2) = 0.138$   
 $S = 1.08$   
2234 reflections

173 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.63$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.88$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O1}^i$	0.95	1.90	2.839 (6)	171
$\text{O1}-\text{H1}\cdots\text{N2}$	0.82	1.84	2.573 (6)	149

Symmetry code: (i)  $x + \frac{1}{2}, y, -z + \frac{3}{2}$ .

Data collection: APEX2 (Bruker, 2004); cell refinement: SAINT-Plus (Bruker, 2001); data reduction: SAINT-Plus; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008b); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008b); molecular graphics: SHELXTL (Sheldrick, 2008b); software used to prepare material for publication: SHELXTL.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2784).

## References

- Bruker (2001). SAINT-Plus. Bruker AXS Inc., Madison, Wisconsin, USA.  
Bruker (2004). APEX2. Bruker AXS Inc., Madison, Wisconsin, USA.  
Elerman, Y. & Kabak, M. (1997). *Acta Cryst.* **C53**, 372–374.  
Eltayeb, N. E., Teoh, S. G., Quah, C. K., Fun, H.-K. & Adnan, R. (2009). *Acta Cryst.* **E65**, o1613–o1614.  
Huang, X.-C., Zhang, J.-P. & Chen, X.-M. (2004). *J. Am. Chem. Soc.* **126**, 13218–13219.  
Sheldrick, G. M. (2008a). SADABS. University of Göttingen, Germany.  
Sheldrick, G. M. (2008b). *Acta Cryst.* **A64**, 112–122.  
Xiao, H.-Q., Zhang, M.-Z. & Wang, W. (2009). *Acta Cryst.* **E65**, o1256.

## supporting information

*Acta Cryst.* (2010). E66, o3132 [https://doi.org/10.1107/S1600536810045277]

**2-(5,6-Dibromo-7-methyl-3*H*-imidazo[4,5-*b*]pyridin-2-yl)phenol****Haixia Wang****S1. Comment**

Due to excellent coordination abilities the imidazole and its derivatives have already been introduced into the transition metal complexes (Huang *et al.*, 2004). Herewith we present the title compound (I) - a new imidazole derivative.

In (I) (Fig. 1), intramolecular O—H $\cdots$ N hydrogen bond (Table 2) influence the molecular conformation, so all non-H atoms are nearly coplanar with the mean deviation of 0.033 Å. The dihedral angle between the 5,6-dibromo-7-methyl-3*H*-imidazo[4,5-*b*]pyridine plane and the phenol plane is 2.1 (2) °. The bond lengths and angles are normal and comparable to those observed in the reported benzimidazole compounds (Xiao *et al.*, 2009; Eltayeb *et al.*, 2009; Elerman & Kabak 1997).

In the crystal structure, intermolecular N—H $\cdots$ O hydrogen bonds (Table 2) link the molecules into chains propagated in direction [100]. Weak intermolecular  $\pi$ — $\pi$  interactions (Table 1) consolidate further the crystal packing.

**S2. Experimental**

The title compound was synthesized by the reaction of 4-methyl-2,3-diamino-5,6-dibromopyridine and 2-hydroxybenzaldehyde with the ratio 1:1 in ethanol. After the mixture was refluxed several hours, the resulting clear yellow solution was allowed to evaporate slowly in air, and orange-yellow block-like crystals suitable for X-ray diffraction were obtained with a yield 47% about ten days later.

**S3. Refinement**

All the H atoms bonded to the C atoms were placed using the HFIX commands in *SHELXL-97* with C—H distances of 0.93 and 0.96 Å, and were refined as riding, with  $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C})$ . H atoms bonded to O and N atoms were found from difference Fourier maps with the bond lengths restrained to 0.82 and 0.96 Å, respectively, and were refined as riding, with  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$  and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ .




Figure 1

The molecular structure of the title compound showing the atom-labelling scheme and 30% probability displacement ellipsoids.

## 2-(5,6-Dibromo-7-methyl-3H-imidazo[4,5-*b*]pyridin-2-yl)phenol

### Crystal data

$C_{13}H_9Br_2N_3O$

$M_r = 383.05$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 13.181 (5) \text{ \AA}$

$b = 8.494 (3) \text{ \AA}$

$c = 22.692 (8) \text{ \AA}$

$V = 2540.5 (16) \text{ \AA}^3$

$Z = 8$

$F(000) = 1488$

$D_x = 2.003 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 1452 reflections

$\theta = 2.9\text{--}24.6^\circ$

$\mu = 6.38 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, orange–yellow

$0.31 \times 0.28 \times 0.24 \text{ mm}$

### Data collection

Bruker APEXII CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 2008a)

$T_{\min} = 0.243$ ,  $T_{\max} = 0.310$

11656 measured reflections

2234 independent reflections

1706 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.041$

$\theta_{\max} = 25.0^\circ$ ,  $\theta_{\min} = 1.8^\circ$

$h = -15 \rightarrow 15$

$k = -9 \rightarrow 10$

$l = -24 \rightarrow 26$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.043$

$wR(F^2) = 0.138$

$S = 1.08$

2234 reflections

173 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0692P)^2 + 6.558P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.63 \text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.88 \text{ e \AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.27968 (6)	1.22305 (9)	0.48970 (3)	0.0618 (3)
Br2	0.51701 (6)	1.19032 (9)	0.54180 (3)	0.0669 (3)
O1	0.0310 (3)	0.7707 (6)	0.73898 (19)	0.0541 (12)
H1	0.0614	0.8111	0.7113	0.081*
N1	0.3413 (3)	0.8522 (5)	0.7102 (2)	0.0380 (10)
H1A	0.4011	0.8215	0.7306	0.046*
N2	0.1806 (3)	0.8887 (5)	0.68102 (19)	0.0365 (10)
N3	0.4280 (4)	1.0090 (7)	0.6313 (2)	0.0615 (15)
C1	0.3468 (4)	0.9445 (6)	0.6601 (2)	0.0348 (12)
C2	0.2452 (4)	0.9659 (6)	0.6416 (2)	0.0350 (12)
C3	0.2226 (4)	1.0517 (7)	0.5909 (2)	0.0432 (13)
C4	0.3060 (4)	1.1146 (7)	0.5612 (2)	0.0414 (13)
C5	0.4049 (4)	1.0950 (6)	0.5811 (2)	0.0400 (13)
C6	0.1127 (5)	1.0806 (9)	0.5643 (3)	0.0680 (19)
H6A	0.0923	1.1873	0.5717	0.102*
H6B	0.0653	1.0098	0.5825	0.102*
H6C	0.1140	1.0621	0.5226	0.102*
C7	0.2410 (4)	0.8221 (6)	0.7213 (2)	0.0357 (12)
C8	0.2041 (4)	0.7275 (6)	0.7696 (2)	0.0354 (12)
C9	0.0988 (5)	0.7064 (7)	0.7768 (3)	0.0431 (13)
C10	0.0636 (5)	0.6169 (8)	0.8237 (3)	0.0578 (17)
H10	-0.0057	0.6022	0.8288	0.069*
C11	0.1315 (5)	0.5488 (8)	0.8632 (3)	0.0604 (18)
H11	0.1074	0.4895	0.8947	0.072*
C12	0.2329 (5)	0.5690 (8)	0.8558 (3)	0.0536 (16)
H12	0.2779	0.5226	0.8821	0.064*
C13	0.2693 (4)	0.6560 (7)	0.8104 (3)	0.0453 (14)
H13	0.3390	0.6686	0.8063	0.054*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.0851 (6)	0.0621 (5)	0.0383 (4)	0.0019 (4)	-0.0058 (3)	0.0082 (3)
Br2	0.0640 (5)	0.0757 (5)	0.0609 (5)	-0.0265 (4)	0.0124 (3)	0.0054 (4)
O1	0.029 (2)	0.079 (3)	0.054 (3)	0.0054 (19)	0.0065 (19)	0.007 (2)
N1	0.027 (2)	0.048 (3)	0.039 (3)	-0.0008 (19)	-0.0055 (19)	0.001 (2)

N2	0.028 (2)	0.045 (3)	0.037 (3)	0.0009 (18)	-0.0012 (19)	-0.002 (2)
N3	0.056 (3)	0.068 (4)	0.060 (4)	-0.006 (3)	0.003 (3)	-0.005 (3)
C1	0.029 (3)	0.043 (3)	0.033 (3)	-0.004 (2)	0.000 (2)	-0.003 (2)
C2	0.033 (3)	0.040 (3)	0.032 (3)	-0.001 (2)	0.000 (2)	-0.004 (2)
C3	0.048 (3)	0.046 (3)	0.036 (3)	0.007 (3)	-0.002 (3)	-0.005 (3)
C4	0.050 (3)	0.041 (3)	0.034 (3)	-0.002 (3)	0.000 (3)	-0.003 (2)
C5	0.045 (3)	0.040 (3)	0.036 (3)	-0.007 (2)	0.006 (3)	0.000 (2)
C6	0.071 (5)	0.076 (5)	0.057 (4)	0.012 (4)	0.011 (4)	0.010 (4)
C7	0.031 (3)	0.040 (3)	0.036 (3)	-0.001 (2)	0.004 (2)	-0.007 (2)
C8	0.038 (3)	0.034 (3)	0.034 (3)	-0.002 (2)	0.004 (2)	-0.004 (2)
C9	0.044 (3)	0.044 (3)	0.041 (3)	0.001 (3)	0.009 (3)	-0.006 (3)
C10	0.054 (4)	0.061 (4)	0.059 (4)	-0.008 (3)	0.023 (3)	-0.001 (3)
C11	0.082 (5)	0.051 (4)	0.048 (4)	-0.004 (3)	0.018 (3)	0.006 (3)
C12	0.066 (4)	0.052 (4)	0.043 (4)	0.001 (3)	-0.004 (3)	0.008 (3)
C13	0.042 (3)	0.047 (3)	0.047 (4)	-0.001 (3)	-0.006 (3)	0.003 (3)

*Geometric parameters (Å, °)*

Br1—C4	1.897 (6)	C4—C5	1.389 (8)
Br2—C5	1.907 (5)	C6—H6A	0.9600
O1—C9	1.354 (7)	C6—H6B	0.9600
O1—H1	0.8200	C6—H6C	0.9600
N1—C7	1.369 (7)	C7—C8	1.442 (8)
N1—C1	1.382 (7)	C8—C13	1.402 (8)
N1—H1A	0.9504	C8—C9	1.408 (8)
N2—C7	1.338 (7)	C9—C10	1.388 (8)
N2—C2	1.398 (7)	C10—C11	1.392 (9)
N3—C1	1.370 (7)	C10—H10	0.9300
N3—C5	1.387 (8)	C11—C12	1.358 (9)
C1—C2	1.415 (7)	C11—H11	0.9300
C2—C3	1.394 (8)	C12—C13	1.355 (8)
C3—C4	1.396 (8)	C12—H12	0.9300
C3—C6	1.589 (9)	C13—H13	0.9300
Cg1...Cg2 <sup>i</sup>	3.760 (3)	Cg1...Cg3 <sup>ii</sup>	3.723 (3)
C9—O1—H1	109.5	C3—C6—H6C	109.5
C7—N1—C1	107.9 (4)	H6A—C6—H6C	109.5
C7—N1—H1A	131.3	H6B—C6—H6C	109.5
C1—N1—H1A	120.8	N2—C7—N1	111.7 (5)
C7—N2—C2	105.9 (4)	N2—C7—C8	123.6 (5)
C1—N3—C5	115.6 (5)	N1—C7—C8	124.7 (5)
N3—C1—N1	131.4 (5)	C13—C8—C9	118.1 (5)
N3—C1—C2	123.1 (5)	C13—C8—C7	122.4 (5)
N1—C1—C2	105.6 (4)	C9—C8—C7	119.5 (5)
C3—C2—N2	130.1 (5)	O1—C9—C10	119.1 (6)
C3—C2—C1	121.0 (5)	O1—C9—C8	121.6 (5)
N2—C2—C1	109.0 (5)	C10—C9—C8	119.3 (6)

C2—C3—C4	115.5 (5)	C9—C10—C11	120.4 (6)
C2—C3—C6	126.1 (5)	C9—C10—H10	119.8
C4—C3—C6	118.4 (5)	C11—C10—H10	119.8
C5—C4—C3	122.4 (5)	C12—C11—C10	120.0 (6)
C5—C4—Br1	120.5 (4)	C12—C11—H11	120.0
C3—C4—Br1	117.1 (4)	C10—C11—H11	120.0
N3—C5—C4	122.5 (5)	C13—C12—C11	120.8 (6)
N3—C5—Br2	115.9 (4)	C13—C12—H12	119.6
C4—C5—Br2	121.6 (4)	C11—C12—H12	119.6
C3—C6—H6A	109.5	C12—C13—C8	121.5 (5)
C3—C6—H6B	109.5	C12—C13—H13	119.3
H6A—C6—H6B	109.5	C8—C13—H13	119.3

Symmetry codes: (i)  $-x+1/2, y+1/2, z$ ; (ii)  $-x+1/2, y-1/2, z$ .

*Hydrogen-bond geometry (Å, °)*

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1A...O1 <sup>iii</sup>	0.95	1.90	2.839 (6)	171
O1—H1...N2	0.82	1.84	2.573 (6)	149

Symmetry code: (iii)  $x+1/2, y, -z+3/2$ .