

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(E)-1-[1-(4-Chlorophenyl)ethyl]-3,5-dimethyl-N-nitro-1,3,5-triazinan-2-imine

Huai-gang Su and Liang-zhong Xu*

 College of Chemistry and Molecular Engineering, Qingdao University of Science and Technology, Qingdao 266042, People's Republic of China
 Correspondence e-mail: qknh@s@yahoo.com.cn


Received 4 November 2010; accepted 9 November 2010

 Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.041; wR factor = 0.104; data-to-parameter ratio = 13.4.

In the title compound, $\text{C}_{13}\text{H}_{18}\text{ClN}_5\text{O}_2$, the 1,3,5-triazinane ring exhibits an envelope conformation with an *E* form. The chlorophenyl ring and the nitro group are each twisted with respect to the mean plane of the triazinane ring, making dihedral angles of 67.30 (9) and 83.54 (8)°, respectively. In the crystal, weak intermolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds build up a corrugated layer parallel to the (101) plane.

Related literature

The title compound was synthesized as a new compound with better insecticidal activity. For similar compounds with insecticidal properties, see: Koln *et al.* (2002). For related structures, see: Zhao *et al.* (2008); Hu *et al.* (2008); Xu *et al.* (2010) For puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

 $\text{C}_{13}\text{H}_{18}\text{ClN}_5\text{O}_2$
 $M_r = 311.77$

 Monoclinic, $P2_1/c$
 $a = 7.2483$ (14) Å
 $b = 29.568$ (6) Å
 $c = 7.2306$ (14) Å
 $\beta = 108.75$ (3)°
 $V = 1467.4$ (5) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.27$ mm⁻¹
 $T = 113$ K
 $0.20 \times 0.16 \times 0.12$ mm

Data collection

 Rigaku Saturn CCD area-detector diffractometer
 Absorption correction: multi-scan (ABSCOR; Higashi, 1995)
 $T_{\min} = 0.947$, $T_{\max} = 0.968$

 9753 measured reflections
 2585 independent reflections
 2303 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.039$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.104$
 $S = 1.08$
 2585 reflections

 193 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.46$ e Å⁻³
 $\Delta\rho_{\min} = -0.51$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C8}-\text{H8A}\cdots\text{O2}^{\text{i}}$	0.98	2.29	3.273 (2)	178
$\text{C10}-\text{H10A}\cdots\text{O1}^{\text{ii}}$	0.99	2.43	3.278 (2)	143
$\text{C11}-\text{H11B}\cdots\text{O2}^{\text{i}}$	0.99	2.49	3.434 (3)	160

 Symmetry codes: (i) $x + 1, y, z$; (ii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *RAPID-AUTO* (Rigaku, 2004); cell refinement: *RAPID-AUTO*; data reduction: *RAPID-AUTO*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *ORTEP III* (Burnett & Johnson, 1996) and *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXTL*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2619).

References

- Burnett, M. N. & Johnson, C. K. (1996). *ORTEP III*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Hu, Z.-Q., Yang, X.-D., An, G.-W., Yang, Z. & Xu, L.-Z. (2008). *Acta Cryst. E* **64**, o121.
- Koln, J. S., Wolfenbuttel, K. L., Dusseldorf, W. S., Bergisch Gladbach, A. H. & Koln, T. T. (2002). US Patent 0869120.
- Rigaku (2004). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Xu, L.-Z., Yin, R.-F. & Li, H.-X. (2010). *Acta Cryst. E* **66**, o867.
- Zhao, C., Yang, W., Hu, Y., Shen, L. & Lu, X. (2008). *Acta Cryst. E* **64**, o1515.

supporting information

Acta Cryst. (2010). E66, o3201 [https://doi.org/10.1107/S1600536810046246]

(E)-1-[1-(4-Chlorophenyl)ethyl]-3,5-dimethyl-N-nitro-1,3,5-triazinan-2-imine**Huai-gang Su and Liang-zhong Xu****S1. Comment**

The title compound was synthesized as a new compound with better insecticide activity. Lots of similar insecticide compounds with (I) were synthesized (Koln *et al.*, 2002). We report here the crystal structure of (I).

The 1,3,5-triazinan ring exhibits envelope conformation with puckering parameters $Q = 0.4777(19)\text{Å}$, $\theta = 57.8(2)^\circ$ and $\varphi = 238.5(3)^\circ$ (Cremer & Pople, 1975). The chlorophenyl ring as well as the nitro group are twisted with respect to the mean plane of the triazinan ring making dihedral angles of $67.30(9)^\circ$ and $83.54(8)^\circ$ respectively (Fig. 1). All bond lengths and angles are normal and in a good agreement with those recently reported (Hu *et al.*, 2008; Zhao *et al.*, 2008; Xu *et al.*, 2010).

Weak intermolecular C–H \cdots O hydrogen bonds build up a corrugated layer parallel to the (1 0 1) plane (Table 1).

S2. Experimental

A mixture of 1,5-dimethyl-2-(nitromethylene)-1,3,5-triazinane (20.64 g, 0.12 mol), potassium carbonate (20.7 g, 0.15 mol), potassium iodization (2 g) and ethyl acetate (150 ml) was stirred and heated in a 500 ml flask. The mixture was slowly heated to 353 K - 363 K and kept for 1 h. Then, 1-chloro-4-(1-chloroethyl)benzene (21 g, 0.12 mol, dissolved in 100 ml of ethyl acetate) was added dropwise into the flask, and the mixture was further stirred at 353 K - 363 K for 15 h. After cooling, the precipitate was filtered, washed with ethyl acetate and water, and recrystallized from ethyl acetate to obtain flaxen powder. Yield: 86%.

S3. Refinement

All H atoms were placed in calculated positions, with C–H = 0.93–0.98 Å, and included in the final cycles of refinement using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aryl, methylene and $1.5U_{\text{eq}}(\text{C})$ for methyl H atoms.




Figure 1

View of the title compound (I) with the atoms labeling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as small spheres of arbitrary radii.

(*E*)-1-[1-(4-Chlorophenyl)ethyl]-3,5-dimethyl-*N*-nitro-1,3,5-triazinan-2-imine

Crystal data

$C_{13}H_{18}ClN_5O_2$

$M_r = 311.77$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 7.2483$ (14) Å

$b = 29.568$ (6) Å

$c = 7.2306$ (14) Å

$\beta = 108.75$ (3)°

$V = 1467.4$ (5) Å³

$Z = 4$

$F(000) = 656$

$D_x = 1.411$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 3278 reflections

$\theta = 1.4$ – 27.9 °

$\mu = 0.27$ mm⁻¹

$T = 113$ K

Needle, colourless

$0.20 \times 0.16 \times 0.12$ mm

Data collection

Rigaku Saturn CCD area-detector
diffractometer

Radiation source: rotating anode

Confocal monochromator

Detector resolution: 7.31 pixels mm⁻¹

ω and φ scans

Absorption correction: multi-scan

(*ABSCOR*; Higashi, 1995)

$T_{\min} = 0.947$, $T_{\max} = 0.968$

9753 measured reflections

2585 independent reflections

2303 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.039$

$\theta_{\max} = 25.0$ °, $\theta_{\min} = 1.4$ °

$h = -8 \rightarrow 8$

$k = -35 \rightarrow 34$

$l = -8 \rightarrow 8$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.041$

$wR(F^2) = 0.104$

$S = 1.08$

2585 reflections

193 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0488P)^2 + 0.7943P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.46 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.51 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.37472 (9)	0.56523 (2)	0.15716 (10)	0.0546 (2)
O1	-0.25418 (19)	0.29503 (4)	0.41241 (18)	0.0236 (3)
O2	-0.48276 (18)	0.33395 (5)	0.47562 (18)	0.0254 (3)
N1	-0.0516 (2)	0.36893 (5)	0.2166 (2)	0.0173 (3)
N2	-0.2858 (2)	0.32380 (5)	0.0044 (2)	0.0175 (3)
N3	0.0492 (2)	0.31375 (5)	0.0208 (2)	0.0181 (3)
N4	-0.3580 (2)	0.36095 (5)	0.2595 (2)	0.0197 (3)
N5	-0.3641 (2)	0.32910 (5)	0.3838 (2)	0.0181 (3)
C1	0.3035 (3)	0.44147 (6)	0.3422 (3)	0.0261 (4)
H1	0.3813	0.4153	0.3853	0.031*
C2	0.3844 (3)	0.47936 (7)	0.2862 (3)	0.0321 (5)
H2	0.5159	0.4789	0.2876	0.038*
C3	0.2725 (3)	0.51775 (7)	0.2283 (3)	0.0323 (5)
C4	0.0823 (3)	0.51896 (7)	0.2250 (3)	0.0326 (5)
H4	0.0071	0.5457	0.1871	0.039*
C5	0.0016 (3)	0.48064 (6)	0.2776 (3)	0.0265 (4)
H5	-0.1307	0.4813	0.2736	0.032*
C6	0.1093 (3)	0.44127 (6)	0.3362 (3)	0.0206 (4)
C7	0.0118 (3)	0.39993 (6)	0.3875 (3)	0.0200 (4)
H7	-0.1081	0.4105	0.4143	0.024*
C8	0.1367 (3)	0.37472 (6)	0.5690 (3)	0.0231 (4)
H8A	0.2520	0.3622	0.5446	0.035*
H8B	0.0605	0.3501	0.5991	0.035*
H8C	0.1776	0.3957	0.6797	0.035*
C9	-0.2273 (2)	0.34984 (5)	0.1609 (2)	0.0166 (4)
C10	-0.1503 (3)	0.31248 (6)	-0.1043 (3)	0.0199 (4)
H10A	-0.1808	0.2819	-0.1616	0.024*
H10B	-0.1683	0.3343	-0.2127	0.024*
C11	0.0888 (3)	0.35748 (6)	0.1119 (3)	0.0184 (4)
H11A	0.0810	0.3807	0.0110	0.022*

H11B	0.2229	0.3578	0.2056	0.022*
C12	0.0994 (3)	0.27654 (6)	0.1633 (3)	0.0231 (4)
H12A	0.0757	0.2475	0.0940	0.035*
H12B	0.0188	0.2785	0.2489	0.035*
H12C	0.2373	0.2787	0.2420	0.035*
C13	-0.4771 (3)	0.30192 (6)	-0.0593 (3)	0.0226 (4)
H13A	-0.5701	0.3204	-0.0193	0.034*
H13B	-0.4672	0.2719	0.0007	0.034*
H13C	-0.5222	0.2988	-0.2018	0.034*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0535 (4)	0.0359 (3)	0.0547 (4)	-0.0206 (3)	-0.0102 (3)	0.0202 (3)
O1	0.0254 (7)	0.0221 (7)	0.0255 (7)	0.0071 (5)	0.0111 (6)	0.0057 (5)
O2	0.0188 (7)	0.0410 (8)	0.0203 (7)	0.0038 (6)	0.0120 (5)	0.0028 (6)
N1	0.0174 (8)	0.0186 (7)	0.0171 (7)	0.0002 (6)	0.0071 (6)	-0.0011 (6)
N2	0.0142 (7)	0.0210 (8)	0.0176 (7)	0.0000 (6)	0.0057 (6)	0.0004 (6)
N3	0.0165 (8)	0.0198 (8)	0.0185 (7)	-0.0008 (6)	0.0062 (6)	-0.0019 (6)
N4	0.0190 (8)	0.0220 (8)	0.0207 (8)	0.0039 (6)	0.0100 (6)	0.0035 (6)
N5	0.0151 (7)	0.0233 (8)	0.0155 (7)	0.0002 (6)	0.0045 (6)	-0.0007 (6)
C1	0.0241 (10)	0.0198 (9)	0.0300 (10)	-0.0014 (7)	0.0026 (8)	0.0005 (8)
C2	0.0260 (11)	0.0310 (11)	0.0346 (11)	-0.0077 (8)	0.0033 (9)	0.0021 (9)
C3	0.0390 (12)	0.0229 (10)	0.0254 (10)	-0.0100 (9)	-0.0032 (9)	0.0030 (8)
C4	0.0439 (13)	0.0204 (10)	0.0262 (11)	0.0063 (9)	0.0012 (9)	0.0000 (8)
C5	0.0286 (10)	0.0269 (10)	0.0223 (10)	0.0049 (8)	0.0059 (8)	-0.0039 (8)
C6	0.0249 (10)	0.0188 (9)	0.0164 (9)	-0.0012 (7)	0.0044 (7)	-0.0038 (7)
C7	0.0206 (9)	0.0214 (9)	0.0184 (9)	0.0015 (7)	0.0067 (7)	-0.0046 (7)
C8	0.0247 (10)	0.0262 (10)	0.0189 (9)	-0.0005 (8)	0.0075 (8)	-0.0005 (7)
C9	0.0175 (9)	0.0155 (8)	0.0165 (8)	0.0039 (7)	0.0052 (7)	0.0038 (7)
C10	0.0196 (9)	0.0253 (10)	0.0159 (8)	-0.0008 (7)	0.0073 (7)	-0.0021 (7)
C11	0.0175 (9)	0.0204 (9)	0.0195 (9)	-0.0012 (7)	0.0088 (7)	-0.0010 (7)
C12	0.0223 (10)	0.0215 (9)	0.0250 (10)	0.0027 (7)	0.0070 (8)	0.0004 (7)
C13	0.0146 (9)	0.0290 (10)	0.0232 (9)	-0.0024 (7)	0.0045 (7)	-0.0022 (8)

Geometric parameters (Å, °)

C11—C3	1.740 (2)	C4—C5	1.383 (3)
O1—N5	1.2595 (19)	C4—H4	0.9500
O2—N5	1.2522 (19)	C5—C6	1.390 (3)
N1—C9	1.332 (2)	C5—H5	0.9500
N1—C7	1.488 (2)	C6—C7	1.516 (3)
N1—C11	1.490 (2)	C7—C8	1.527 (3)
N2—C9	1.321 (2)	C7—H7	1.0000
N2—C13	1.464 (2)	C8—H8A	0.9800
N2—C10	1.480 (2)	C8—H8B	0.9800
N3—C11	1.437 (2)	C8—H8C	0.9800
N3—C10	1.439 (2)	C10—H10A	0.9900

N3—C12	1.471 (2)	C10—H10B	0.9900
N4—N5	1.312 (2)	C11—H11A	0.9900
N4—C9	1.395 (2)	C11—H11B	0.9900
C1—C2	1.384 (3)	C12—H12A	0.9800
C1—C6	1.394 (3)	C12—H12B	0.9800
C1—H1	0.9500	C12—H12C	0.9800
C2—C3	1.379 (3)	C13—H13A	0.9800
C2—H2	0.9500	C13—H13B	0.9800
C3—C4	1.372 (3)	C13—H13C	0.9800
C9—N1—C7	121.49 (14)	C6—C7—H7	107.3
C9—N1—C11	119.58 (14)	C8—C7—H7	107.3
C7—N1—C11	118.87 (13)	C7—C8—H8A	109.5
C9—N2—C13	122.48 (15)	C7—C8—H8B	109.5
C9—N2—C10	120.15 (14)	H8A—C8—H8B	109.5
C13—N2—C10	117.16 (14)	C7—C8—H8C	109.5
C11—N3—C10	108.91 (14)	H8A—C8—H8C	109.5
C11—N3—C12	112.59 (14)	H8B—C8—H8C	109.5
C10—N3—C12	113.18 (14)	N2—C9—N1	121.19 (15)
N5—N4—C9	111.03 (14)	N2—C9—N4	119.45 (15)
O2—N5—O1	120.71 (14)	N1—C9—N4	119.10 (15)
O2—N5—N4	117.46 (14)	N3—C10—N2	111.35 (14)
O1—N5—N4	121.83 (14)	N3—C10—H10A	109.4
C2—C1—C6	120.84 (18)	N2—C10—H10A	109.4
C2—C1—H1	119.6	N3—C10—H10B	109.4
C6—C1—H1	119.6	N2—C10—H10B	109.4
C3—C2—C1	119.5 (2)	H10A—C10—H10B	108.0
C3—C2—H2	120.3	N3—C11—N1	111.63 (14)
C1—C2—H2	120.3	N3—C11—H11A	109.3
C4—C3—C2	121.07 (19)	N1—C11—H11A	109.3
C4—C3—C11	119.65 (16)	N3—C11—H11B	109.3
C2—C3—C11	119.28 (18)	N1—C11—H11B	109.3
C3—C4—C5	119.04 (18)	H11A—C11—H11B	108.0
C3—C4—H4	120.5	N3—C12—H12A	109.5
C5—C4—H4	120.5	N3—C12—H12B	109.5
C4—C5—C6	121.61 (19)	H12A—C12—H12B	109.5
C4—C5—H5	119.2	N3—C12—H12C	109.5
C6—C5—H5	119.2	H12A—C12—H12C	109.5
C5—C6—C1	117.94 (17)	H12B—C12—H12C	109.5
C5—C6—C7	119.32 (17)	N2—C13—H13A	109.5
C1—C6—C7	122.73 (16)	N2—C13—H13B	109.5
N1—C7—C6	109.73 (14)	H13A—C13—H13B	109.5
N1—C7—C8	110.65 (14)	N2—C13—H13C	109.5
C6—C7—C8	114.18 (15)	H13A—C13—H13C	109.5
N1—C7—H7	107.3	H13B—C13—H13C	109.5

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C8—H8 <i>A</i> \cdots O2 ⁱ	0.98	2.29	3.273 (2)	178
C10—H10 <i>A</i> \cdots O1 ⁱⁱ	0.99	2.43	3.278 (2)	143
C11—H11 <i>B</i> \cdots O2 ⁱ	0.99	2.49	3.434 (3)	160

Symmetry codes: (i) $x+1, y, z$; (ii) $x, -y+1/2, z-1/2$.