

Expression, purification, crystallization and preliminary X-ray diffraction analysis of galactokinase from *Pyrococcus horikoshii*. Erratum

Eiji Inagaki,^a Keiko Sakamoto,^a Naomi Obayashi,^b Takaho Terada,^b Mikako Shirouzu,^b Yoshitaka Bessho,^b Chizu Kuroishi,^a Seiki Kuramitsu,^{a,c} Akeo Shinkai^a and Shigeyuki Yokoyama^{a,b,d*}

^aRIKEN SPring-8 Center, Harima Institute, 1-1-1 Kouto, Sayo, Hyogo 679-5148, Japan, ^bRIKEN Genomic Sciences Center, 1-7-22 Suehiro-cho, Tsurumi, Yokohama 230-0045, Japan, ^cDepartment of Biology, Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan, and ^dDepartment of Biophysics and Biochemistry, Graduate School of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan. Correspondence e-mail: yokoyama@biochem.s.u-tokyo.ac.jp

An error is corrected in the paper by Inagaki *et al.* [(2006), *Acta Cryst.* **F62**, 169–171].

This erratum refers to the paper by Inagaki *et al.* [(2006), *Acta Cryst.* **F62**, 169–171]. In §2.3 the amount of silicone and paraffin oil mixture used in the crystallization of the apo form of the enzyme was given incorrectly. The correct amount is 15 µl.

References

Inagaki, E., Sakamoto, K., Obayashi, N., Terada, T., Shirouzu, M., Bessho, Y., Kuroishi, C., Kuramitsu, S., Shinkai, A. & Yokoyama, S. (2006). *Acta Cryst.* **F62**, 169–171.