

Deer mouse hemoglobin exhibits a lowered oxygen affinity owing to mobility of the E helix. Corrigendum

Noriko Inoguchi,^a Jake R. Oshlo,^b Chandrasekhar Natarajan,^a Roy E. Weber,^c Angela Fago,^c Jay F. Storz^a and Hideaki Moriyama^{a*}

^aSchool of Biological Sciences, University of Nebraska-Lincoln, Lincoln, Nebraska, USA, ^bNebraska Wesleyan University, Lincoln, Nebraska, USA, and ^cZoophysiology, Department of Bioscience, Aarhus University, Aarhus, Denmark

Correspondence e-mail: hmoriyama2@unl.edu

The affiliation of two authors in the article by Inoguchi *et al.* [(2013). *Acta Cryst.* **F69**, 393–398] is corrected.

In the article by Inoguchi *et al.* (2013) the affiliation for two of the authors, Angela Fago and Roy E. Weber, was given incorrectly. The correct affiliation is Zoophysiology, Department of Bioscience, Aarhus University, Aarhus, Denmark.

References

Inoguchi, N., Oshlo, J. R., Natarajan, C., Weber, R. E., Fago, A., Storz, J. F. & Moriyama, H. (2013). *Acta Cryst.* **F69**, 393–398.