

Pentacarbonyl(*N,N*-dimethylbenzylamine)-tungstenTariq Mahmud,^a Javed Iqbal,^a
Muhammad Irshad,^a Mark R. J.
Elsegood^b and Vickie McKee^{b*}^aInstitute of Chemistry, University of the Punjab,
Lahore 54590, Pakistan, and ^bChemistry
Department, Loughborough University,
Loughborough, Leicestershire LE11 3TU,
England

Correspondence e-mail: v.mckee@lboro.ac.uk

Key indicators

Single-crystal X-ray study
T = 150 K
Mean $\sigma(\text{C}-\text{C}) = 0.011 \text{ \AA}$
R factor = 0.033
wR factor = 0.074
Data-to-parameter ratio = 20.2For details of how these key indicators were
automatically derived from the article, see
<http://journals.iucr.org/e>.

The title compound, $[\text{W}(\text{C}_9\text{H}_{13}\text{N})(\text{CO})_5]$, was prepared by irradiation of $\text{W}(\text{CO})_6$ in tetrahydrofuran in the presence of *N,N*-dimethylbenzylamine. The geometry at the W atom is approximately octahedral, with the *cis* bond angles in the range $86.3(3)$ – $95.6(2)^\circ$. The bond to the tertiary amine is long [$2.371(5) \text{ \AA}$] and, as might be expected, the bond to the *trans* carbonyl is quite short [$\text{W}-\text{C} = 1.964(7) \text{ \AA}$]. The remaining $\text{W}-\text{CO}$ bonds lie in the range $2.033(6)$ – $2.049(6) \text{ \AA}$. Similar bonding patterns have been observed in related $\text{W}(\text{CO})_5(\text{amine})$ complexes,



Received 8 August 2005

Accepted 10 August 2005


Online 17 August 2005

Comment

(*N,N*-Dimethylbenzylamine)pentacarbonyltungsten was prepared by irradiation of $\text{W}(\text{CO})_6$ in tetrahydrofuran (THF) in the presence of the amine. Presumably, the reaction proceeds via an intermediate THF complex (Aroney *et al.*, 1994, and references therein):



Although a number of cyclometallated complexes of tungsten with this ligand have been reported previously (van der Schaaf *et al.*, 1993), no carbonyl complex has been structurally characterized. Also, in our hands, no cyclometallated complex was isolated.



The structure of (*N,N*-dimethylbenzylamine)pentacarbonyltungsten, (I), is shown in Fig. 1. The geometry at the W atom is approximately octahedral, with the *cis* bond angles in the range $86.3(3)$ – $95.6(2)^\circ$. The bond to the tertiary amine is long [$2.371(5) \text{ \AA}$] and, as might be expected, the bond to the *trans* carbonyl is quite short [$\text{W1}-\text{C10} = 1.964(7) \text{ \AA}$]. The remaining $\text{W}-\text{CO}$ bonds lie in the range $2.033(6)$ –




Figure 1
Perspective view of the complex, with displacement ellipsoids drawn at the 50% probability level.

2.049 (6) Å. Similar bonding patterns have been observed in related $W(CO)_5(\text{amine})$ complexes [see, for example, Long *et al.* (2002) and Moralejo *et al.* (1991)]. There are no obvious π - π or edge-to-face interactions.

Experimental

$W(CO)_6$ (0.351 g, 1.0 mmol) and *N,N*-dimethylbenzylamine (0.30 ml, 2.0 mmol) were dissolved in sodium-dried THF (20 ml). The mixture was stirred under N_2 and irradiated with UV light for 4 h, yielding a yellow solution. The progress of the reaction was monitored by following the CO stretching band at 1975 cm^{-1} by IR. The volume of the solvent was reduced under vacuum and n-hexane added to induce crystallization (yield 0.078 g, 17%). The sample was not pure and did not give satisfactory microanalysis. The EI mass spectrum of the complex showed a cluster corresponding to the parent ion $W(C_6H_5CH_2N(CH_3)_2)(CO)_5$ centered at m/e 459 and the isotope pattern matched that predicted from theory. Clusters corresponding to sequential loss of CO groups were observed at m/e of 431 [$W(C_6H_5CH_2N(CH_3)_2)(CO)_4$], 403 [$W(C_6H_5CH_2N(CH_3)_2)(CO)_3$] and 375 [$W(C_6H_5CH_2N(CH_3)_2)(CO)_2$]. Clusters at m/e 345, 317 and 135 were assigned to $W(C_6H_5CH_2N)(CO)_2$, $W(C_6H_5CH_2N)(CO)$ and [$C_6H_5CH_2N(CH_3)_2 + H^+$], respectively. Clusters corresponding to $W(CO)_6$, $W(CO)_5$, $W(CO)_4$, $W(CO)_3$, $W(CO)_2$, $W(CO)$ and W were also observed. The $W(CO)_6$ was most likely present as an impurity in the sample. 1H NMR ($CDCl_3$): 2.78 (s, 6H, CH_3), 4.22 (s, 2H, CH_2), 7.25–7.34 (m, 5H, aromatic). ^{13}C NMR: 55.3 (CH_3), 73.6 (CH_2), 128.5 (aromatic C_3 , C_5), 129.0 (aromatic C_4), 132.0 (aromatic, C_2 , C_6), 191, 199, 202 (carbonyl). IR (KBr, cm^{-1}): 3425 (m), 1952 (m), 1060 (w), 932 (m), 853 (m) 774 (m), 592 (s).

Crystal data

$[W(C_9H_{13}N)(CO)_5]$
 $M_r = 459.10$
Orthorhombic, $Pbca$
 $a = 13.7829$ (11) Å
 $b = 12.5247$ (10) Å
 $c = 18.2985$ (14) Å
 $V = 3158.8$ (4) Å³
 $Z = 8$
 $D_x = 1.931\text{ Mg m}^{-3}$

Mo $K\alpha$ radiation
Cell parameters from 4675 reflections
 $\theta = 2.5$ – 28.1°
 $\mu = 7.33\text{ mm}^{-1}$
 $T = 150$ (2) K
Plate, yellow
 $0.28 \times 0.20 \times 0.05\text{ mm}$

Data collection

Bruker SMART 1000 CCD diffractometer
 ω rotation with narrow-frame scans
Absorption correction: multi-scan (SADABS; Sheldrick, 2003)
 $T_{\min} = 0.233$, $T_{\max} = 0.711$
18147 measured reflections

3870 independent reflections
2534 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.059$
 $\theta_{\max} = 29.0^\circ$
 $h = -13 \rightarrow 18$
 $k = -16 \rightarrow 16$
 $l = -24 \rightarrow 22$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.033$
 $wR(F^2) = 0.074$
 $S = 1.03$
3870 reflections
192 parameters
H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0125P)^2 + 14.4005P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 1.10\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.85\text{ e \AA}^{-3}$

Table 1

Selected geometric parameters (Å, °).

W1–N1	2.371 (5)	O11–C11	1.144 (8)
W1–C10	1.964 (7)	O12–C12	1.152 (8)
W1–C11	2.041 (6)	O13–C13	1.156 (7)
W1–C12	2.034 (6)	O14–C14	1.142 (7)
W1–C13	2.033 (6)	N1–C1	1.492 (9)
W1–C14	2.048 (6)	N1–C2	1.496 (9)
O10–C10	1.157 (9)	N1–C3	1.514 (8)
N1–W1–C10	176.8 (2)	C13–W1–C14	172.6 (3)
N1–W1–C11	93.9 (2)	W1–N1–C1	110.0 (4)
N1–W1–C12	89.8 (2)	W1–N1–C2	111.7 (4)
N1–W1–C13	91.5 (2)	W1–N1–C3	109.1 (4)
N1–W1–C14	95.6 (2)	C1–N1–C2	107.4 (5)
C10–W1–C11	88.4 (3)	C1–N1–C3	109.3 (5)
C10–W1–C12	88.0 (3)	C2–N1–C3	109.3 (5)
C10–W1–C13	86.3 (3)	N1–C3–C4	115.8 (6)
C10–W1–C14	86.7 (3)	W1–C10–O10	177.2 (6)
C11–W1–C12	174.8 (3)	W1–C11–O11	176.4 (6)
C11–W1–C13	90.5 (2)	W1–C12–O12	174.0 (6)
C11–W1–C14	86.8 (2)	W1–C13–O13	173.9 (6)
C12–W1–C13	93.1 (3)	W1–C14–O14	174.5 (5)
C12–W1–C14	89.1 (3)		

H atoms bonded to C atoms were inserted at calculated positions and refined using a riding model. The constrained C–H distances were 0.95, 0.98 and 0.99 Å for aryl, methyl, and methylene H atoms, respectively. The H atoms of methylene and aryl groups were refined with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ and those of the methyl groups with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$. The highest residual electron-density peak is 0.88 Å⁻³ from the W atom.

Data collection: SMART (Bruker, 1998); cell refinement: SAINT (Bruker, 1998); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2001); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and local programs.

The authors thank the University of the Punjab, Lahore, Pakistan, and the Charles Wallace Pakistan Trust for financial support, and are also grateful to Loughborough University for providing facilities and to the EPSRC Mass Spectrometry Service Centre, University of Wales, Swansea, for the mass spectrum.

References

- Aroney, M. J., Buys, I. E., Davies, M. S. & Hambley, T. W. (1994). *J. Chem. Soc. Dalton Trans.* pp. 2827–2834.
- Bruker (1998). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Long, N. J., Parker, D. G., Speyer, P. R., White, A. J. P. & Williams, D. J. (2002). *J. Chem. Soc. Dalton Trans.* pp. 2142–2150.
- Moralejo, C., Langford, C. H. & Bird, P. H. (1991). *Can. J. Chem.* **69**, 2033–2037.
- Schaaf, P. A. van der , Grove, D. M., Smeets, W. J. J., Spek, A. L. & van Koten, G. (1993) *Organometallics*, **12**, 3955–3963.
- Sheldrick, G. M. (2001). *SHELXTL*. Version 6.12. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2003). *SADABS*. Version 2.10. Bruker AXS Inc., Madison, Wisconsin, USA.

supporting information

Acta Cryst. (2005). E61, m1789–m1791 [https://doi.org/10.1107/S1600536805025560]

Pentacarbonyl(*N,N*-dimethylbenzylamine)tungsten

Tariq Mahmud, Javed Iqbal, Muhammad Irshad, Mark R. J. Elsegood and Vickie McKee

(N,N-dimethylbenzylamine)pentacarbonyltungsten*Crystal data*

[W(C₉H₁₃N)(CO)₅]

$M_r = 459.10$

Orthorhombic, *Pbca*

Hall symbol: -P 2ac 2ab

$a = 13.7829$ (11) Å

$b = 12.5247$ (10) Å

$c = 18.2985$ (14) Å

$V = 3158.8$ (4) Å³

$Z = 8$

$F(000) = 1744$

$D_x = 1.931$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4675 reflections

$\theta = 2.5$ – 28.1°

$\mu = 7.33$ mm⁻¹

$T = 150$ K

Plate, yellow

$0.28 \times 0.20 \times 0.05$ mm

Data collection

Bruker SMART 1000 CCD
diffractometer

Radiation source: sealed tube

Graphite monochromator

ω rotation with narrow frames scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 2003)

$T_{\min} = 0.233$, $T_{\max} = 0.711$

18147 measured reflections

3870 independent reflections

2534 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.059$

$\theta_{\max} = 29.0^\circ$, $\theta_{\min} = 2.2^\circ$

$h = -13 \rightarrow 18$

$k = -16 \rightarrow 16$

$l = -24 \rightarrow 22$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.033$

$wR(F^2) = 0.074$

$S = 1.03$

3870 reflections

192 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0125P)^2 + 14.4005P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 1.10$ e Å⁻³

$\Delta\rho_{\min} = -0.85$ e Å⁻³

Special details

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
W1	0.07804 (2)	0.77665 (2)	0.20210 (1)	0.0237 (1)
O10	0.0153 (4)	0.8367 (4)	0.0435 (3)	0.0530 (19)
O11	0.1652 (3)	1.0111 (4)	0.2164 (3)	0.0460 (18)
O12	0.0043 (4)	0.5425 (4)	0.1638 (3)	0.050 (2)
O13	-0.1313 (3)	0.8659 (4)	0.2432 (3)	0.0437 (17)
O14	0.2815 (3)	0.7141 (4)	0.1313 (3)	0.0410 (17)
N1	0.1143 (4)	0.7259 (4)	0.3239 (3)	0.0270 (14)
C1	0.0264 (5)	0.6796 (7)	0.3594 (4)	0.048 (3)
C2	0.1456 (6)	0.8194 (5)	0.3690 (4)	0.046 (3)
C3	0.1944 (5)	0.6432 (5)	0.3230 (4)	0.0313 (19)
C4	0.2273 (5)	0.6028 (5)	0.3973 (4)	0.032 (2)
C5	0.3071 (5)	0.6478 (6)	0.4319 (4)	0.041 (3)
C6	0.3377 (6)	0.6098 (7)	0.4996 (5)	0.056 (3)
C7	0.2906 (6)	0.5259 (7)	0.5324 (4)	0.055 (3)
C8	0.2133 (6)	0.4800 (6)	0.4991 (5)	0.051 (3)
C9	0.1813 (5)	0.5179 (5)	0.4313 (4)	0.039 (2)
C10	0.0410 (5)	0.8148 (5)	0.1018 (4)	0.0323 (19)
C11	0.1338 (4)	0.9266 (5)	0.2138 (4)	0.032 (2)
C12	0.0289 (5)	0.6265 (5)	0.1815 (4)	0.0330 (19)
C13	-0.0554 (4)	0.8303 (5)	0.2320 (4)	0.032 (2)
C14	0.2101 (4)	0.7340 (5)	0.1599 (3)	0.0283 (19)
H1A	0.00200	0.62020	0.32980	0.0720*
H1B	-0.02380	0.73470	0.36340	0.0720*
H1C	0.04330	0.65360	0.40830	0.0720*
H2A	0.16320	0.79500	0.41810	0.0680*
H2B	0.09240	0.87090	0.37230	0.0680*
H2C	0.20190	0.85340	0.34600	0.0680*
H3A	0.25120	0.67420	0.29760	0.0380*
H3B	0.17200	0.58130	0.29380	0.0380*
H5	0.34110	0.70490	0.40920	0.0500*
H6	0.39160	0.64210	0.52330	0.0670*
H7	0.31220	0.50010	0.57850	0.0660*
H8	0.18070	0.42200	0.52190	0.0620*
H9	0.12730	0.48490	0.40820	0.0470*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
W1	0.0238 (1)	0.0235 (1)	0.0238 (1)	-0.0016 (1)	0.0004 (1)	0.0004 (1)
O10	0.055 (3)	0.068 (4)	0.036 (3)	-0.005 (3)	-0.006 (3)	0.013 (3)
O11	0.048 (3)	0.028 (2)	0.062 (4)	-0.008 (2)	-0.003 (3)	0.002 (2)
O12	0.064 (4)	0.028 (3)	0.057 (4)	-0.014 (2)	-0.007 (3)	-0.007 (2)
O13	0.029 (3)	0.047 (3)	0.055 (3)	0.005 (2)	0.003 (2)	0.005 (3)
O14	0.034 (3)	0.047 (3)	0.042 (3)	0.005 (2)	0.011 (2)	-0.002 (3)
N1	0.026 (2)	0.028 (2)	0.027 (3)	0.001 (2)	0.001 (2)	0.000 (3)

C1	0.032 (4)	0.075 (5)	0.037 (5)	0.001 (4)	0.006 (3)	0.018 (4)
C2	0.066 (5)	0.034 (4)	0.038 (4)	0.017 (4)	-0.012 (4)	-0.015 (3)
C3	0.033 (3)	0.027 (3)	0.034 (4)	0.006 (3)	0.002 (3)	-0.005 (3)
C4	0.038 (4)	0.030 (3)	0.027 (4)	0.012 (3)	-0.001 (3)	0.001 (3)
C5	0.035 (4)	0.053 (5)	0.036 (4)	0.005 (3)	0.002 (3)	0.004 (4)
C6	0.049 (5)	0.074 (6)	0.044 (5)	0.011 (4)	-0.014 (4)	-0.007 (5)
C7	0.063 (6)	0.069 (6)	0.032 (5)	0.026 (5)	-0.004 (4)	0.011 (4)
C8	0.064 (5)	0.047 (4)	0.042 (5)	0.010 (4)	0.006 (4)	0.016 (4)
C9	0.048 (4)	0.026 (3)	0.043 (5)	0.001 (3)	-0.006 (4)	0.005 (3)
C10	0.029 (3)	0.032 (3)	0.036 (4)	-0.009 (3)	0.004 (3)	0.005 (3)
C11	0.029 (3)	0.037 (4)	0.031 (4)	0.003 (3)	0.004 (3)	-0.005 (3)
C12	0.031 (3)	0.034 (3)	0.034 (4)	-0.001 (3)	0.001 (3)	0.008 (3)
C13	0.030 (4)	0.033 (3)	0.032 (4)	-0.007 (3)	0.001 (3)	0.004 (3)
C14	0.033 (3)	0.025 (3)	0.027 (4)	-0.002 (3)	-0.002 (3)	-0.003 (3)

Geometric parameters (Å, °)

W1—N1	2.371 (5)	C5—C6	1.393 (12)
W1—C10	1.964 (7)	C6—C7	1.373 (12)
W1—C11	2.041 (6)	C7—C8	1.355 (12)
W1—C12	2.034 (6)	C8—C9	1.400 (11)
W1—C13	2.033 (6)	C1—H1A	0.9800
W1—C14	2.048 (6)	C1—H1B	0.9800
O10—C10	1.157 (9)	C1—H1C	0.9800
O11—C11	1.144 (8)	C2—H2A	0.9800
O12—C12	1.152 (8)	C2—H2B	0.9800
O13—C13	1.156 (7)	C2—H2C	0.9800
O14—C14	1.142 (7)	C3—H3A	0.9900
N1—C1	1.492 (9)	C3—H3B	0.9900
N1—C2	1.496 (9)	C5—H5	0.9500
N1—C3	1.514 (8)	C6—H6	0.9500
C3—C4	1.520 (10)	C7—H7	0.9500
C4—C5	1.389 (10)	C8—H8	0.9500
C4—C9	1.386 (9)	C9—H9	0.9500
N1—W1—C10	176.8 (2)	W1—C11—O11	176.4 (6)
N1—W1—C11	93.9 (2)	W1—C12—O12	174.0 (6)
N1—W1—C12	89.8 (2)	W1—C13—O13	173.9 (6)
N1—W1—C13	91.5 (2)	W1—C14—O14	174.5 (5)
N1—W1—C14	95.6 (2)	N1—C1—H1A	109.00
C10—W1—C11	88.4 (3)	N1—C1—H1B	109.00
C10—W1—C12	88.0 (3)	N1—C1—H1C	109.00
C10—W1—C13	86.3 (3)	H1A—C1—H1B	109.00
C10—W1—C14	86.7 (3)	H1A—C1—H1C	110.00
C11—W1—C12	174.8 (3)	H1B—C1—H1C	109.00
C11—W1—C13	90.5 (2)	N1—C2—H2A	109.00
C11—W1—C14	86.8 (2)	N1—C2—H2B	110.00
C12—W1—C13	93.1 (3)	N1—C2—H2C	109.00

C12—W1—C14	89.1 (3)	H2A—C2—H2B	110.00
C13—W1—C14	172.6 (3)	H2A—C2—H2C	109.00
W1—N1—C1	110.0 (4)	H2B—C2—H2C	109.00
W1—N1—C2	111.7 (4)	N1—C3—H3A	108.00
W1—N1—C3	109.1 (4)	N1—C3—H3B	108.00
C1—N1—C2	107.4 (5)	C4—C3—H3A	108.00
C1—N1—C3	109.3 (5)	C4—C3—H3B	108.00
C2—N1—C3	109.3 (5)	H3A—C3—H3B	107.00
N1—C3—C4	115.8 (6)	C4—C5—H5	120.00
C3—C4—C5	120.6 (6)	C6—C5—H5	120.00
C3—C4—C9	121.4 (6)	C5—C6—H6	120.00
C5—C4—C9	118.0 (7)	C7—C6—H6	120.00
C4—C5—C6	120.5 (7)	C6—C7—H7	120.00
C5—C6—C7	120.5 (8)	C8—C7—H7	120.00
C6—C7—C8	120.0 (8)	C7—C8—H8	120.00
C7—C8—C9	120.2 (7)	C9—C8—H8	120.00
C4—C9—C8	120.9 (7)	C4—C9—H9	119.00
W1—C10—O10	177.2 (6)	C8—C9—H9	120.00
C11—W1—N1—C1	129.4 (5)	C1—N1—C3—C4	-60.3 (7)
C11—W1—N1—C2	10.3 (5)	C2—N1—C3—C4	57.0 (7)
C11—W1—N1—C3	-110.7 (4)	N1—C3—C4—C5	-95.4 (8)
C12—W1—N1—C1	-54.3 (5)	N1—C3—C4—C9	87.0 (8)
C12—W1—N1—C2	-173.4 (5)	C3—C4—C9—C8	178.8 (7)
C12—W1—N1—C3	65.6 (4)	C5—C4—C9—C8	1.1 (10)
C13—W1—N1—C1	38.8 (5)	C9—C4—C5—C6	-1.6 (11)
C13—W1—N1—C2	-80.3 (5)	C3—C4—C5—C6	-179.3 (7)
C13—W1—N1—C3	158.7 (4)	C4—C5—C6—C7	1.3 (12)
C14—W1—N1—C1	-143.5 (5)	C5—C6—C7—C8	-0.5 (13)
C14—W1—N1—C2	97.5 (5)	C6—C7—C8—C9	0.0 (12)
C14—W1—N1—C3	-23.5 (4)	C7—C8—C9—C4	-0.4 (12)
W1—N1—C3—C4	179.4 (4)		
