

Propane-1,3-diaminium-2-carboxy-pyridine-6-carboxylate-pyridine-2,6-dicarboxylic acid-water (1/2/2/8)

Janet Soleimannejad,^a Hossein Aghabozorg,^{b*} Elham Motyeian,^b Mohammad Ghadermazi,^c Jafar Attar Gharamaleki^b and Harry Adams^d

^aFaculty of Science, Department of Chemistry, Ilam University, Iran, ^bFaculty of Chemistry, Teacher Training University, 49 Mofateh Avenue 15614, Tehran, Iran, ^cDepartment of Chemistry, Faculty of Science, University of Kurdistan, Sanandaj, Iran, and ^dDepartment of Chemistry, Sheffield University, Sheffield S3 7HF, England
Correspondence e-mail: haghbozorg@yahoo.com


Received 13 November 2007; accepted 3 December 2007

Key indicators: single-crystal X-ray study; $T = 150\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.051; wR factor = 0.139; data-to-parameter ratio = 16.4.

The title proton-transfer compound, $\text{C}_3\text{H}_{12}\text{N}_2^{2+} \cdot 2\text{C}_7\text{H}_4\text{NO}_4^- \cdots 2\text{C}_7\text{H}_5\text{NO}_4 \cdot 8\text{H}_2\text{O}$ or $(\text{pnH}_2)(\text{pydcH}_2)_2 \cdot 2(\text{pydcH}_2) \cdot 8\text{H}_2\text{O}$, was obtained by the reaction of pyridine-2,6-dicarboxylic acid (pydcH_2) and propane-1,3-diamine (pn) in aqueous solution. Both neutral and monoanionic forms of the diacid are observed in the crystal structure. The negative charge of two monoanions is balanced by the dicationic propane-1,3-diaminium species. In addition, considerable $\pi-\pi$ stacking interactions between the aromatic rings of the (pydcH_2)⁻ and (pydcH_2) fragments [with centroid–centroid distances of 3.5108 (11)–3.5949 (11) Å] are observed. The most important feature of this crystal structure is the presence of a large number of O–H···O, O–H···N, N–H···O, N–H···N, C–H···O and C–H···N hydrogen bonds, with $D \cdots A$ ranging from 2.445 (2) to 3.485 (3) Å. These interactions as well as ion pairing and $\pi-\pi$ stacking connect the various fragments into a supramolecular structure.

Related literature

For related literature, see: Aghabozorg, Attar Gharamaleki, Ghadermazi, Ghasemikhah & Soleimannejad (2007); Aghabozorg, Attar Gharamaleki, Ghasemikhah, Ghadermazi & Soleimannejad, 2007; Aghabozorg, Daneshvar *et al.* (2007).



Experimental

Crystal data

$\text{C}_3\text{H}_{12}\text{N}_2^{2+} \cdot 2\text{C}_7\text{H}_4\text{NO}_4^- \cdots 2\text{C}_7\text{H}_5\text{NO}_4 \cdot 8\text{H}_2\text{O}$	$\beta = 99.914(1)^\circ$
$M_r = 886.73$	$V = 3924.60(10)\text{ \AA}^3$
Monoclinic, $P2_1/c$	$Z = 4$
$a = 13.5425(2)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 13.5237(2)\text{ \AA}$	$\mu = 0.13\text{ mm}^{-1}$
$c = 21.7538(3)\text{ \AA}$	$T = 150(2)\text{ K}$
	$0.26 \times 0.12 \times 0.12\text{ mm}$

Data collection

Bruker SMART 1000 diffractometer	56174 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 1998)	9031 independent reflections
$T_{\min} = 0.967$, $T_{\max} = 0.985$	6702 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.054$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$	550 parameters
$wR(F^2) = 0.139$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\text{max}} = 0.84\text{ e \AA}^{-3}$
9031 reflections	$\Delta\rho_{\text{min}} = -0.51\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O2–H2A···O19	0.95	1.94	2.830 (2)	154
O5–H5A4A···O20	0.95	1.67	2.592 (2)	164
O7–H7A···O3	0.95	1.51	2.455 (2)	177
O7–H7A···O4	0.95	2.57	3.119 (2)	117
O9–H9A···O22	0.95	1.64	2.588 (2)	172
O11–H11A···O13 ⁱ	0.95	1.50	2.445 (2)	177
O11–H11A···O14 ⁱ	0.95	2.61	3.181 (2)	119
O16–H16A···O24	0.95	1.60	2.551 (2)	174
O17–H17A···O15	0.95	1.90	2.838 (2)	167
O17–H17A···N4	0.95	2.43	2.956 (2)	115
O17–H17B···O14	0.95	1.87	2.821 (2)	175
O18–H18B···O1	0.95	1.89	2.815 (2)	164
O18–H18A···O17 ⁱⁱ	0.95	1.88	2.809 (2)	166
O19–H19B···N1	0.95	2.22	3.092 (2)	152
O19–H19A···O13 ⁱ	0.95	2.34	3.174 (2)	146
O20–H20B···O12 ⁱⁱⁱ	0.95	1.87	2.793 (2)	164
O20–H20A···O22	0.95	1.95	2.875 (3)	165
O21–H21B···O10	0.95	1.89	2.834 (2)	170
O21–H21A···O19 ⁱⁱⁱ	0.95	1.85	2.784 (2)	166
O22–H22A···O21	0.95	1.88	2.818 (2)	168
O22–H22B···O17 ^{iv}	0.95	1.81	2.742 (2)	167
O23–H23B···O6	0.95	1.85	2.749 (2)	157
O23–H23A···O12 ⁱⁱⁱ	0.95	1.96	2.899 (2)	170
O24–H24B···O18 ^v	0.95	1.79	2.706 (2)	161
O24–H24A···O12 ^{vi}	0.95	1.86	2.761 (2)	157
N5–H5A···N2	0.95	2.40	3.342 (3)	171
N5–H5A···O6	0.95	2.61	3.227 (2)	123
N5–H5B···O4	0.95	1.95	2.879 (2)	166
N5–H5C···O23 ^{vii}	0.95	2.06	2.991 (3)	166
N6–H6A···O14 ⁱ	0.95	2.01	2.938 (2)	164
N6–H6B···O21 ^{viii}	0.95	1.90	2.849 (2)	174
N6–H6C···N3	0.95	2.17	3.070 (2)	158

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
C5—H5 \cdots O18 ⁱⁱⁱ	0.95	2.56	3.429 (3)	153
C10—H10 \cdots O1 ^{ix}	0.95	2.50	3.335 (3)	147
C11—H11 \cdots O18 ^{ix}	0.95	2.52	3.427 (3)	159
C19—H19 \cdots O15 ^x	0.95	2.46	3.266 (3)	143
C25—H25 \cdots O24 ^{xi}	0.95	2.56	3.485 (3)	165
C26—H26 \cdots O16 ^{xi}	0.95	2.56	3.287 (3)	134
C31—H31A \cdots N2	0.99	2.52	3.391 (3)	146
C31—H31B \cdots O10	0.99	2.54	3.175 (3)	122

Symmetry codes: (i) $-x+1, y+\frac{1}{2}, -z+\frac{1}{2}$; (ii) $-x+1, -y+1, -z+1$; (iii) $x, -y+\frac{3}{2}, z-\frac{1}{2}$; (iv) $x, -y+\frac{1}{2}, z-\frac{1}{2}$; (v) $x-1, -y+\frac{1}{2}, z-\frac{1}{2}$; (vi) $-x, y-\frac{1}{2}, -z+\frac{1}{2}$; (vii) $-x+1, -y+2, -z$; (viii) $-x+1, -y+1, -z$; (ix) $x-1, -y+\frac{3}{2}, z-\frac{1}{2}$; (x) $-x, y+\frac{1}{2}, -z+\frac{1}{2}$; (xi) $-x, -y, -z$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1998); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL* (Bruker, 2005); software used to prepare material for publication: *SHELXTL*.

Financial support from Ilam University and the Teacher Training University is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2048).

References

- Aghabozorg, H., Attar Gharamaleki, J., Ghadermazi, M., Ghasemikhah, P. & Soleimannejad, J. (2007). *Acta Cryst. E63*, m1803–m1804.
- Aghabozorg, H., Attar Gharamaleki, J., Ghasemikhah, P., Ghadermazi, M. & Soleimannejad, J. (2007). *Acta Cryst. E63*, m1710–m1711.
- Aghabozorg, H., Daneshvar, S., Motyeian, E., Ghadermazi, M. & Attar Gharamaleki, J. (2007). *Acta Cryst. E63*, m2468–m2469.
- Bruker (1998). *SMART* (Version 5.0), *SAINT* (Version 4.0) and *SADABS* (Version 2.0). Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2005). *SHELXTL*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.

supporting information

Acta Cryst. (2008). E64, o231–o232 [https://doi.org/10.1107/S1600536807065270]

Propane-1,3-diaminium-2-carboxypyridine-6-carboxylate-pyridine-2,6-dicarboxylic acid-water (1/2/2/8)

Janet Soleimannejad, Hossein Aghabozorg, Elham Motyeian, Mohammad Ghadermazi, Jafar Attar Gharamaleki and Harry Adams

S1. Comment

Intermolecular interactions, such as hydrogen bonding, π - π stacking, ion pairing and donor-acceptor interactions, are famous for making aggregates of molecules. Hydrogen bonding has been described as the most important interaction in supramolecular chemistry. Dicarboxylic acids possess a good potential to be used as proton donors in the synthesis of proton transfer compounds. Among these diacids, pyridine-2,6-dicarboxylic acid has been used by our research group for preparing of such compounds. For example, (pydaH)(pydcH) in which pyridine-2,6-diamine (pyda) was used as a proton acceptor. The resulting compounds with some remaining sites as electron donors can coordinate to metallic ions (Aghabozorg, Attar Gharamaleki, Ghadermazi, Ghasemikhah & Soleimannejad, 2007; Aghabozorg, Attar Gharamaleki, Ghasemikhah, Ghadermazi & Soleimannejad, 2007; Aghabozorg, Daneshvar *et al.*, 2007).


Here we report a new proton transfer compound obtained from (pydcH₂) as a proton donor and propane-1,3-diamine (pn) as an acceptor. The molecular structure of the title compound is shown in Fig. 1. The intermolecular hydrogen bond distances are listed in Table 1.

The structure of this compound contains two neutral pydcH₂ molecules, two monoanionic (pydcH)⁻, one (pnH₂)²⁺ species and eight uncoordinated water molecules. The negative charge of two monoanions is neutralized by dicationic propane-1,3-diaminium fragments.


A considerable π - π stacking interactions between aromatic rings of (pydcH₂) and (pydcH)⁻ fragments with centroid-centroid distances of 3.5108 (11)–3.5949 (11) Å are observed in the prepared compound (Fig. 2). A remarkable feature in the crystal structure of compound (I) is the presence of a large number of O—H···O, O—H···N, N—H···O, N—H···N, C—H···O and C—H···N hydrogen bonds interactions ranging from 2.450 (2) to 3.488 (3) Å. The shortest hydrogen bond is O11—H11A···O13 (-x + 1, y + 1/2, -z + 1/2) with D···A = 2.450 (2) Å, a very strong interaction (Table 1). These extensive hydrogen bond interactions as well as ion pairing and π - π stacking connect the different components to form a three-dimensional supramolecular structure (Fig. 3).

S2. Experimental


The reaction of pyridine-2,6-dicarboxylic acid (pydcH₂) (330 mg, 2 mmol) with propane-1,3-diamine (pn) (76 mg, 1 mmol) in a 2:1 molar ratio in tetrahydrofuran (THF) led to the formation of a white precipitate which was filtered off and dried. The resulting powder was dissolved in water to give colorless crystals of the title compound after four weeks.

**Figure 1**

Molecular structure of the title compound (I), displacement ellipsoids are drawn at the 50% probability level. Water molecules are omitted for clarity.

**Figure 2**

$\pi\cdots\pi$ Stacking interactions between aromatic rings of (pydcH₂) and (pydcH[·]) fragments with centroid-centroid distances of:
X1A \cdots X1B, 3.5949 (11) Å (x, y, z); X1B \cdots X1C, 3.5191 (11) Å ($1 - x, -1/2 + y, 1/2 - z$); X1C \cdots X1D, 3.5789 (11) Å ($1 - x, 1/2 + y, 1/2 - z$); X1A \cdots X1D, 3.5108 (11) Å ($x, 1 + y, z$).

**Figure 3**

Unit-cell packing diagram of the compound (I). Hydrogen bonds are indicated by dashed lines.

Propane-1,3-diaminium–2-carboxypyridine-6-carboxylate– pyridine-2,6-dicarboxylic acid–water (1/2/2/8)

Crystal data



$M_r = 886.73$

Monoclinic, $P2_1/c$

$a = 13.5425$ (2) Å

$b = 13.5237$ (2) Å

$c = 21.7538$ (3) Å

$\beta = 99.914$ (1)°

$V = 3924.60$ (10) Å³

$Z = 4$

$F(000) = 1864$

$D_x = 1.501$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 9156 reflections

$\theta = 2.4\text{--}29.1$ °

$\mu = 0.13$ mm⁻¹

$T = 150$ K

Block, colourless

0.26 × 0.12 × 0.12 mm

Data collection

Bruker SMART 1000

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution: 100 pixels mm⁻¹

ω scans

Absorption correction: multi-scan

(SADABS; Bruker, 1998)

$T_{\min} = 0.967$, $T_{\max} = 0.985$

56174 measured reflections

9031 independent reflections

6702 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.054$

$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 1.5^\circ$
 $h = -17 \rightarrow 17$

$k = -17 \rightarrow 17$
 $l = -28 \rightarrow 28$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.139$
 $S = 1.02$
9031 reflections
550 parameters
0 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0592P)^2 + 3.5811P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.84 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.51 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.90840 (11)	0.77843 (12)	0.54180 (7)	0.0254 (3)
O2	0.74609 (11)	0.79981 (11)	0.51063 (7)	0.0249 (3)
H2A	0.6958	0.8087	0.4748	0.030*
O3	0.60302 (12)	0.87473 (13)	0.31596 (7)	0.0307 (4)
O4	0.68349 (11)	0.88557 (12)	0.23488 (7)	0.0262 (3)
O5	0.18211 (12)	0.78608 (13)	-0.00659 (7)	0.0320 (4)
H5AA	0.2027	0.7723	-0.0453	0.038*
O6	0.33622 (12)	0.84668 (13)	0.02551 (7)	0.0292 (4)
O7	0.45325 (11)	0.89020 (13)	0.23505 (7)	0.0298 (4)
H7A	0.5096	0.8838	0.2675	0.036*
O8	0.35911 (12)	0.86950 (14)	0.30873 (7)	0.0333 (4)
O9	0.15292 (11)	0.52710 (12)	-0.01050 (6)	0.0245 (3)
H9A	0.1799	0.5233	-0.0480	0.029*
O10	0.31288 (12)	0.55101 (13)	0.03474 (7)	0.0315 (4)
O11	0.39207 (11)	0.63847 (12)	0.24320 (6)	0.0262 (3)
H11A	0.4456	0.6339	0.2776	0.031*
O12	0.28767 (11)	0.61399 (11)	0.31033 (6)	0.0235 (3)
O13	0.46668 (11)	0.12691 (14)	0.17039 (7)	0.0322 (4)
O14	0.37185 (11)	0.11228 (12)	0.24412 (6)	0.0258 (3)
O15	0.02539 (11)	0.06135 (12)	0.19062 (7)	0.0255 (3)
O16	-0.04375 (11)	0.02480 (12)	0.09246 (6)	0.0248 (3)
H16A	-0.1019	0.0061	0.1086	0.030*
O17	0.18876 (11)	0.09699 (12)	0.28868 (7)	0.0263 (3)

H17A	0.1416	0.0806	0.2525	0.032*
H17B	0.2490	0.1058	0.2726	0.032*
O18	0.85758 (12)	0.72397 (12)	0.65729 (7)	0.0276 (3)
H18B	0.8642	0.7497	0.6176	0.033*
H18A	0.8308	0.7822	0.6725	0.033*
O19	0.55978 (12)	0.80178 (13)	0.42843 (7)	0.0337 (4)
H19B	0.6221	0.8311	0.4244	0.040*
H19A	0.5239	0.7533	0.4019	0.040*
O20	0.20445 (14)	0.74513 (13)	-0.11979 (8)	0.0385 (4)
H20B	0.2437	0.7866	-0.1413	0.046*
H20A	0.2206	0.6770	-0.1216	0.046*
O21	0.42936 (12)	0.54219 (13)	-0.06159 (7)	0.0338 (4)
H21B	0.3970	0.5418	-0.0262	0.041*
H21A	0.4678	0.6011	-0.0607	0.041*
O22	0.22400 (13)	0.53350 (13)	-0.11351 (7)	0.0348 (4)
H22A	0.2948	0.5373	-0.1021	0.042*
H22B	0.2014	0.4911	-0.1479	0.042*
O23	0.42837 (14)	0.89677 (16)	-0.07328 (8)	0.0464 (5)
H23B	0.3852	0.8938	-0.0432	0.056*
H23A	0.3866	0.8866	-0.1125	0.056*
O24	-0.20133 (11)	-0.03538 (11)	0.13049 (7)	0.0236 (3)
H24B	-0.1884	-0.1000	0.1470	0.028*
H24A	-0.2294	0.0031	0.1597	0.028*
C1	0.83900 (16)	0.79459 (15)	0.49983 (9)	0.0200 (4)
C2	0.85368 (15)	0.80829 (15)	0.43368 (9)	0.0187 (4)
C3	0.94700 (16)	0.79837 (15)	0.41643 (10)	0.0213 (4)
H3	1.0049	0.7844	0.4465	0.026*
C4	0.95318 (17)	0.80945 (16)	0.35423 (10)	0.0244 (5)
H4	1.0156	0.8019	0.3405	0.029*
C5	0.86744 (16)	0.83174 (15)	0.31202 (10)	0.0222 (4)
H5	0.8702	0.8397	0.2690	0.027*
C6	0.77715 (16)	0.84232 (15)	0.33360 (9)	0.0196 (4)
C7	0.68120 (16)	0.86929 (16)	0.29052 (9)	0.0223 (4)
C8	0.25328 (15)	0.82141 (16)	0.03516 (9)	0.0207 (4)
C9	0.22600 (15)	0.82952 (15)	0.09889 (9)	0.0179 (4)
C10	0.12850 (15)	0.81571 (15)	0.10843 (9)	0.0199 (4)
H10	0.0767	0.7994	0.0747	0.024*
C11	0.10811 (15)	0.82621 (15)	0.16848 (10)	0.0215 (4)
H11	0.0417	0.8191	0.1765	0.026*
C12	0.18632 (16)	0.84712 (15)	0.21610 (10)	0.0212 (4)
H12	0.1748	0.8543	0.2577	0.025*
C13	0.28225 (15)	0.85764 (15)	0.20248 (9)	0.0193 (4)
C14	0.36955 (16)	0.87368 (16)	0.25443 (10)	0.0228 (4)
C15	0.22408 (15)	0.54705 (16)	0.03722 (9)	0.0209 (4)
C16	0.18464 (15)	0.56360 (15)	0.09693 (9)	0.0177 (4)
C17	0.08312 (15)	0.55850 (15)	0.09973 (10)	0.0204 (4)
H17	0.0350	0.5465	0.0633	0.024*
C18	0.05401 (16)	0.57144 (16)	0.15715 (10)	0.0224 (4)

H18	-0.0148	0.5680	0.1608	0.027*
C19	0.12607 (15)	0.58934 (15)	0.20898 (9)	0.0197 (4)
H19	0.1079	0.5973	0.2490	0.024*
C20	0.22577 (15)	0.59546 (14)	0.20150 (9)	0.0172 (4)
C21	0.30780 (15)	0.61719 (15)	0.25620 (9)	0.0190 (4)
C22	0.38272 (15)	0.11248 (16)	0.18930 (9)	0.0208 (4)
C23	0.29575 (15)	0.09393 (14)	0.13742 (9)	0.0178 (4)
C24	0.30923 (15)	0.08566 (15)	0.07564 (9)	0.0192 (4)
H24	0.3738	0.0925	0.0647	0.023*
C25	0.22567 (16)	0.06715 (15)	0.03039 (9)	0.0201 (4)
H25	0.2320	0.0614	-0.0123	0.024*
C26	0.13305 (16)	0.05727 (15)	0.04837 (9)	0.0197 (4)
H26	0.0746	0.0452	0.0183	0.024*
C27	0.12751 (15)	0.06546 (14)	0.11150 (9)	0.0173 (4)
C28	0.03140 (15)	0.05113 (15)	0.13594 (9)	0.0193 (4)
C29	0.57849 (17)	0.82728 (18)	0.08483 (10)	0.0278 (5)
H29A	0.6472	0.8472	0.0803	0.033*
H29B	0.5371	0.8245	0.0427	0.033*
C30	0.58164 (16)	0.72587 (18)	0.11447 (10)	0.0266 (5)
H30A	0.6195	0.7297	0.1576	0.032*
H30B	0.6176	0.6798	0.0907	0.032*
C31	0.47782 (16)	0.68576 (17)	0.11606 (10)	0.0255 (5)
H31A	0.4435	0.7295	0.1422	0.031*
H31B	0.4385	0.6861	0.0733	0.031*
N1	0.76996 (13)	0.82962 (13)	0.39359 (8)	0.0192 (4)
N2	0.30298 (13)	0.85093 (12)	0.14479 (8)	0.0185 (3)
N3	0.25553 (12)	0.58254 (12)	0.14653 (7)	0.0172 (3)
N4	0.20715 (12)	0.08387 (12)	0.15546 (7)	0.0172 (3)
N5	0.53588 (15)	0.90263 (15)	0.12309 (8)	0.0285 (4)
H5A	0.4682	0.8858	0.1245	0.034*
H5B	0.5770	0.9026	0.1631	0.034*
H5C	0.5359	0.9684	0.1078	0.034*
N6	0.48067 (13)	0.58346 (14)	0.14140 (8)	0.0230 (4)
H6A	0.5210	0.5842	0.1817	0.028*
H6B	0.5084	0.5374	0.1161	0.028*
H6C	0.4155	0.5676	0.1490	0.028*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0257 (8)	0.0309 (9)	0.0189 (7)	0.0012 (6)	0.0019 (6)	0.0031 (6)
O2	0.0228 (8)	0.0324 (9)	0.0200 (7)	0.0025 (6)	0.0051 (6)	0.0006 (6)
O3	0.0221 (8)	0.0496 (11)	0.0201 (8)	-0.0014 (7)	0.0027 (6)	-0.0003 (7)
O4	0.0266 (8)	0.0358 (9)	0.0157 (7)	-0.0062 (7)	0.0017 (6)	0.0001 (6)
O5	0.0288 (9)	0.0478 (11)	0.0183 (8)	-0.0069 (7)	0.0008 (6)	-0.0078 (7)
O6	0.0266 (8)	0.0433 (10)	0.0188 (7)	-0.0046 (7)	0.0074 (6)	-0.0021 (7)
O7	0.0195 (8)	0.0508 (11)	0.0184 (7)	-0.0055 (7)	0.0007 (6)	-0.0019 (7)
O8	0.0287 (9)	0.0554 (11)	0.0162 (8)	0.0009 (8)	0.0048 (6)	-0.0034 (7)

O9	0.0243 (8)	0.0328 (9)	0.0156 (7)	-0.0044 (6)	0.0007 (6)	-0.0031 (6)
O10	0.0249 (8)	0.0493 (11)	0.0212 (8)	-0.0053 (7)	0.0066 (6)	-0.0076 (7)
O11	0.0165 (7)	0.0465 (10)	0.0151 (7)	-0.0009 (7)	0.0014 (6)	-0.0020 (7)
O12	0.0285 (8)	0.0276 (8)	0.0153 (7)	-0.0007 (6)	0.0066 (6)	-0.0005 (6)
O13	0.0181 (8)	0.0608 (12)	0.0173 (7)	-0.0034 (7)	0.0022 (6)	0.0041 (7)
O14	0.0211 (8)	0.0402 (9)	0.0157 (7)	-0.0010 (7)	0.0022 (6)	0.0008 (6)
O15	0.0225 (8)	0.0363 (9)	0.0187 (7)	-0.0022 (6)	0.0065 (6)	-0.0048 (6)
O16	0.0193 (7)	0.0366 (9)	0.0186 (7)	-0.0064 (6)	0.0035 (6)	-0.0035 (6)
O17	0.0251 (8)	0.0379 (9)	0.0163 (7)	-0.0009 (7)	0.0050 (6)	-0.0014 (6)
O18	0.0338 (9)	0.0276 (8)	0.0234 (8)	0.0022 (7)	0.0107 (7)	0.0040 (6)
O19	0.0290 (9)	0.0442 (10)	0.0285 (9)	0.0022 (7)	0.0061 (7)	-0.0040 (7)
O20	0.0527 (11)	0.0310 (9)	0.0367 (10)	-0.0008 (8)	0.0213 (8)	0.0039 (7)
O21	0.0350 (9)	0.0400 (10)	0.0301 (9)	-0.0047 (8)	0.0156 (7)	-0.0063 (7)
O22	0.0425 (10)	0.0408 (10)	0.0223 (8)	-0.0057 (8)	0.0087 (7)	-0.0043 (7)
O23	0.0338 (10)	0.0815 (15)	0.0254 (9)	0.0036 (10)	0.0095 (7)	0.0090 (9)
O24	0.0228 (8)	0.0285 (8)	0.0205 (7)	-0.0024 (6)	0.0060 (6)	-0.0031 (6)
C1	0.0236 (11)	0.0171 (10)	0.0190 (10)	-0.0004 (8)	0.0035 (8)	0.0001 (8)
C2	0.0215 (10)	0.0160 (10)	0.0188 (10)	-0.0015 (8)	0.0043 (8)	-0.0009 (8)
C3	0.0209 (10)	0.0189 (10)	0.0237 (10)	0.0004 (8)	0.0027 (8)	0.0010 (8)
C4	0.0252 (11)	0.0234 (11)	0.0269 (11)	0.0000 (9)	0.0104 (9)	0.0010 (9)
C5	0.0275 (11)	0.0213 (11)	0.0192 (10)	-0.0028 (8)	0.0075 (8)	-0.0014 (8)
C6	0.0258 (11)	0.0158 (10)	0.0169 (9)	-0.0039 (8)	0.0033 (8)	-0.0023 (8)
C7	0.0236 (11)	0.0239 (11)	0.0188 (10)	-0.0064 (8)	0.0019 (8)	-0.0028 (8)
C8	0.0200 (10)	0.0221 (10)	0.0195 (10)	0.0009 (8)	0.0020 (8)	0.0004 (8)
C9	0.0208 (10)	0.0155 (9)	0.0176 (9)	0.0003 (8)	0.0037 (8)	0.0009 (8)
C10	0.0208 (10)	0.0174 (10)	0.0205 (10)	-0.0015 (8)	0.0002 (8)	0.0033 (8)
C11	0.0177 (10)	0.0205 (10)	0.0275 (11)	0.0004 (8)	0.0072 (8)	0.0035 (8)
C12	0.0244 (11)	0.0208 (10)	0.0195 (10)	0.0020 (8)	0.0073 (8)	0.0010 (8)
C13	0.0218 (10)	0.0183 (10)	0.0180 (10)	-0.0006 (8)	0.0034 (8)	0.0001 (8)
C14	0.0217 (11)	0.0268 (11)	0.0202 (10)	0.0008 (9)	0.0042 (8)	-0.0024 (8)
C15	0.0222 (11)	0.0224 (10)	0.0180 (10)	-0.0022 (8)	0.0030 (8)	-0.0009 (8)
C16	0.0190 (10)	0.0168 (10)	0.0172 (9)	0.0004 (8)	0.0030 (7)	0.0012 (8)
C17	0.0197 (10)	0.0174 (10)	0.0226 (10)	-0.0009 (8)	-0.0004 (8)	0.0002 (8)
C18	0.0177 (10)	0.0219 (11)	0.0288 (11)	0.0006 (8)	0.0073 (8)	0.0011 (9)
C19	0.0206 (10)	0.0192 (10)	0.0211 (10)	0.0010 (8)	0.0083 (8)	0.0001 (8)
C20	0.0198 (10)	0.0161 (10)	0.0165 (9)	0.0021 (7)	0.0051 (8)	0.0015 (7)
C21	0.0222 (10)	0.0188 (10)	0.0166 (9)	0.0027 (8)	0.0053 (8)	-0.0003 (8)
C22	0.0184 (10)	0.0261 (11)	0.0176 (10)	0.0006 (8)	0.0023 (8)	0.0032 (8)
C23	0.0198 (10)	0.0162 (10)	0.0171 (9)	0.0002 (8)	0.0023 (8)	0.0009 (8)
C24	0.0210 (10)	0.0188 (10)	0.0188 (10)	0.0000 (8)	0.0059 (8)	0.0002 (8)
C25	0.0272 (11)	0.0190 (10)	0.0146 (9)	-0.0025 (8)	0.0049 (8)	-0.0005 (8)
C26	0.0222 (10)	0.0186 (10)	0.0175 (9)	-0.0028 (8)	0.0014 (8)	-0.0008 (8)
C27	0.0192 (10)	0.0149 (9)	0.0179 (9)	0.0002 (7)	0.0036 (8)	0.0010 (7)
C28	0.0219 (10)	0.0183 (10)	0.0177 (10)	0.0014 (8)	0.0036 (8)	-0.0002 (8)
C29	0.0225 (11)	0.0417 (14)	0.0202 (10)	-0.0082 (10)	0.0064 (9)	-0.0055 (10)
C30	0.0169 (10)	0.0398 (13)	0.0235 (11)	-0.0014 (9)	0.0046 (8)	-0.0042 (10)
C31	0.0187 (10)	0.0327 (12)	0.0249 (11)	0.0005 (9)	0.0033 (8)	0.0008 (9)
N1	0.0226 (9)	0.0185 (9)	0.0168 (8)	-0.0022 (7)	0.0036 (7)	-0.0006 (7)

N2	0.0205 (9)	0.0185 (8)	0.0171 (8)	0.0006 (7)	0.0043 (7)	0.0010 (7)
N3	0.0188 (8)	0.0176 (8)	0.0160 (8)	-0.0001 (7)	0.0053 (6)	0.0001 (7)
N4	0.0183 (8)	0.0179 (8)	0.0153 (8)	0.0010 (6)	0.0028 (6)	0.0002 (6)
N5	0.0297 (10)	0.0355 (11)	0.0199 (9)	-0.0053 (8)	0.0034 (8)	-0.0007 (8)
N6	0.0186 (9)	0.0329 (10)	0.0181 (8)	0.0010 (7)	0.0052 (7)	-0.0035 (7)

Geometric parameters (\AA , $^{\circ}$)

O1—C1	1.212 (3)	C8—C9	1.499 (3)
O2—C1	1.321 (2)	C9—N2	1.345 (3)
O2—H2A	0.9499	C9—C10	1.384 (3)
O3—C7	1.278 (3)	C10—C11	1.388 (3)
O4—C7	1.236 (2)	C10—H10	0.9500
O5—C8	1.296 (3)	C11—C12	1.377 (3)
O5—H5AA	0.9501	C11—H11	0.9500
O6—C8	1.226 (3)	C12—C13	1.389 (3)
O7—C14	1.295 (3)	C12—H12	0.9500
O7—H7A	0.9501	C13—N2	1.335 (3)
O8—C14	1.215 (3)	C13—C14	1.504 (3)
O9—C15	1.317 (2)	C15—C16	1.504 (3)
O9—H9A	0.9499	C16—N3	1.340 (3)
O10—C15	1.214 (3)	C16—C17	1.388 (3)
O11—C21	1.255 (2)	C17—C18	1.384 (3)
O11—H11A	0.9499	C17—H17	0.9500
O12—C21	1.254 (2)	C18—C19	1.380 (3)
O13—C22	1.289 (2)	C18—H18	0.9500
O14—C22	1.227 (2)	C19—C20	1.390 (3)
O15—C28	1.214 (2)	C19—H19	0.9500
O16—C28	1.314 (2)	C20—N3	1.338 (2)
O16—H16A	0.9500	C20—C21	1.510 (3)
O17—H17A	0.9500	C22—C23	1.505 (3)
O17—H17B	0.9499	C23—N4	1.332 (3)
O18—H18B	0.9501	C23—C24	1.392 (3)
O18—H18A	0.9499	C24—C25	1.389 (3)
O19—H19B	0.9501	C24—H24	0.9500
O19—H19A	0.9500	C25—C26	1.383 (3)
O20—H20B	0.9502	C25—H25	0.9500
O20—H20A	0.9501	C26—C27	1.393 (3)
O21—H21B	0.9500	C26—H26	0.9500
O21—H21A	0.9500	C27—N4	1.336 (3)
O22—H22A	0.9500	C27—C28	1.501 (3)
O22—H22B	0.9500	C29—N5	1.493 (3)
O23—H23B	0.9500	C29—C30	1.513 (3)
O23—H23A	0.9499	C29—H29A	0.9900
O24—H24B	0.9500	C29—H29B	0.9900
O24—H24A	0.9501	C30—C31	1.513 (3)
C1—C2	1.498 (3)	C30—H30A	0.9900
C2—N1	1.337 (3)	C30—H30B	0.9900

C2—C3	1.385 (3)	C31—N6	1.487 (3)
C3—C4	1.378 (3)	C31—H31A	0.9900
C3—H3	0.9500	C31—H31B	0.9900
C4—C5	1.384 (3)	N5—H5A	0.9500
C4—H4	0.9500	N5—H5B	0.9500
C5—C6	1.391 (3)	N5—H5C	0.9500
C5—H5	0.9500	N6—H6A	0.9502
C6—N1	1.336 (3)	N6—H6B	0.9499
C6—C7	1.510 (3)	N6—H6C	0.9500
C1—O2—H2A	115.5	C19—C18—H18	120.4
C8—O5—H5AA	113.4	C17—C18—H18	120.4
C14—O7—H7A	112.3	C18—C19—C20	118.65 (18)
C15—O9—H9A	110.5	C18—C19—H19	120.7
C21—O11—H11A	114.1	C20—C19—H19	120.7
C28—O16—H16A	113.3	N3—C20—C19	123.04 (18)
H17A—O17—H17B	102.8	N3—C20—C21	115.83 (17)
H18B—O18—H18A	96.6	C19—C20—C21	121.13 (17)
H19B—O19—H19A	127.0	O12—C21—O11	124.91 (19)
H20B—O20—H20A	113.5	O12—C21—C20	118.92 (18)
H21B—O21—H21A	108.4	O11—C21—C20	116.17 (17)
H22A—O22—H22B	114.4	O14—C22—O13	124.81 (19)
H23B—O23—H23A	105.7	O14—C22—C23	121.27 (18)
H24B—O24—H24A	108.7	O13—C22—C23	113.92 (17)
O1—C1—O2	120.96 (18)	N4—C23—C24	123.43 (18)
O1—C1—C2	122.15 (19)	N4—C23—C22	115.11 (17)
O2—C1—C2	116.88 (17)	C24—C23—C22	121.44 (18)
N1—C2—C3	123.70 (18)	C25—C24—C23	118.21 (18)
N1—C2—C1	114.54 (17)	C25—C24—H24	120.9
C3—C2—C1	121.76 (18)	C23—C24—H24	120.9
C4—C3—C2	117.9 (2)	C26—C25—C24	119.01 (18)
C4—C3—H3	121.1	C26—C25—H25	120.5
C2—C3—H3	121.1	C24—C25—H25	120.5
C3—C4—C5	119.3 (2)	C25—C26—C27	118.46 (19)
C3—C4—H4	120.4	C25—C26—H26	120.8
C5—C4—H4	120.4	C27—C26—H26	120.8
C4—C5—C6	118.98 (19)	N4—C27—C26	123.16 (18)
C4—C5—H5	120.5	N4—C27—C28	114.39 (17)
C6—C5—H5	120.5	C26—C27—C28	122.43 (18)
N1—C6—C5	122.19 (19)	O15—C28—O16	124.19 (19)
N1—C6—C7	115.94 (18)	O15—C28—C27	122.74 (18)
C5—C6—C7	121.87 (18)	O16—C28—C27	113.06 (17)
O4—C7—O3	125.2 (2)	N5—C29—C30	111.50 (17)
O4—C7—C6	119.12 (19)	N5—C29—H29A	109.3
O3—C7—C6	115.69 (18)	C30—C29—H29A	109.3
O6—C8—O5	125.11 (19)	N5—C29—H29B	109.3
O6—C8—C9	121.12 (18)	C30—C29—H29B	109.3
O5—C8—C9	113.76 (18)	H29A—C29—H29B	108.0

N2—C9—C10	123.73 (18)	C29—C30—C31	112.13 (19)
N2—C9—C8	114.79 (17)	C29—C30—H30A	109.2
C10—C9—C8	121.48 (18)	C31—C30—H30A	109.2
C9—C10—C11	118.47 (19)	C29—C30—H30B	109.2
C9—C10—H10	120.8	C31—C30—H30B	109.2
C11—C10—H10	120.8	H30A—C30—H30B	107.9
C12—C11—C10	118.49 (19)	N6—C31—C30	112.15 (18)
C12—C11—H11	120.8	N6—C31—H31A	109.2
C10—C11—H11	120.8	C30—C31—H31A	109.2
C11—C12—C13	119.14 (19)	N6—C31—H31B	109.2
C11—C12—H12	120.4	C30—C31—H31B	109.2
C13—C12—H12	120.4	H31A—C31—H31B	107.9
N2—C13—C12	123.32 (19)	C6—N1—C2	117.92 (18)
N2—C13—C14	116.79 (18)	C13—N2—C9	116.79 (17)
C12—C13—C14	119.84 (18)	C20—N3—C16	117.42 (17)
O8—C14—O7	125.4 (2)	C23—N4—C27	117.72 (16)
O8—C14—C13	121.05 (19)	C29—N5—H5A	108.8
O7—C14—C13	113.53 (17)	C29—N5—H5B	106.8
O10—C15—O9	124.74 (19)	H5A—N5—H5B	112.7
O10—C15—C16	122.26 (18)	C29—N5—H5C	114.9
O9—C15—C16	113.00 (17)	H5A—N5—H5C	107.1
N3—C16—C17	123.50 (18)	H5B—N5—H5C	106.6
N3—C16—C15	114.34 (17)	C31—N6—H6A	107.8
C17—C16—C15	122.16 (18)	C31—N6—H6B	112.5
C18—C17—C16	118.13 (19)	H6A—N6—H6B	108.7
C18—C17—H17	120.9	C31—N6—H6C	108.0
C16—C17—H17	120.9	H6A—N6—H6C	104.2
C19—C18—C17	119.23 (19)	H6B—N6—H6C	115.1
O1—C1—C2—N1	-177.34 (19)	C18—C19—C20—C21	178.60 (19)
O2—C1—C2—N1	3.4 (3)	N3—C20—C21—O12	-168.41 (18)
O1—C1—C2—C3	3.2 (3)	C19—C20—C21—O12	11.4 (3)
O2—C1—C2—C3	-176.03 (18)	N3—C20—C21—O11	12.3 (3)
N1—C2—C3—C4	-1.2 (3)	C19—C20—C21—O11	-167.82 (19)
C1—C2—C3—C4	178.17 (19)	O14—C22—C23—N4	4.0 (3)
C2—C3—C4—C5	1.3 (3)	O13—C22—C23—N4	-176.68 (19)
C3—C4—C5—C6	0.0 (3)	O14—C22—C23—C24	-174.4 (2)
C4—C5—C6—N1	-1.5 (3)	O13—C22—C23—C24	4.9 (3)
C4—C5—C6—C7	178.33 (19)	N4—C23—C24—C25	0.8 (3)
N1—C6—C7—O4	177.41 (19)	C22—C23—C24—C25	179.11 (19)
C5—C6—C7—O4	-2.4 (3)	C23—C24—C25—C26	-0.3 (3)
N1—C6—C7—O3	-1.6 (3)	C24—C25—C26—C27	-0.6 (3)
C5—C6—C7—O3	178.6 (2)	C25—C26—C27—N4	1.0 (3)
O6—C8—C9—N2	-10.0 (3)	C25—C26—C27—C28	-177.05 (18)
O5—C8—C9—N2	169.95 (18)	N4—C27—C28—O15	4.3 (3)
O6—C8—C9—C10	170.0 (2)	C26—C27—C28—O15	-177.5 (2)
O5—C8—C9—C10	-10.1 (3)	N4—C27—C28—O16	-174.48 (17)
N2—C9—C10—C11	1.2 (3)	C26—C27—C28—O16	3.7 (3)

C8—C9—C10—C11	−178.76 (19)	N5—C29—C30—C31	65.6 (2)
C9—C10—C11—C12	−1.8 (3)	C29—C30—C31—N6	176.28 (17)
C10—C11—C12—C13	0.4 (3)	C5—C6—N1—C2	1.6 (3)
C11—C12—C13—N2	1.7 (3)	C7—C6—N1—C2	−178.23 (18)
C11—C12—C13—C14	−175.90 (19)	C3—C2—N1—C6	−0.2 (3)
N2—C13—C14—O8	−171.6 (2)	C1—C2—N1—C6	−179.62 (17)
C12—C13—C14—O8	6.2 (3)	C12—C13—N2—C9	−2.3 (3)
N2—C13—C14—O7	7.5 (3)	C14—C13—N2—C9	175.36 (18)
C12—C13—C14—O7	−174.77 (19)	C10—C9—N2—C13	0.8 (3)
O10—C15—C16—N3	−0.2 (3)	C8—C9—N2—C13	−179.19 (18)
O9—C15—C16—N3	179.01 (17)	C19—C20—N3—C16	0.5 (3)
O10—C15—C16—C17	−179.6 (2)	C21—C20—N3—C16	−179.69 (17)
O9—C15—C16—C17	−0.4 (3)	C17—C16—N3—C20	1.1 (3)
N3—C16—C17—C18	−1.5 (3)	C15—C16—N3—C20	−178.31 (17)
C15—C16—C17—C18	177.83 (19)	C24—C23—N4—C27	−0.4 (3)
C16—C17—C18—C19	0.4 (3)	C22—C23—N4—C27	−178.80 (17)
C17—C18—C19—C20	1.1 (3)	C26—C27—N4—C23	−0.5 (3)
C18—C19—C20—N3	−1.6 (3)	C28—C27—N4—C23	177.68 (17)

Hydrogen-bond geometry (\AA , °)

$D\cdots H$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
O2—H24 \cdots O19	0.95	1.94	2.830 (2)	154
O5—H5A \cdots O20	0.95	1.67	2.592 (2)	164
O7—H7A \cdots O3	0.95	1.51	2.455 (2)	177
O7—H7A \cdots O4	0.95	2.57	3.119 (2)	117
O9—H9A \cdots O22	0.95	1.64	2.588 (2)	172
O11—H11A \cdots O13 ⁱ	0.95	1.50	2.445 (2)	177
O11—H11A \cdots O14 ⁱ	0.95	2.61	3.181 (2)	119
O16—H16A \cdots O24	0.95	1.60	2.551 (2)	174
O17—H17A \cdots O15	0.95	1.90	2.838 (2)	167
O17—H17A \cdots N4	0.95	2.43	2.956 (2)	115
O17—H17B \cdots O14	0.95	1.87	2.821 (2)	175
O18—H18B \cdots O1	0.95	1.89	2.815 (2)	164
O18—H18A \cdots O17 ⁱⁱ	0.95	1.88	2.809 (2)	166
O19—H19B \cdots N1	0.95	2.22	3.092 (2)	152
O19—H19A \cdots O13 ⁱ	0.95	2.34	3.174 (2)	146
O20—H20B \cdots O12 ⁱⁱⁱ	0.95	1.87	2.793 (2)	164
O20—H20A \cdots O22	0.95	1.95	2.875 (3)	165
O21—H21B \cdots O10	0.95	1.89	2.834 (2)	170
O21—H21A \cdots O19 ⁱⁱⁱ	0.95	1.85	2.784 (2)	166
O22—H22A \cdots O21	0.95	1.88	2.818 (2)	168
O22—H22B \cdots O17 ^{iv}	0.95	1.81	2.742 (2)	167
O23—H23B \cdots O6	0.95	1.85	2.749 (2)	157
O23—H23A \cdots O12 ⁱⁱⁱ	0.95	1.96	2.899 (2)	170
O24—H24B \cdots O18 ^v	0.95	1.79	2.706 (2)	161
O24—H24A \cdots O12 ^{vi}	0.95	1.86	2.761 (2)	157
N5—H5A \cdots N2	0.95	2.40	3.342 (3)	171

N5—H5A···O6	0.95	2.61	3.227 (2)	123
N5—H5B···O4	0.95	1.95	2.879 (2)	166
N5—H5C···O23 ^{vii}	0.95	2.06	2.991 (3)	166
N6—H6A···O14 ⁱ	0.95	2.01	2.938 (2)	164
N6—H6B···O21 ^{viii}	0.95	1.90	2.849 (2)	174
N6—H6C···N3	0.95	2.17	3.070 (2)	158
C5—H5···O18 ⁱⁱⁱ	0.95	2.56	3.429 (3)	153
C10—H10···O1 ^{ix}	0.95	2.50	3.335 (3)	147
C11—H11···O18 ^{ix}	0.95	2.52	3.427 (3)	159
C19—H19···O15 ^x	0.95	2.46	3.266 (3)	143
C25—H25···O24 ^{xi}	0.95	2.56	3.485 (3)	165
C26—H26···O16 ^{xi}	0.95	2.56	3.287 (3)	134
C31—H31A···N2	0.99	2.52	3.391 (3)	146
C31—H31B···O10	0.99	2.54	3.175 (3)	122

Symmetry codes: (i) $-x+1, y+1/2, -z+1/2$; (ii) $-x+1, -y+1, -z+1$; (iii) $x, -y+3/2, z-1/2$; (iv) $x, -y+1/2, z-1/2$; (v) $x-1, -y+1/2, z-1/2$; (vi) $-x, y-1/2, -z+1/2$; (vii) $-x+1, -y+2, -z$; (viii) $-x+1, -y+1, -z$; (ix) $x-1, -y+3/2, z-1/2$; (x) $-x, y+1/2, -z+1/2$; (xi) $-x, -y, -z$.