

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1,2-Bis[amino(pyrimidin-2-yl)methylene]-hydrazine dihydrate

Shuo Shi,^{a*} Tian-ming Yao,^a Xiao-ting Geng,^a Lin Chen^b and Liang-nian Ji^b^aDepartment of Chemistry, Tongji University, Shanghai, People's Republic of China,and ^bSchool of Chemistry and Chemical Engineering, Sun Yat-Sen University, Guangzhou, People's Republic of China

Correspondence e-mail: shishuo@mail.tongji.edu.cn


Received 27 November 2007; accepted 6 December 2007

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.036; wR factor = 0.102; data-to-parameter ratio = 9.7.

The centrosymmetric organic molecule in the title compound, $\text{C}_{10}\text{H}_{10}\text{N}_8 \cdot 2\text{H}_2\text{O}$, is essentially flat and has a *trans* configuration. The molecules are linked by intermolecular $\text{O}-\text{H} \cdots \text{N}$, $\text{N}-\text{H} \cdots \text{O}$ and $\text{N}-\text{H} \cdots \text{N}$ hydrogen bonds to form a linear chain structure.

Related literature

For related structures, see: Armstrong *et al.* (1998); Case (1965); Thompson *et al.* (1998); Xu *et al.* (1997, 1998, 2000, 2001).



Experimental

Crystal data

 $\text{C}_{10}\text{H}_{10}\text{N}_8 \cdot 2\text{H}_2\text{O}$ $M_r = 278.15$ Triclinic, $P\bar{1}$ $a = 6.109$ (2) Å $b = 7.502$ (3) Å $c = 7.588$ (3) Å $\alpha = 105.112$ (6)° $\beta = 106.975$ (7)° $\gamma = 99.193$ (6)° $V = 310.41$ (19) Å³ $Z = 1$ Mo $K\alpha$ radiation $\mu = 0.11$ mm⁻¹ $T = 293$ (2) K

0.48 × 0.22 × 0.18 mm

Data collection

Bruker SMART APEX CCD
area-detector diffractometer
Absorption correction: multi-scan
(*SADABS*; Bruker, 2000)
 $T_{\min} = 0.949$, $T_{\max} = 0.980$

1526 measured reflections
1036 independent reflections
778 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.008$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.036$ $wR(F^2) = 0.102$ $S = 1.04$

1036 reflections

107 parameters

H atoms treated by a mixture of
independent and constrained
refinement

 $\Delta\rho_{\max} = 0.15$ e Å⁻³ $\Delta\rho_{\min} = -0.14$ e Å⁻³

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N3}-\text{H3B} \cdots \text{N1}^{\text{ii}}$	0.85 (2)	2.59 (2)	3.276 (2)	138.5 (16)
$\text{N3}-\text{H3C} \cdots \text{O1W}^{\text{iii}}$	0.89 (2)	2.17 (3)	3.043 (3)	166.7 (19)
$\text{O1W}-\text{H1WA} \cdots \text{N2}^{\text{iv}}$	0.79 (3)	2.20 (3)	2.979 (2)	168 (3)
$\text{O1W}-\text{H1WB} \cdots \text{N4}$	0.91 (3)	2.16 (3)	3.055 (2)	172 (2)

Symmetry codes: (ii) $-x + 2, -y + 2, -z + 2$; (iii) $x + 1, y, z$; (iv) $-x, -y + 1, -z + 1$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SMART*; data reduction: *SAINT-Plus* and *SHELXTL* (Bruker, 1998); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

The authors thank the National Natural Science Foundation of China, the Research Fund for the Doctoral Program of Higher Education, and the Program for Young Excellent Talents in Tongji University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2403).

References

- Armstrong, J. A., Barnes, J. C. & Weakley, T. J. R. (1998). *Acta Cryst.* **C54**, 1923–1925.
- Bruker (1998). *SMART*, *SAINT-Plus* and *SHELXTL*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2000). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Case, F. H. (1965). *J. Org. Chem.* **30**, 931–933.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Thompson, L. K., Xu, Z. Q., Goeta, A. E., Howard, J. A. K., Clase, H. J. & Miller, D. O. (1998). *Inorg. Chem.* **37**, 3217–3229.
- Xu, Z. Q., Thompson, L. K., Black, D. A., Ralph, C., Miller, D. O., Leech, M. A. & Howard, J. A. K. (2001). *J. Chem. Soc. Dalton Trans.* pp. 2042–2048.
- Xu, Z. Q., Thompson, L. K. & Miller, D. O. (1997). *Inorg. Chem.* **36**, 3985–3995.
- Xu, Z. Q., Thompson, L. K., Miller, D. O., Clase, H. J., Howard, J. A. K. & Goeta, A. E. (1998). *Inorg. Chem.* **37**, 3620–3627.
- Xu, Z. Q., White, S., Thompson, L. K., Miller, D. O., Ohba, M., Okawa, H., Wilson, C. & Howard, J. A. K. (2000). *J. Chem. Soc. Dalton Trans.* pp. 1751–1757.

supporting information

Acta Cryst. (2008). E64, o272 [https://doi.org/10.1107/S1600536807065889]

1,2-Bis[amino(pyrimidin-2-yl)methylene]hydrazine dihydrate

Shuo Shi, Tian-ming Yao, Xiao-ting Geng, Lin Chen and Liang-nian Ji

S1. Comment

The title compound, (I) (Fig. 1), can be regarded as a dihydrazidine. It is formed as the major product from mixing 2-cyanopyrimidine and hydrazine in ethanol (Case, 1965) and the minor product is Pyrimidine-2-carboxamide hydrazone, (II)(Scheme. 1). Compound (I) has now been shown to have *trans* geometry (Fig. 1), with all atoms essentially coplanar. The overall *trans* configuration is therefore due mainly to steric repulsion effects. The title compound contains a single N—N bond, presents several possible mononucleating and dinucleating coordination modes and, also, the potential for free rotation about the N—N bond. The flexible geometries result from the ability of the systems to rotate freely about the single N—N bond of the diazine fragment of the compound.

S2. Refinement

All H atoms were placed in geometrically positions and constrained to ride on their parent atoms, with N—H distances in the range 0.85—0.89 Å and C—H = 0.93 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C or N})$ for all H atoms.




Figure 1

The molecular structure of (I), with atom labels.

1,2-Bis[amino(pyrimidin-2-yl)methylene]hydrazine dihydrate

Crystal data

$\text{C}_{10}\text{H}_{10}\text{N}_8 \cdot 2\text{H}_2\text{O}$

$M_r = 278.15$

Triclinic, $P\bar{1}$

$a = 6.109$ (2) Å

$b = 7.502$ (3) Å

$c = 7.588$ (3) Å

$\alpha = 105.112$ (6)°

$\beta = 106.975$ (7)°

$\gamma = 99.193$ (6)°

$V = 310.41$ (19) Å³

$Z = 1$

$F(000) = 146$

$D_x = 1.489$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

$\mu = 0.11 \text{ mm}^{-1}$
 $T = 293 \text{ K}$

Prism, yellow
 $0.48 \times 0.22 \times 0.18 \text{ mm}$

Data collection

Bruker SMART APEX CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan
 (SADABS; Bruker, 2000)
 $T_{\min} = 0.949, T_{\max} = 0.980$

1526 measured reflections
 1036 independent reflections
 778 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.008$
 $\theta_{\max} = 25.0^\circ, \theta_{\min} = 2.9^\circ$
 $h = -7 \rightarrow 7$
 $k = -8 \rightarrow 8$
 $l = -9 \rightarrow 8$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.102$
 $S = 1.04$
 1036 reflections
 107 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0576P)^2 + 0.0469P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.15 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.14 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.6787 (3)	0.9111 (3)	1.2198 (3)	0.0443 (5)
H1A	0.7942	1.0004	1.3310	0.053*
C2	0.4674 (4)	0.8331 (3)	1.2312 (3)	0.0452 (5)
H2A	0.4367	0.8673	1.3468	0.054*
C3	0.3041 (4)	0.7028 (3)	1.0649 (3)	0.0440 (5)
H3A	0.1575	0.6498	1.0681	0.053*
C4	0.5561 (3)	0.7320 (2)	0.9025 (2)	0.0309 (4)
C5	0.6110 (3)	0.6704 (2)	0.7210 (2)	0.0307 (4)
N1	0.7253 (3)	0.8640 (2)	1.0550 (2)	0.0382 (4)
N2	0.3437 (3)	0.6477 (2)	0.8988 (2)	0.0385 (4)
N3	0.8159 (3)	0.7655 (3)	0.7217 (3)	0.0469 (5)
H3B	0.900 (3)	0.857 (3)	0.825 (3)	0.042 (6)*
H3C	0.856 (4)	0.732 (3)	0.617 (3)	0.052 (6)*

N4	0.4600 (2)	0.5260 (2)	0.5788 (2)	0.0334 (4)
O1W	0.0384 (3)	0.6771 (2)	0.4092 (2)	0.0477 (4)
H1WA	-0.074 (5)	0.593 (4)	0.338 (4)	0.079 (10)*
H1WB	0.153 (5)	0.624 (4)	0.462 (4)	0.088 (10)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0481 (12)	0.0404 (11)	0.0308 (11)	-0.0017 (9)	0.0107 (9)	0.0008 (9)
C2	0.0579 (13)	0.0404 (11)	0.0357 (11)	0.0055 (10)	0.0225 (10)	0.0069 (9)
C3	0.0444 (11)	0.0439 (11)	0.0446 (12)	0.0037 (9)	0.0236 (9)	0.0112 (10)
C4	0.0288 (10)	0.0301 (9)	0.0312 (10)	0.0067 (7)	0.0086 (8)	0.0085 (8)
C5	0.0248 (9)	0.0309 (9)	0.0318 (10)	0.0033 (7)	0.0087 (8)	0.0066 (8)
N1	0.0375 (9)	0.0355 (9)	0.0317 (9)	0.0005 (7)	0.0088 (7)	0.0036 (7)
N2	0.0336 (8)	0.0403 (9)	0.0355 (9)	0.0017 (7)	0.0132 (7)	0.0055 (7)
N3	0.0373 (10)	0.0480 (11)	0.0388 (11)	-0.0092 (8)	0.0184 (8)	-0.0062 (9)
N4	0.0295 (8)	0.0374 (9)	0.0285 (8)	0.0037 (7)	0.0114 (7)	0.0043 (7)
O1W	0.0387 (9)	0.0457 (9)	0.0505 (9)	0.0026 (8)	0.0134 (7)	0.0097 (8)

Geometric parameters (Å, °)

C1—N1	1.335 (2)	C4—C5	1.487 (2)
C1—C2	1.366 (3)	C5—N4	1.296 (2)
C1—H1A	0.9300	C5—N3	1.336 (2)
C2—C3	1.361 (3)	N3—H3B	0.85 (2)
C2—H2A	0.9300	N3—H3C	0.89 (2)
C3—N2	1.325 (3)	N4—N4 ⁱ	1.407 (3)
C3—H3A	0.9300	O1W—H1WA	0.79 (3)
C4—N1	1.328 (2)	O1W—H1WB	0.91 (3)
C4—N2	1.339 (2)		
N1—C1—C2	122.40 (17)	N2—C4—C5	117.39 (15)
N1—C1—H1A	118.8	N4—C5—N3	125.86 (17)
C2—C1—H1A	118.8	N4—C5—C4	117.26 (15)
C3—C2—C1	116.68 (18)	N3—C5—C4	116.84 (16)
C3—C2—H2A	121.7	C4—N1—C1	116.03 (16)
C1—C2—H2A	121.7	C3—N2—C4	115.50 (16)
N2—C3—C2	123.30 (19)	C5—N3—H3B	116.4 (13)
N2—C3—H3A	118.4	C5—N3—H3C	119.7 (14)
C2—C3—H3A	118.4	H3B—N3—H3C	123.9 (19)
N1—C4—N2	126.04 (17)	C5—N4—N4 ⁱ	111.67 (16)
N1—C4—C5	116.56 (15)	H1WA—O1W—H1WB	108 (3)

Symmetry code: (i) $-x+1, -y+1, -z+1$.*Hydrogen-bond geometry (Å, °)*

$D—H\cdots A$	$D—H$	$H\cdots A$	$D\cdots A$	$D—H\cdots A$
N3—H3B \cdots N1 ⁱⁱ	0.85 (2)	2.59 (2)	3.276 (2)	138.5 (16)

N3—H3C···O1W ⁱⁱ	0.89 (2)	2.17 (3)	3.043 (3)	166.7 (19)
O1W—H1WA···N2 ^{iv}	0.79 (3)	2.20 (3)	2.979 (2)	168 (3)
O1W—H1WB···N4	0.91 (3)	2.16 (3)	3.055 (2)	172 (2)

Symmetry codes: (ii) $-x+2, -y+2, -z+2$; (iii) $x+1, y, z$; (iv) $-x, -y+1, -z+1$.