

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## A monoclinic polymorph of 1,3-bis(2-pyridylaminomethyl)benzene

Li-Na Zhu,<sup>a</sup> Shan Gao<sup>a</sup> and Seik Weng Ng<sup>b\*</sup>

<sup>a</sup>School of Chemistry and Materials Science, Heilongjiang University, Harbin 150080, People's Republic of China, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: seikweng@um.edu.my


Received 16 November 2007; accepted 27 November 2007

Key indicators: single-crystal X-ray study;  $T = 295$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.047;  $wR$  factor = 0.150; data-to-parameter ratio = 16.9.

The molecules of the title compound,  $\text{C}_{18}\text{H}_{18}\text{N}_4$ , are linked by two different  $\text{N}-\text{H}\cdots\text{N}_{\text{pyridyl}}$  hydrogen bonds into a linear chain.

## Related literature

In the orthorhombic polymorph, the molecule lies on a twofold rotation axis; adjacent molecules are hydrogen-bonded into a linear chain, see: Zhu *et al.* (2007).



## Experimental

## Crystal data

$\text{C}_{18}\text{H}_{18}\text{N}_4$   
 $M_r = 290.36$   
 Monoclinic,  $C2/c$   
 $a = 35.647$  (2) Å  
 $b = 5.7899$  (4) Å  
 $c = 14.9019$  (9) Å  
 $\beta = 94.074$  (2)°

$V = 3067.9$  (3) Å<sup>3</sup>  
 $Z = 8$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.08$  mm<sup>-1</sup>  
 $T = 295$  (2) K  
 $0.32 \times 0.28 \times 0.24$  mm

## Data collection

Rigaku R-Axis RAPID  
 diffractometer  
 Absorption correction: multi-scan  
 (ABSCOR; Higashi, 1995)  
 $T_{\min} = 0.589$ ,  $T_{\max} = 0.982$

13991 measured reflections  
 3512 independent reflections  
 1833 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.047$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.150$   
 $S = 1.02$   
 3512 reflections  
 208 parameters  
 2 restraints

H atoms treated by a mixture of  
 independent and constrained  
 refinement  
 $\Delta\rho_{\max} = 0.18$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.17$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N2}-\text{H2N}\cdots\text{N1}^i$	0.86 (1)	2.19 (1)	3.047 (2)	175 (2)
$\text{N3}-\text{H3N}\cdots\text{N4}^{ii}$	0.86 (1)	2.24 (1)	3.104 (2)	177 (2)

Symmetry codes: (i)  $-x + \frac{1}{2}, -y + \frac{1}{2}, -z + 1$ ; (ii)  $-x + 1, -y + 1, -z + 1$ .

Data collection: *RAPID-AUTO* (Rigaku Corporation, 1998); cell refinement: *RAPID-AUTO*; data reduction: *CrystalStructure* (Rigaku/MSK, 2002); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2007).

We thank the Heilongjiang Province Natural Science Foundation (No. B200501), the Scientific Fund for Remarkable Teachers of Heilongjiang Province (No. 1054 G036), Heilongjiang University and the University of Malaya for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SG2210).

## References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
 Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.  
 Rigaku Corporation (1998). *RAPID-AUTO*. Rigaku Corporation, Tokyo, Japan.  
 Rigaku/MSK (2002). *CrystalStructure*. Rigaku/MSK, The Woodlands, Texas, USA.  
 Sheldrick, G. M. (1997). *SHELXL97* and *SHELXS97*. University of Göttingen, Germany.  
 Westrip, S. P. (2007). *publCIF*. In preparation.  
 Zhu, L.-N., Gao, S. & Huo, L.-H. (2007). *Acta Cryst.* **E63**, o4399.

## supporting information

*Acta Cryst.* (2008). E64, o252 [https://doi.org/10.1107/S1600536807063878]

## A monoclinic polymorph of 1,3-bis(2-pyridylaminomethyl)benzene

Li-Na Zhu, Shan Gao and Seik Weng Ng

## S1. Experimental

The compound was synthesized as described (Zhu *et al.*, 2007), and crystals were obtained upon recrystallization from methanol.

## S2. Refinement

Carbon-bound H atoms were placed in calculated positions [ $C-H$  0.93–0.97 Å and  $U_{iso}(H)$  1.25 $U_{eq}(C)$ ], were included in the refinement in the riding-model approximation. The nitrogen-bound H atoms were located and difference Fourier map and were refined with a distance restraint of  $N-H$  0.86±0.01 Å.




Figure 1

Thermal ellipsoid plot of a portion of the hydrogen-bonded structure; displacement ellipsoids are drawn at the 50% probability level, and H atoms as spheres of arbitrary radius.

## 1,3-bis(2-pyridylaminomethyl)benzene

*Crystal data*

$C_{18}H_{18}N_4$

$M_r = 290.36$

Monoclinic,  $C2/c$

Hall symbol:  $-C 2yc$

$a = 35.647$  (2) Å

$b = 5.7899$  (4) Å

$c = 14.9019$  (9) Å

$\beta = 94.074$  (2)°

$V = 3067.9$  (3) Å<sup>3</sup>

$Z = 8$

$F(000) = 1232$

$D_x = 1.257$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$   
 Cell parameters from 7473 reflections  
 $\theta = 3.0\text{--}27.5^\circ$   
 $\mu = 0.08 \text{ mm}^{-1}$

$T = 295 \text{ K}$   
 Prism, colorless  
 $0.32 \times 0.28 \times 0.24 \text{ mm}$

*Data collection*

Rigaku R-AXIS RAPID  
 diffractometer  
 Radiation source: fine-focus sealed tube  
 Graphite monochromator  
 Detector resolution: 10.000 pixels  $\text{mm}^{-1}$   
 $\omega$ -scans  
 Absorption correction: multi-scan  
 (ABSCOR; Higashi, 1995)  
 $T_{\min} = 0.589$ ,  $T_{\max} = 0.982$

13991 measured reflections  
 3512 independent reflections  
 1833 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.047$   
 $\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 3.0^\circ$   
 $h = -46 \rightarrow 46$   
 $k = -7 \rightarrow 7$   
 $l = -19 \rightarrow 19$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.048$   
 $wR(F^2) = 0.150$   
 $S = 1.02$   
 3512 reflections  
 208 parameters  
 2 restraints  
 Primary atom site location: structure-invariant  
 direct methods  
 Secondary atom site location: difference Fourier  
 map

Hydrogen site location: inferred from  
 neighbouring sites  
 H atoms treated by a mixture of independent  
 and constrained refinement  
 $w = 1/[\sigma^2(F_o^2) + (0.0752P)^2]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.18 \text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.17 \text{ e \AA}^{-3}$   
 Extinction correction: SHELXL,  
 $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$   
 Extinction coefficient: 0.0013 (5)

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.26513 (4)	0.1786 (3)	0.38622 (11)	0.0551 (4)
N2	0.28167 (4)	0.5057 (3)	0.46611 (11)	0.0571 (4)
H2N	0.2672 (5)	0.452 (3)	0.5052 (12)	0.076 (7)*
N3	0.47595 (4)	0.7010 (3)	0.58328 (13)	0.0615 (5)
H3N	0.4725 (6)	0.599 (3)	0.5412 (11)	0.080 (7)*
N4	0.53909 (4)	0.6575 (3)	0.56996 (12)	0.0632 (5)
C1	0.27020 (6)	0.0330 (4)	0.31870 (14)	0.0648 (6)
H1	0.2536	-0.0903	0.3102	0.078*
C2	0.29845 (6)	0.0538 (4)	0.26093 (14)	0.0717 (6)
H2	0.3009	-0.0517	0.2146	0.086*
C3	0.32289 (6)	0.2357 (4)	0.27402 (15)	0.0682 (6)
H3	0.3424	0.2542	0.2364	0.082*
C4	0.31870 (5)	0.3903 (4)	0.34224 (14)	0.0597 (5)
H4	0.3352	0.5141	0.3513	0.072*
C5	0.28904 (5)	0.3579 (3)	0.39817 (12)	0.0491 (4)
C6	0.30779 (5)	0.6811 (3)	0.49945 (14)	0.0566 (5)
H6A	0.2950	0.7808	0.5398	0.068*
H6B	0.3144	0.7747	0.4490	0.068*
C7	0.34377 (5)	0.5957 (3)	0.54853 (12)	0.0469 (4)

C8	0.34573 (6)	0.3867 (3)	0.59374 (13)	0.0566 (5)
H8	0.3249	0.2898	0.5912	0.068*
C9	0.37829 (6)	0.3207 (3)	0.64242 (15)	0.0630 (6)
H9	0.3792	0.1803	0.6728	0.076*
C10	0.40950 (5)	0.4621 (3)	0.64621 (13)	0.0601 (5)
H10	0.4313	0.4171	0.6796	0.072*
C11	0.40859 (5)	0.6714 (3)	0.60048 (12)	0.0496 (5)
C12	0.37567 (5)	0.7345 (3)	0.55229 (12)	0.0480 (4)
H12	0.3748	0.8741	0.5214	0.058*
C13	0.44274 (5)	0.8269 (3)	0.60434 (15)	0.0573 (5)
H13A	0.4384	0.9527	0.5619	0.069*
H13B	0.4466	0.8928	0.6641	0.069*
C14	0.51105 (5)	0.7920 (3)	0.59598 (13)	0.0525 (5)
C15	0.51840 (5)	1.0098 (3)	0.63396 (13)	0.0600 (5)
H15	0.4987	1.1056	0.6482	0.072*
C16	0.55490 (6)	1.0800 (4)	0.64988 (14)	0.0664 (6)
H16	0.5602	1.2243	0.6751	0.080*
C17	0.58374 (6)	0.9363 (4)	0.62849 (15)	0.0667 (6)
H17	0.6088	0.9778	0.6412	0.080*
C18	0.57426 (6)	0.7309 (4)	0.58794 (16)	0.0681 (6)
H18	0.5937	0.6357	0.5718	0.082*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
N1	0.0488 (9)	0.0608 (10)	0.0551 (10)	−0.0034 (7)	0.0003 (7)	−0.0014 (8)
N2	0.0473 (9)	0.0641 (10)	0.0604 (10)	−0.0077 (8)	0.0084 (8)	−0.0103 (9)
N3	0.0474 (9)	0.0637 (11)	0.0737 (12)	−0.0021 (8)	0.0062 (8)	−0.0216 (9)
N4	0.0485 (10)	0.0596 (10)	0.0820 (12)	−0.0003 (7)	0.0071 (9)	−0.0096 (9)
C1	0.0689 (13)	0.0630 (12)	0.0615 (12)	−0.0010 (10)	−0.0031 (11)	−0.0067 (10)
C2	0.0773 (15)	0.0803 (15)	0.0573 (13)	0.0147 (12)	0.0043 (12)	−0.0080 (11)
C3	0.0612 (13)	0.0913 (15)	0.0530 (12)	0.0132 (12)	0.0111 (10)	0.0059 (12)
C4	0.0474 (11)	0.0717 (12)	0.0603 (12)	−0.0037 (9)	0.0058 (9)	0.0065 (10)
C5	0.0391 (9)	0.0574 (10)	0.0503 (10)	0.0008 (8)	−0.0006 (8)	0.0036 (9)
C6	0.0507 (11)	0.0534 (11)	0.0649 (12)	−0.0010 (9)	−0.0013 (9)	−0.0014 (9)
C7	0.0471 (10)	0.0456 (10)	0.0484 (10)	0.0008 (8)	0.0054 (8)	−0.0038 (8)
C8	0.0578 (12)	0.0499 (11)	0.0628 (12)	−0.0049 (9)	0.0086 (10)	0.0036 (9)
C9	0.0671 (13)	0.0556 (12)	0.0666 (13)	0.0046 (10)	0.0052 (11)	0.0162 (10)
C10	0.0543 (12)	0.0654 (12)	0.0600 (12)	0.0093 (10)	−0.0009 (10)	0.0063 (10)
C11	0.0487 (11)	0.0519 (10)	0.0486 (10)	0.0042 (8)	0.0065 (9)	−0.0029 (8)
C12	0.0507 (10)	0.0437 (9)	0.0500 (10)	0.0027 (8)	0.0056 (8)	0.0009 (8)
C13	0.0468 (11)	0.0598 (11)	0.0648 (12)	0.0014 (9)	0.0011 (9)	−0.0083 (10)
C14	0.0478 (11)	0.0531 (11)	0.0563 (11)	−0.0012 (8)	0.0014 (9)	−0.0037 (9)
C15	0.0580 (12)	0.0596 (12)	0.0618 (12)	−0.0011 (9)	0.0008 (10)	−0.0120 (10)
C16	0.0708 (14)	0.0676 (13)	0.0598 (12)	−0.0146 (11)	−0.0022 (11)	−0.0083 (10)
C17	0.0519 (12)	0.0822 (14)	0.0650 (13)	−0.0120 (11)	−0.0034 (10)	0.0022 (11)
C18	0.0488 (12)	0.0708 (13)	0.0848 (16)	0.0002 (10)	0.0050 (11)	−0.0039 (12)

## Geometric parameters (Å, °)

N1—C1	1.335 (2)	C7—C8	1.384 (2)
N1—C5	1.347 (2)	C7—C12	1.390 (2)
N2—C5	1.365 (2)	C8—C9	1.378 (3)
N2—C6	1.442 (2)	C8—H8	0.9300
N2—H2N	0.86 (1)	C9—C10	1.379 (3)
N3—C14	1.358 (2)	C9—H9	0.9300
N3—C13	1.444 (2)	C10—C11	1.389 (2)
N3—H3N	0.86 (1)	C10—H10	0.9300
N4—C18	1.333 (2)	C11—C12	1.381 (3)
N4—C14	1.346 (2)	C11—C13	1.512 (2)
C1—C2	1.376 (3)	C12—H12	0.9300
C1—H1	0.9300	C13—H13A	0.9700
C2—C3	1.372 (3)	C13—H13B	0.9700
C2—H2	0.9300	C14—C15	1.399 (3)
C3—C4	1.371 (3)	C15—C16	1.368 (3)
C3—H3	0.9300	C15—H15	0.9300
C4—C5	1.405 (2)	C16—C17	1.377 (3)
C4—H4	0.9300	C16—H16	0.9300
C6—C7	1.514 (3)	C17—C18	1.366 (3)
C6—H6A	0.9700	C17—H17	0.9300
C6—H6B	0.9700	C18—H18	0.9300
C1—N1—C5	117.91 (15)	C7—C8—H8	119.7
C5—N2—C6	122.91 (15)	C8—C9—C10	120.23 (18)
C5—N2—H2N	115.5 (15)	C8—C9—H9	119.9
C6—N2—H2N	115.1 (15)	C10—C9—H9	119.9
C14—N3—C13	122.39 (16)	C9—C10—C11	120.47 (19)
C14—N3—H3N	116.7 (14)	C9—C10—H10	119.8
C13—N3—H3N	115.2 (15)	C11—C10—H10	119.8
C18—N4—C14	117.70 (17)	C12—C11—C10	118.35 (17)
N1—C1—C2	124.1 (2)	C12—C11—C13	120.98 (16)
N1—C1—H1	118.0	C10—C11—C13	120.67 (17)
C2—C1—H1	118.0	C11—C12—C7	122.05 (16)
C3—C2—C1	117.6 (2)	C11—C12—H12	119.0
C3—C2—H2	121.2	C7—C12—H12	119.0
C1—C2—H2	121.2	N3—C13—C11	111.21 (15)
C4—C3—C2	120.34 (18)	N3—C13—H13A	109.4
C4—C3—H3	119.8	C11—C13—H13A	109.4
C2—C3—H3	119.8	N3—C13—H13B	109.4
C3—C4—C5	118.65 (19)	C11—C13—H13B	109.4
C3—C4—H4	120.7	H13A—C13—H13B	108.0
C5—C4—H4	120.7	N4—C14—N3	115.54 (16)
N1—C5—N2	115.33 (14)	N4—C14—C15	121.11 (17)
N1—C5—C4	121.35 (17)	N3—C14—C15	123.36 (16)
N2—C5—C4	123.30 (17)	C16—C15—C14	119.09 (18)
N2—C6—C7	116.16 (15)	C16—C15—H15	120.5

N2—C6—H6A	108.2	C14—C15—H15	120.5
C7—C6—H6A	108.2	C15—C16—C17	119.81 (19)
N2—C6—H6B	108.2	C15—C16—H16	120.1
C7—C6—H6B	108.2	C17—C16—H16	120.1
H6A—C6—H6B	107.4	C18—C17—C16	117.59 (19)
C8—C7—C12	118.21 (17)	C18—C17—H17	121.2
C8—C7—C6	122.17 (16)	C16—C17—H17	121.2
C12—C7—C6	119.55 (15)	N4—C18—C17	124.49 (18)
C9—C8—C7	120.67 (17)	N4—C18—H18	117.8
C9—C8—H8	119.7	C17—C18—H18	117.8
C5—N1—C1—C2	0.4 (3)	C9—C10—C11—C13	-179.93 (17)
N1—C1—C2—C3	0.2 (3)	C10—C11—C12—C7	0.1 (3)
C1—C2—C3—C4	-0.4 (3)	C13—C11—C12—C7	-179.27 (16)
C2—C3—C4—C5	0.1 (3)	C8—C7—C12—C11	-1.0 (3)
C1—N1—C5—N2	177.71 (17)	C6—C7—C12—C11	176.15 (16)
C1—N1—C5—C4	-0.7 (3)	C14—N3—C13—C11	-169.32 (18)
C6—N2—C5—N1	168.25 (17)	C12—C11—C13—N3	-130.03 (18)
C6—N2—C5—C4	-13.4 (3)	C10—C11—C13—N3	50.6 (2)
C3—C4—C5—N1	0.4 (3)	C18—N4—C14—N3	-174.51 (19)
C3—C4—C5—N2	-177.85 (19)	C18—N4—C14—C15	5.1 (3)
C5—N2—C6—C7	-67.6 (2)	C13—N3—C14—N4	-177.51 (17)
N2—C6—C7—C8	-27.6 (2)	C13—N3—C14—C15	2.9 (3)
N2—C6—C7—C12	155.37 (16)	N4—C14—C15—C16	-4.1 (3)
C12—C7—C8—C9	1.2 (3)	N3—C14—C15—C16	175.4 (2)
C6—C7—C8—C9	-175.93 (18)	C14—C15—C16—C17	0.0 (3)
C7—C8—C9—C10	-0.4 (3)	C15—C16—C17—C18	2.9 (3)
C8—C9—C10—C11	-0.6 (3)	C14—N4—C18—C17	-2.1 (3)
C9—C10—C11—C12	0.7 (3)	C16—C17—C18—N4	-1.9 (3)

---

*Hydrogen-bond geometry (Å, °)*

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N2—H2N...N1 <sup>i</sup>	0.86 (1)	2.19 (1)	3.047 (2)	175 (2)
N3—H3N...N4 <sup>ii</sup>	0.86 (1)	2.24 (1)	3.104 (2)	177 (2)

Symmetry codes: (i)  $-x+1/2, -y+1/2, -z+1$ ; (ii)  $-x+1, -y+1, -z+1$ .