

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-(Biphenyl-4-carbonyl)-N'-(4-chlorophenyl)thiourea

Bohari M. Yamin and M. Asyikin M. Arif*

School of Chemical Sciences and Food Technology, Universiti Kebangsaan Malaysia, UKM 43500 Bangi Selangor, Malaysia

Correspondence e-mail: mayaasyikin_ns@yahoo.com


Received 16 November 2007; accepted 22 November 2007

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.049; wR factor = 0.122; data-to-parameter ratio = 15.4.

In the title compound, $\text{C}_{20}\text{H}_{15}\text{ClN}_2\text{OS}$, the benzene rings of the biphenyl group are at an angle of $44.23(12)^\circ$. The $\text{C}_4\text{N}_2\text{OS}$ central thiourea fragment makes dihedral angles with the benzene carbonyl and chlorobenzene rings of $55.96(9)$ and $64.09(9)^\circ$, respectively. The *trans-cis* geometry of the thiourea group is stabilized by the intramolecular hydrogen bond between the carbonyl O atom and the H atom of the *cis*-thioamide. In the crystal structure, molecules are linked by $\text{N}-\text{H}\cdots\text{S}$ and $\text{N}-\text{H}\cdots\text{O}$ intermolecular hydrogen bonds to form one-dimensional chains along the c axis. $\text{C}-\text{H}\cdots\pi$ interactions also contribute to the stability of the molecule.

Related literature

For related literature, see: Allen *et al.* (1987); Arif & Yamin (2007).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{15}\text{ClN}_2\text{OS}$
 $M_r = 366.85$
 Monoclinic, $P2_1/c$
 $a = 16.039(7)$ Å
 $b = 6.087(3)$ Å
 $c = 18.096(8)$ Å
 $\beta = 94.780(8)^\circ$

$V = 1760.5(14)$ Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.35$ mm⁻¹
 $T = 298(2)$ K
 $0.49 \times 0.46 \times 0.10$ mm

Data collection

Bruker SMART APEX CCD
 area-detector diffractometer
 Absorption correction: multi-scan
 (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.848$, $T_{\max} = 0.966$

9371 measured reflections
 3475 independent reflections
 2278 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.122$
 $S = 1.02$
 3475 reflections

226 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.28$ e Å⁻³
 $\Delta\rho_{\min} = -0.22$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

| $D-\text{H}\cdots A$ | $D-\text{H}$ | $\text{H}\cdots A$ | $D\cdots A$ | $D-\text{H}\cdots A$ |
|--|--------------|--------------------|-------------|----------------------|
| $\text{N2}-\text{H2}\cdots\text{O1}$ | 0.86 | 2.00 | 2.683 (3) | 135 |
| $\text{N1}-\text{H1}\cdots\text{S1}^{\text{i}}$ | 0.86 | 2.55 | 3.362 (3) | 157 |
| $\text{N2}-\text{H2}\cdots\text{O1}^{\text{ii}}$ | 0.86 | 2.58 | 3.210 (3) | 131 |
| $\text{Cl1}-\text{H1A}\cdots\text{Cg3}^{\text{iii}}$ | 0.93 | 2.98 | 3.586 (3) | 124 |

Symmetry codes: (i) $-x + 2, -y + 1, -z + 1$; (ii) $-x + 2, y, -z + \frac{1}{2}$. Cg3 is the centroid of atoms $\text{C15}-\text{C20}$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINTE* (Bruker, 2000); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997a); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997a); molecular graphics: *SHELXTL* (Sheldrick, 1997b); software used to prepare material for publication: *SHELXTL*, *PARST* (Nardelli, 1995) and *PLATON* (Spek, 2003).

The authors thank the Ministry of Higher Education for research grant No. UKM-ST-01-FRGS-0003-2006, and the Universiti Kebangsaan Malaysia and the Universiti Malaysia Sarawak for the facilities and scholarship to MAMA.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2027).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Arif, M. A. M. & Yamin, B. M. (2007). *Acta Cryst. E* **63**, o3594.
 Bruker (2000). *SADABS* (Version 2.01), *SMART* (Version 5.630) and *SAINTE* (Version 6.36a). Bruker AXS Inc., Madison, Wisconsin, USA.
 Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
 Sheldrick, G. M. (1997a). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
 Sheldrick, G. M. (1997b). *SHELXTL*. Version 5.1. Bruker AXS Inc., Madison, Wisconsin, USA.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supporting information

Acta Cryst. (2008). E64, o104 [https://doi.org/10.1107/S1600536807062265]

N*-(Biphenyl-4-carbonyl)-*N'*-(4-chlorophenyl)thiourea*Bohari M. Yamin and M. Asyikin M. Arif****S1. Comment**

The title compound (Fig. 1) is an isomeric analog of the previously reported *N*-(biphenyl-4-carbonyl)-*N'*-(2-chlorophenyl) thiourea (II) (Arif and Yamin, 2007). The dihedral angle between the two benzene rings in the biphenyl fragment is 44.23 (12)°, which is double the value of 20.71 (17)° in (II). The examination on the planarity of the central thiourea fragment S1/N1/N2/C14 and the chlorophenyl plane (C15—C20)/C11, indicates that they are planar. The central thiourea fragment makes dihedral angles with the benzene carbonyl and chlorobenzene rings of 55.96 (9) and 64.09 (9)°, respectively. The *trans-cis* geometry in the thiourea moiety is stabilized by the N2—H2···O1 intramolecular hydrogen bond (Table 1).


In the crystal structure symmetry related molecules are linked by N1—H1···S1ⁱ and N2—H2···O1ⁱⁱ intermolecular hydrogen bonds to form one-dimensional chains along the *c* axis (Fig. 2 and Table 1). The molecule is also stabilized by a C1—H1A··· π interaction; the distance between H1A and the (C15—C20) ring centroid is 2.98 Å, and the angle about the hydrogen atom is 124°.

S2. Experimental


A solution of 4-chloroaniline (0.63 g, 2.5 mmol) in 20 ml acetone was added dropwise to a two-necked round-bottomed flask containing an equimolar amount of biphenylcarbomoylthiocyanate (0.60 g, 2.5 mmol) in 20 ml of acetone. The mixture was refluxed for about 3 h. The light yellow solution was filtered and the filtrate allowed to evaporate at room temperature. Colourless crystals were obtained after five days (yield 0.71 g, 85%, m.p.: 164–166°C).

S3. Refinement

H atoms on C and N atoms were positioned geometrically with C—H = 0.93 and N—H = 0.86 Å, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{parent atom})$.

**Figure 1**

Molecular structure of compound (I), with displacement ellipsoid drawn at the 50% probability level. The dashed line indicates the intramolecular hydrogen bond.

**Figure 2**

A view along the *b* axis of the crystal packing of compound (I). The dashed line indicates the intermolecular N—H...S and N—H...O hydrogen bonds (see Table 1 for details).

N-(Biphenyl-4-carbonyl)-*N'*-(4-chlorophenyl)thiourea*Crystal data*C₂₀H₁₅ClN₂OS $M_r = 366.85$ Monoclinic, *P*2₁/*c*Hall symbol: -*P* 2₁*c* $a = 16.039$ (7) Å $b = 6.087$ (3) Å $c = 18.096$ (8) Å $\beta = 94.780$ (8)° $V = 1760.5$ (14) Å³ $Z = 4$ $F(000) = 760$ $D_x = 1.384$ Mg m⁻³

Melting point: 164–166°C K

Mo *K*α radiation, $\lambda = 0.71073$ Å

Cell parameters from 1713 reflections

 $\theta = 2.2$ – 26.0 ° $\mu = 0.35$ mm⁻¹ $T = 298$ K

Block, colourless

 $0.49 \times 0.46 \times 0.10$ mm*Data collection*Bruker SMART APEX CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 ω scan

Absorption correction: multi-scan

(SADABS; Bruker, 2000)

 $T_{\min} = 0.848$, $T_{\max} = 0.966$

9371 measured reflections

3475 independent reflections

2278 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.037$ $\theta_{\max} = 26.0$ °, $\theta_{\min} = 2.2$ ° $h = -19 \rightarrow 18$ $k = -7 \rightarrow 7$ $l = -22 \rightarrow 12$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.049$ $wR(F^2) = 0.122$ $S = 1.02$

3475 reflections

226 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0596P)^2 + 0.0946P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.28$ e Å⁻³ $\Delta\rho_{\min} = -0.22$ e Å⁻³*Special details*

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

| | <i>x</i> | <i>y</i> | <i>z</i> | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|-----|--------------|--------------|--------------|----------------------------------|
| Cl1 | 1.37861 (5) | 0.39472 (17) | 0.18867 (5) | 0.1065 (4) |
| S1 | 1.08379 (4) | 0.38002 (11) | 0.42312 (3) | 0.0553 (2) |
| O1 | 0.95079 (10) | 0.9893 (3) | 0.32501 (9) | 0.0591 (5) |
| N1 | 0.97515 (11) | 0.7034 (3) | 0.40513 (10) | 0.0483 (5) |

| | | | | |
|-----|--------------|------------|--------------|------------|
| H1 | 0.9564 | 0.6440 | 0.4435 | 0.058* |
| N2 | 1.07779 (10) | 0.7027 (3) | 0.32382 (10) | 0.0487 (5) |
| H2 | 1.0537 | 0.8194 | 0.3057 | 0.058* |
| C1 | 0.79626 (14) | 0.7695 (4) | 0.42417 (12) | 0.0505 (6) |
| H1A | 0.8026 | 0.6333 | 0.4020 | 0.061* |
| C2 | 0.72746 (13) | 0.8084 (4) | 0.46287 (13) | 0.0504 (6) |
| H2A | 0.6872 | 0.6993 | 0.4656 | 0.060* |
| C3 | 0.71742 (13) | 1.0086 (4) | 0.49788 (12) | 0.0445 (6) |
| C4 | 0.64556 (14) | 1.0439 (4) | 0.54340 (13) | 0.0481 (6) |
| C5 | 0.62285 (15) | 0.8839 (4) | 0.59148 (14) | 0.0601 (7) |
| H5 | 0.6532 | 0.7538 | 0.5957 | 0.072* |
| C6 | 0.55587 (17) | 0.9127 (5) | 0.63367 (16) | 0.0752 (9) |
| H6 | 0.5416 | 0.8032 | 0.6661 | 0.090* |
| C7 | 0.51023 (17) | 1.1049 (6) | 0.62738 (16) | 0.0749 (9) |
| H7 | 0.4645 | 1.1246 | 0.6550 | 0.090* |
| C8 | 0.53267 (16) | 1.2669 (5) | 0.58009 (16) | 0.0759 (9) |
| H8 | 0.5026 | 1.3976 | 0.5763 | 0.091* |
| C9 | 0.59943 (15) | 1.2364 (5) | 0.53843 (15) | 0.0627 (7) |
| H9 | 0.6139 | 1.3468 | 0.5064 | 0.075* |
| C10 | 0.77729 (14) | 1.1700 (4) | 0.49070 (13) | 0.0525 (6) |
| H10 | 0.7713 | 1.3058 | 0.5132 | 0.063* |
| C11 | 0.84582 (14) | 1.1326 (4) | 0.45067 (13) | 0.0519 (6) |
| H11 | 0.8850 | 1.2432 | 0.4458 | 0.062* |
| C12 | 0.85579 (13) | 0.9293 (4) | 0.41789 (12) | 0.0431 (5) |
| C13 | 0.93135 (13) | 0.8825 (4) | 0.37742 (12) | 0.0451 (6) |
| C14 | 1.04533 (13) | 0.6039 (4) | 0.38029 (12) | 0.0425 (5) |
| C15 | 1.15067 (13) | 0.6247 (4) | 0.29167 (11) | 0.0426 (5) |
| C16 | 1.22033 (14) | 0.7565 (5) | 0.29525 (12) | 0.0572 (7) |
| H16 | 1.2199 | 0.8927 | 0.3185 | 0.069* |
| C17 | 1.29127 (15) | 0.6842 (5) | 0.26385 (15) | 0.0679 (8) |
| H17 | 1.3389 | 0.7719 | 0.2655 | 0.081* |
| C18 | 1.29059 (15) | 0.4827 (5) | 0.23047 (14) | 0.0605 (7) |
| C19 | 1.22169 (16) | 0.3514 (4) | 0.22703 (14) | 0.0594 (7) |
| H19 | 1.2226 | 0.2141 | 0.2046 | 0.071* |
| C20 | 1.15066 (15) | 0.4242 (4) | 0.25714 (13) | 0.0530 (6) |
| H20 | 1.1027 | 0.3376 | 0.2541 | 0.064* |

Atomic displacement parameters (\AA^2)

| | U^{11} | U^{22} | U^{33} | U^{12} | U^{13} | U^{23} |
|-----|-------------|-------------|-------------|--------------|-------------|--------------|
| C11 | 0.0737 (5) | 0.1498 (9) | 0.1022 (7) | 0.0428 (5) | 0.0449 (5) | 0.0262 (6) |
| S1 | 0.0576 (4) | 0.0604 (4) | 0.0499 (4) | 0.0152 (3) | 0.0171 (3) | 0.0127 (3) |
| O1 | 0.0684 (11) | 0.0584 (11) | 0.0528 (10) | 0.0138 (9) | 0.0183 (8) | 0.0152 (9) |
| N1 | 0.0496 (11) | 0.0554 (13) | 0.0417 (11) | 0.0119 (10) | 0.0137 (9) | 0.0092 (9) |
| N2 | 0.0495 (11) | 0.0530 (12) | 0.0454 (11) | 0.0088 (9) | 0.0138 (9) | 0.0096 (10) |
| C1 | 0.0504 (14) | 0.0455 (15) | 0.0549 (15) | 0.0053 (12) | 0.0007 (11) | -0.0092 (12) |
| C2 | 0.0434 (13) | 0.0483 (15) | 0.0594 (15) | -0.0066 (11) | 0.0039 (11) | -0.0050 (12) |
| C3 | 0.0415 (12) | 0.0462 (15) | 0.0456 (13) | 0.0027 (11) | 0.0029 (10) | 0.0012 (11) |

| | | | | | | |
|-----|-------------|-------------|-------------|--------------|-------------|--------------|
| C4 | 0.0419 (13) | 0.0507 (15) | 0.0519 (14) | -0.0052 (11) | 0.0046 (10) | -0.0057 (12) |
| C5 | 0.0554 (15) | 0.0614 (18) | 0.0644 (17) | -0.0039 (13) | 0.0110 (13) | 0.0014 (14) |
| C6 | 0.0655 (18) | 0.090 (2) | 0.0726 (19) | -0.0176 (17) | 0.0198 (15) | 0.0045 (17) |
| C7 | 0.0504 (16) | 0.100 (3) | 0.077 (2) | -0.0100 (17) | 0.0222 (14) | -0.0116 (19) |
| C8 | 0.0542 (17) | 0.084 (2) | 0.091 (2) | 0.0153 (15) | 0.0165 (15) | -0.0057 (18) |
| C9 | 0.0575 (16) | 0.0625 (18) | 0.0699 (18) | 0.0081 (14) | 0.0166 (13) | 0.0053 (14) |
| C10 | 0.0577 (15) | 0.0411 (14) | 0.0606 (16) | 0.0008 (12) | 0.0165 (12) | -0.0076 (12) |
| C11 | 0.0524 (14) | 0.0462 (15) | 0.0590 (15) | -0.0046 (12) | 0.0157 (11) | -0.0034 (12) |
| C12 | 0.0420 (12) | 0.0461 (14) | 0.0411 (13) | 0.0065 (11) | 0.0038 (10) | 0.0020 (11) |
| C13 | 0.0470 (13) | 0.0482 (15) | 0.0399 (13) | 0.0018 (11) | 0.0017 (10) | -0.0025 (12) |
| C14 | 0.0403 (12) | 0.0489 (14) | 0.0390 (12) | 0.0017 (11) | 0.0066 (9) | 0.0001 (11) |
| C15 | 0.0441 (12) | 0.0497 (14) | 0.0347 (12) | 0.0004 (11) | 0.0077 (9) | 0.0060 (11) |
| C16 | 0.0613 (16) | 0.0607 (18) | 0.0505 (15) | -0.0115 (13) | 0.0101 (12) | -0.0079 (13) |
| C17 | 0.0453 (15) | 0.098 (2) | 0.0610 (17) | -0.0151 (15) | 0.0092 (12) | 0.0054 (17) |
| C18 | 0.0505 (15) | 0.083 (2) | 0.0497 (15) | 0.0137 (15) | 0.0154 (12) | 0.0120 (15) |
| C19 | 0.0757 (18) | 0.0522 (16) | 0.0529 (15) | 0.0090 (14) | 0.0196 (13) | 0.0009 (13) |
| C20 | 0.0537 (15) | 0.0583 (17) | 0.0487 (14) | -0.0076 (12) | 0.0138 (11) | -0.0026 (13) |

Geometric parameters (Å, °)

| | | | |
|------------|-------------|-------------|-----------|
| C11—C18 | 1.741 (2) | C6—H6 | 0.9300 |
| S1—C14 | 1.661 (2) | C7—C8 | 1.374 (4) |
| O1—C13 | 1.212 (3) | C7—H7 | 0.9300 |
| N1—C13 | 1.369 (3) | C8—C9 | 1.372 (3) |
| N1—C14 | 1.386 (3) | C8—H8 | 0.9300 |
| N1—H1 | 0.8600 | C9—H9 | 0.9300 |
| N2—C14 | 1.329 (3) | C10—C11 | 1.385 (3) |
| N2—C15 | 1.430 (3) | C10—H10 | 0.9300 |
| N2—H2 | 0.8600 | C11—C12 | 1.387 (3) |
| C1—C12 | 1.374 (3) | C11—H11 | 0.9300 |
| C1—C2 | 1.376 (3) | C12—C13 | 1.495 (3) |
| C1—H1A | 0.9300 | C15—C20 | 1.371 (3) |
| C2—C3 | 1.389 (3) | C15—C16 | 1.373 (3) |
| C2—H2A | 0.9300 | C16—C17 | 1.385 (3) |
| C3—C10 | 1.387 (3) | C16—H16 | 0.9300 |
| C3—C4 | 1.487 (3) | C17—C18 | 1.367 (4) |
| C4—C5 | 1.375 (3) | C17—H17 | 0.9300 |
| C4—C9 | 1.385 (3) | C18—C19 | 1.361 (4) |
| C5—C6 | 1.380 (3) | C19—C20 | 1.376 (3) |
| C5—H5 | 0.9300 | C19—H19 | 0.9300 |
| C6—C7 | 1.380 (4) | C20—H20 | 0.9300 |
| C13—N1—C14 | 129.43 (18) | C11—C10—C3 | 121.2 (2) |
| C13—N1—H1 | 115.3 | C11—C10—H10 | 119.4 |
| C14—N1—H1 | 115.3 | C3—C10—H10 | 119.4 |
| C14—N2—C15 | 123.29 (19) | C10—C11—C12 | 119.7 (2) |
| C14—N2—H2 | 118.4 | C10—C11—H11 | 120.2 |
| C15—N2—H2 | 118.4 | C12—C11—H11 | 120.2 |

| | | | |
|----------------|------------|-----------------|--------------|
| C12—C1—C2 | 120.8 (2) | C1—C12—C11 | 119.4 (2) |
| C12—C1—H1A | 119.6 | C1—C12—C13 | 120.2 (2) |
| C2—C1—H1A | 119.6 | C11—C12—C13 | 120.4 (2) |
| C1—C2—C3 | 120.8 (2) | O1—C13—N1 | 123.8 (2) |
| C1—C2—H2A | 119.6 | O1—C13—C12 | 123.6 (2) |
| C3—C2—H2A | 119.6 | N1—C13—C12 | 112.62 (19) |
| C10—C3—C2 | 118.1 (2) | N2—C14—N1 | 115.62 (19) |
| C10—C3—C4 | 121.6 (2) | N2—C14—S1 | 125.19 (16) |
| C2—C3—C4 | 120.3 (2) | N1—C14—S1 | 119.17 (16) |
| C5—C4—C9 | 118.2 (2) | C20—C15—C16 | 120.7 (2) |
| C5—C4—C3 | 120.5 (2) | C20—C15—N2 | 120.8 (2) |
| C9—C4—C3 | 121.4 (2) | C16—C15—N2 | 118.5 (2) |
| C4—C5—C6 | 121.3 (3) | C15—C16—C17 | 119.2 (3) |
| C4—C5—H5 | 119.3 | C15—C16—H16 | 120.4 |
| C6—C5—H5 | 119.3 | C17—C16—H16 | 120.4 |
| C7—C6—C5 | 119.6 (3) | C18—C17—C16 | 119.3 (2) |
| C7—C6—H6 | 120.2 | C18—C17—H17 | 120.3 |
| C5—C6—H6 | 120.2 | C16—C17—H17 | 120.3 |
| C8—C7—C6 | 119.7 (3) | C19—C18—C17 | 121.6 (2) |
| C8—C7—H7 | 120.1 | C19—C18—C11 | 119.0 (2) |
| C6—C7—H7 | 120.1 | C17—C18—C11 | 119.4 (2) |
| C9—C8—C7 | 120.1 (3) | C18—C19—C20 | 119.3 (3) |
| C9—C8—H8 | 119.9 | C18—C19—H19 | 120.4 |
| C7—C8—H8 | 119.9 | C20—C19—H19 | 120.4 |
| C8—C9—C4 | 121.1 (3) | C15—C20—C19 | 119.9 (2) |
| C8—C9—H9 | 119.5 | C15—C20—H20 | 120.1 |
| C4—C9—H9 | 119.5 | C19—C20—H20 | 120.1 |
| | | | |
| C12—C1—C2—C3 | -1.4 (3) | C14—N1—C13—O1 | -2.3 (4) |
| C1—C2—C3—C10 | 2.0 (3) | C14—N1—C13—C12 | 176.5 (2) |
| C1—C2—C3—C4 | -176.2 (2) | C1—C12—C13—O1 | 123.0 (3) |
| C10—C3—C4—C5 | -134.7 (3) | C11—C12—C13—O1 | -58.3 (3) |
| C2—C3—C4—C5 | 43.4 (3) | C1—C12—C13—N1 | -55.8 (3) |
| C10—C3—C4—C9 | 45.7 (3) | C11—C12—C13—N1 | 122.9 (2) |
| C2—C3—C4—C9 | -136.2 (2) | C15—N2—C14—N1 | 179.21 (19) |
| C9—C4—C5—C6 | 0.2 (4) | C15—N2—C14—S1 | 1.0 (3) |
| C3—C4—C5—C6 | -179.4 (2) | C13—N1—C14—N2 | 4.0 (3) |
| C4—C5—C6—C7 | 0.4 (4) | C13—N1—C14—S1 | -177.63 (18) |
| C5—C6—C7—C8 | -1.0 (4) | C14—N2—C15—C20 | 63.7 (3) |
| C6—C7—C8—C9 | 1.0 (5) | C14—N2—C15—C16 | -117.2 (3) |
| C7—C8—C9—C4 | -0.4 (4) | C20—C15—C16—C17 | -0.4 (4) |
| C5—C4—C9—C8 | -0.2 (4) | N2—C15—C16—C17 | -179.5 (2) |
| C3—C4—C9—C8 | 179.4 (2) | C15—C16—C17—C18 | -0.5 (4) |
| C2—C3—C10—C11 | -0.8 (4) | C16—C17—C18—C19 | 0.3 (4) |
| C4—C3—C10—C11 | 177.3 (2) | C16—C17—C18—C11 | 178.19 (19) |
| C3—C10—C11—C12 | -0.9 (4) | C17—C18—C19—C20 | 0.7 (4) |
| C2—C1—C12—C11 | -0.4 (3) | C11—C18—C19—C20 | -177.18 (19) |
| C2—C1—C12—C13 | 178.3 (2) | C16—C15—C20—C19 | 1.4 (4) |

| | | | |
|-----------------|------------|-----------------|------------|
| C10—C11—C12—C1 | 1.6 (3) | N2—C15—C20—C19 | -179.5 (2) |
| C10—C11—C12—C13 | -177.2 (2) | C18—C19—C20—C15 | -1.6 (4) |

Hydrogen-bond geometry (Å, °)

| <i>D</i> —H \cdots <i>A</i> | <i>D</i> —H | H \cdots <i>A</i> | <i>D</i> \cdots <i>A</i> | <i>D</i> —H \cdots <i>A</i> |
|-----------------------------------|-------------|---------------------|----------------------------|-------------------------------|
| N2—H2 \cdots O1 | 0.86 | 2.00 | 2.683 (3) | 135 |
| N1—H1 \cdots S1 ⁱ | 0.86 | 2.55 | 3.362 (3) | 157 |
| N2—H2 \cdots O1 ⁱⁱ | 0.86 | 2.58 | 3.210 (3) | 131 |
| C1—H1A \cdots Cg3 ⁱⁱ | 0.93 | 2.98 | 3.586 (3) | 124 |

Symmetry codes: (i) $-x+2, -y+1, -z+1$; (ii) $-x+2, y, -z+1/2$.