

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## Ethyl 6-(2-chlorophenyl)-4-methyl-1-(3-oxobutyl)-2-thioxo-1,2,3,6-tetrahydropyrimidine-5-carboxylate

Qing-Peng He,\* Jian-Yong Wang and Ruo-Kun Feng

College of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China

Correspondence e-mail: hxqqxh2008@yahoo.com.cn


Received 5 December 2007; accepted 17 December 2007

Key indicators: single-crystal X-ray study;  $T = 273$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å; disorder in main residue;  $R$  factor = 0.055;  $wR$  factor = 0.166; data-to-parameter ratio = 13.4.

In the title molecule,  $\text{C}_{18}\text{H}_{21}\text{ClN}_2\text{O}_3\text{S}$ , the pyrimidine ring exhibits a half-chair conformation. The ethyl group is disordered between two positions in a ratio 0.74:0.26. In the crystal structure, the molecules are linked into chains along the  $a$  axis by  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bonds.

## Related literature

For the crystal structure of a related compound, see Jiang *et al.* (2007).



## Experimental

## Crystal data

$\text{C}_{18}\text{H}_{21}\text{ClN}_2\text{O}_3\text{S}$	$\gamma = 71.296$ (6) $^\circ$
$M_r = 380.88$	$V = 931.7$ (3) Å <sup>3</sup>
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.5227$ (12) Å	Mo $K\alpha$ radiation
$b = 9.7163$ (15) Å	$\mu = 0.34$ mm <sup>-1</sup>
$c = 14.122$ (2) Å	$T = 273$ (2) K
$\alpha = 72.617$ (6) $^\circ$	$0.14 \times 0.12 \times 0.10$ mm
$\beta = 87.300$ (6) $^\circ$	

## Data collection

Bruker SMART CCD area-detector diffractometer	10470 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	3233 independent reflections
$T_{\min} = 0.954$ , $T_{\max} = 0.967$	2743 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.021$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$	63 restraints
$wR(F^2) = 0.166$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{\text{max}} = 0.92$ e Å <sup>-3</sup>
3233 reflections	$\Delta\rho_{\text{min}} = -0.71$ e Å <sup>-3</sup>
242 parameters	

Table 1

Hydrogen-bond geometry (Å,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}^i$	0.86	2.16	2.984 (3)	160

Symmetry code: (i)  $x - 1, y, z$ .

Data collection: SMART (Siemens, 1996); cell refinement: SAINT (Siemens, 1996); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 1997a); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997a); molecular graphics: SHELXTL (Sheldrick, 1997b); software used to prepare material for publication: SHELXTL.

The authors acknowledge the support of the National Natural Science Foundation of Liaocheng University (No. X051040).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV2372).

## References

- Jiang, H., Yu, C.-X., Tu, S.-J., Wang, X.-S. & Yao, C.-S. (2007). *Acta Cryst.* **E63**, o298–o299.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997a). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Sheldrick, G. M. (1997b). *SHELXTL*. Version 5.1. Bruker AXS Inc., Madison, Wisconsin, USA.
- Siemens (1996). *SMART* and *SAINTE*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

## supporting information

*Acta Cryst.* (2008). E64, o352 [doi:10.1107/S160053680706730X]

## Ethyl 6-(2-chlorophenyl)-4-methyl-1-(3-oxobutyl)-2-thioxo-1,2,3,6-tetrahydro-pyrimidine-5-carboxylate

Qing-Peng He, Jian-Yong Wang and Ruo-Kun Feng

### S1. Comment

Herewith we present the crystal structure of the title compound, (I), which was synthesized through the reaction of 2-chlorobenzaldehyde and ethyl acetoacetate with urea derivative under solvent-free conditions.

In (I) (Fig. 1), bond lengths and angles are normal and comparable with those observed in reported compound (Jiang *et al.*, 2007). The pyrimidine ring exhibits a half-chair conformation with the maximal deviation of 0.168 Å for C6 from mean plane.


In the crystal, the molecules related by translation along axis *a* are linked into chains by N—H···O hydrogen bonds (Table 1).

### S2. Experimental

2-Chlorobenzaldehyde (2 mmol), ethyl acetoacetate (2 mmol), urea derivatives (2.4 mmol) and H<sub>3</sub>BO<sub>3</sub> (0.4 mmol), in glacial acetic acid (10 ml) was heated at 373 K, while stirring for 1 h, then cooled to room temperature, and poured into ice water (50 ml), and recrystallized from EtOH, affording the title compound as a colourless crystalline solid. Elemental analysis: calculated for C<sub>18</sub>H<sub>21</sub>ClN<sub>2</sub>O<sub>3</sub>S: C 56.76, H 5.56, N 7.35%; found: C 56.68, H 5.45, N 7.44%.

### S3. Refinement

All H atoms were placed in geometrically idealized positions (N—H 0.86, C—H 0.93–0.97 Å) and treated as riding on their parent atoms, with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}, \text{N})$ . Atoms C16 and C17 were treated as disordered between two positions with refined occupancies of 0.740 (1) and 0.26 (1), respectively.

**Figure 1**

The molecular structure of (I) showing the atomic numbering and 30% probability displacement ellipsoids. Only major parts of disordered atoms is shown. H atoms omitted for clarity.

### Ethyl 6-(2-chlorophenyl)-4-methyl-1-(3-oxobutyl)-2-thioxo-1,2,3,6-tetrahydropyrimidine-5-carboxylate

#### Crystal data

$C_{18}H_{21}ClN_2O_3S$

$M_r = 380.88$

Triclinic,  $P\bar{1}$

$a = 7.5227$  (12) Å

$b = 9.7163$  (15) Å

$c = 14.122$  (2) Å

$\alpha = 72.617$  (6)°

$\beta = 87.300$  (6)°

$\gamma = 71.296$  (6)°

$V = 931.7$  (3) Å<sup>3</sup>

$Z = 2$

$F(000) = 400$

$D_x = 1.358$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 5002 reflections

$\theta = 2.3$ – $27.4$ °

$\mu = 0.34$  mm<sup>-1</sup>

$T = 273$  K

Block, colourless

$0.14 \times 0.12 \times 0.10$  mm

#### Data collection

Bruker SMART CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.954$ ,  $T_{\max} = 0.967$

10470 measured reflections

3233 independent reflections

2743 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.021$

$\theta_{\max} = 25.0$ °,  $\theta_{\min} = 1.5$ °

$h = -8 \rightarrow 8$

$k = -11 \rightarrow 10$

$l = -16 \rightarrow 16$

*Refinement*Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.055$  $wR(F^2) = 0.166$  $S = 1.00$ 

3233 reflections

242 parameters

63 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0856P)^2 + 1.0486P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.002$  $\Delta\rho_{\max} = 0.92 \text{ e } \text{\AA}^{-3}$  $\Delta\rho_{\min} = -0.71 \text{ e } \text{\AA}^{-3}$ Extinction correction: *SHELXL97* (Sheldrick,  
1997a),  $F_c^* = kF_c[1 + 0.001xF_c^2\lambda^3/\sin(2\theta)]^{-1/4}$ 

Extinction coefficient: 0.019 (5)

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
S1	0.25980 (13)	0.52300 (10)	0.37239 (6)	0.0576 (3)	
C1	0.7929 (7)	0.7860 (5)	0.2049 (3)	0.0787 (12)	
H1A	0.8645	0.7990	0.1467	0.094*	
H1B	0.8473	0.8105	0.2553	0.094*	
H1C	0.6657	0.8523	0.1885	0.094*	
O2	0.8603 (4)	0.1939 (5)	0.0855 (2)	0.0915 (11)	
O3	0.5995 (5)	0.2294 (6)	0.0050 (3)	0.1280 (14)	
N1	0.3010 (3)	0.4024 (3)	0.22578 (18)	0.0459 (6)	
H1	0.1819	0.4188	0.2294	0.055*	
N2	0.5778 (3)	0.3881 (3)	0.29602 (17)	0.0409 (6)	
O1	0.8827 (3)	0.5317 (3)	0.2018 (2)	0.0647 (7)	
C2	0.7951 (4)	0.6252 (4)	0.2424 (2)	0.0507 (7)	
C3	0.6906 (5)	0.5828 (4)	0.3335 (2)	0.0505 (7)	
H3A	0.5621	0.6511	0.3218	0.061*	
H3B	0.7468	0.5994	0.3878	0.061*	
C4	0.6868 (5)	0.4201 (3)	0.3659 (2)	0.0465 (7)	
H4A	0.8148	0.3511	0.3731	0.056*	
H4B	0.6334	0.3999	0.4305	0.056*	
C5	0.3905 (4)	0.4316 (3)	0.2963 (2)	0.0419 (6)	
C6	0.6877 (4)	0.2778 (3)	0.2446 (2)	0.0378 (6)	
H6	0.8002	0.3050	0.2211	0.045*	
C7	0.5756 (4)	0.2894 (3)	0.1548 (2)	0.0431 (7)	
C8	0.3869 (4)	0.3488 (3)	0.1493 (2)	0.0437 (7)	

C9	0.7520 (4)	0.1183 (3)	0.3179 (2)	0.0388 (6)	
C10	0.9389 (4)	0.0353 (3)	0.3472 (2)	0.0445 (7)	
C11	0.9927 (5)	-0.1094 (4)	0.4150 (3)	0.0569 (8)	
H11	1.1193	-0.1625	0.4324	0.068*	
C12	0.8590 (6)	-0.1737 (4)	0.4562 (3)	0.0592 (9)	
H12	0.8942	-0.2705	0.5020	0.071*	
C13	0.6714 (5)	-0.0942 (4)	0.4295 (3)	0.0574 (8)	
H13	0.5800	-0.1373	0.4575	0.069*	
C14	0.6196 (4)	0.0488 (4)	0.3613 (2)	0.0483 (7)	
H14	0.4928	0.1006	0.3438	0.058*	
C15	0.6932 (5)	0.2335 (5)	0.0802 (3)	0.0619 (9)	
C16	0.7016 (11)	0.2217 (13)	-0.0899 (6)	0.1280 (14)	0.740 (10)
H16A	0.6253	0.2918	-0.1489	0.154*	0.740 (10)
H16B	0.8211	0.2397	-0.0892	0.154*	0.740 (10)
C17	0.7244 (13)	0.0688 (10)	-0.0813 (8)	0.119 (3)	0.740 (10)
H17A	0.8124	0.0031	-0.0268	0.178*	0.740 (10)
H17B	0.7708	0.0476	-0.1417	0.178*	0.740 (10)
H17C	0.6055	0.0515	-0.0696	0.178*	0.740 (10)
C16'	0.719 (3)	0.101 (2)	-0.0367 (13)	0.124 (4)	0.260 (10)
H16C	0.8434	0.0499	-0.0034	0.148*	0.260 (10)
H16D	0.6574	0.0267	-0.0349	0.148*	0.260 (10)
C17'	0.725 (3)	0.195 (3)	-0.1337 (12)	0.119 (5)	0.260 (10)
H17D	0.6047	0.2722	-0.1532	0.179*	0.260 (10)
H17E	0.7526	0.1341	-0.1786	0.179*	0.260 (10)
H17F	0.8200	0.2412	-0.1354	0.179*	0.260 (10)
C18	0.2497 (5)	0.3678 (5)	0.0689 (3)	0.0608 (9)	
H18A	0.3170	0.3401	0.0144	0.073*	
H18B	0.1703	0.4721	0.0464	0.073*	
H18C	0.1740	0.3033	0.0946	0.073*	
Cl1	1.11696 (11)	0.11095 (11)	0.30061 (7)	0.0646 (3)	

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0587 (5)	0.0578 (5)	0.0593 (5)	-0.0108 (4)	0.0111 (4)	-0.0317 (4)
C1	0.097 (3)	0.060 (2)	0.081 (3)	-0.031 (2)	0.015 (2)	-0.019 (2)
O2	0.0477 (15)	0.158 (3)	0.0847 (19)	-0.0197 (17)	0.0138 (13)	-0.074 (2)
O3	0.0798 (19)	0.245 (4)	0.096 (2)	-0.039 (2)	0.0175 (16)	-0.120 (3)
N1	0.0346 (12)	0.0521 (15)	0.0538 (14)	-0.0099 (11)	0.0009 (10)	-0.0243 (12)
N2	0.0438 (13)	0.0361 (12)	0.0431 (12)	-0.0083 (10)	-0.0040 (10)	-0.0165 (10)
O1	0.0564 (14)	0.0666 (16)	0.0822 (17)	-0.0224 (12)	0.0144 (12)	-0.0371 (14)
C2	0.0461 (17)	0.0491 (18)	0.0602 (18)	-0.0144 (14)	-0.0057 (14)	-0.0209 (15)
C3	0.0572 (19)	0.0451 (17)	0.0556 (18)	-0.0152 (14)	-0.0027 (14)	-0.0247 (14)
C4	0.0529 (17)	0.0434 (16)	0.0436 (15)	-0.0117 (13)	-0.0077 (13)	-0.0163 (13)
C5	0.0437 (16)	0.0353 (14)	0.0448 (15)	-0.0093 (12)	0.0011 (12)	-0.0130 (12)
C6	0.0353 (14)	0.0370 (14)	0.0429 (14)	-0.0095 (11)	0.0003 (11)	-0.0164 (12)
C7	0.0427 (16)	0.0487 (17)	0.0416 (15)	-0.0152 (13)	0.0013 (12)	-0.0180 (13)
C8	0.0432 (16)	0.0476 (17)	0.0433 (15)	-0.0154 (13)	0.0017 (12)	-0.0169 (13)

C9	0.0405 (15)	0.0351 (14)	0.0444 (14)	-0.0097 (11)	0.0008 (11)	-0.0197 (12)
C10	0.0410 (15)	0.0432 (16)	0.0527 (17)	-0.0110 (12)	-0.0025 (12)	-0.0210 (13)
C11	0.0548 (19)	0.0435 (18)	0.065 (2)	-0.0028 (15)	-0.0165 (16)	-0.0176 (15)
C12	0.077 (2)	0.0376 (17)	0.0578 (19)	-0.0124 (16)	-0.0051 (17)	-0.0123 (14)
C13	0.069 (2)	0.0486 (19)	0.0612 (19)	-0.0262 (17)	0.0147 (16)	-0.0200 (16)
C14	0.0410 (16)	0.0449 (17)	0.0612 (18)	-0.0123 (13)	0.0065 (13)	-0.0216 (14)
C15	0.049 (2)	0.093 (3)	0.0526 (19)	-0.0214 (18)	0.0071 (15)	-0.0365 (19)
C16	0.0798 (19)	0.245 (4)	0.096 (2)	-0.039 (2)	0.0175 (16)	-0.120 (3)
C17	0.115 (5)	0.119 (6)	0.129 (7)	-0.028 (5)	0.036 (5)	-0.063 (5)
C16'	0.095 (6)	0.222 (8)	0.081 (6)	-0.025 (6)	0.020 (6)	-0.116 (6)
C17'	0.104 (9)	0.196 (10)	0.076 (8)	-0.034 (9)	0.028 (7)	-0.088 (8)
C18	0.0491 (19)	0.079 (2)	0.0567 (19)	-0.0157 (17)	-0.0071 (15)	-0.0273 (18)
C11	0.0387 (5)	0.0689 (6)	0.0853 (6)	-0.0185 (4)	-0.0042 (4)	-0.0190 (5)

*Geometric parameters (Å, °)*

S1—C5	1.680 (3)	C8—C18	1.500 (4)
C1—C2	1.487 (5)	C9—C10	1.388 (4)
C1—H1A	0.9600	C9—C14	1.396 (4)
C1—H1B	0.9600	C10—C11	1.387 (5)
C1—H1C	0.9600	C10—C11	1.740 (3)
O2—C15	1.189 (4)	C11—C12	1.368 (5)
O3—C15	1.318 (5)	C11—H11	0.9300
O3—C16	1.523 (7)	C12—C13	1.381 (5)
O3—C16'	1.545 (10)	C12—H12	0.9300
N1—C5	1.371 (4)	C13—C14	1.378 (5)
N1—C8	1.382 (4)	C13—H13	0.9300
N1—H1	0.8600	C14—H14	0.9300
N2—C5	1.335 (4)	C16—C17	1.408 (11)
N2—C4	1.472 (4)	C16—H16A	0.9700
N2—C6	1.483 (3)	C16—H16B	0.9700
O1—C2	1.215 (4)	C17—H17A	0.9600
C2—C3	1.497 (5)	C17—H17B	0.9600
C3—C4	1.519 (4)	C17—H17C	0.9600
C3—H3A	0.9700	C16'—C17'	1.407 (13)
C3—H3B	0.9700	C16'—H16C	0.9700
C4—H4A	0.9700	C16'—H16D	0.9700
C4—H4B	0.9700	C17'—H17D	0.9600
C6—C7	1.512 (4)	C17'—H17E	0.9600
C6—C9	1.521 (4)	C17'—H17F	0.9600
C6—H6	0.9800	C18—H18A	0.9600
C7—C8	1.346 (4)	C18—H18B	0.9600
C7—C15	1.461 (4)	C18—H18C	0.9600
C2—C1—H1A	109.5	C10—C9—C14	116.3 (3)
C2—C1—H1B	109.5	C10—C9—C6	123.7 (3)
H1A—C1—H1B	109.5	C14—C9—C6	120.0 (2)
C2—C1—H1C	109.5	C11—C10—C9	122.3 (3)

H1A—C1—H1C	109.5	C11—C10—C11	117.0 (2)
H1B—C1—H1C	109.5	C9—C10—C11	120.8 (2)
C15—O3—C16	117.8 (4)	C12—C11—C10	119.8 (3)
C15—O3—C16'	109.7 (9)	C12—C11—H11	120.1
C16—O3—C16'	44.4 (9)	C10—C11—H11	120.1
C5—N1—C8	125.2 (2)	C11—C12—C13	119.6 (3)
C5—N1—H1	117.4	C11—C12—H12	120.2
C8—N1—H1	117.4	C13—C12—H12	120.2
C5—N2—C4	120.3 (2)	C14—C13—C12	120.1 (3)
C5—N2—C6	122.7 (2)	C14—C13—H13	120.0
C4—N2—C6	115.6 (2)	C12—C13—H13	120.0
O1—C2—C1	121.0 (3)	C13—C14—C9	121.9 (3)
O1—C2—C3	121.7 (3)	C13—C14—H14	119.0
C1—C2—C3	117.2 (3)	C9—C14—H14	119.0
C2—C3—C4	115.6 (3)	O2—C15—O3	121.8 (3)
C2—C3—H3A	108.4	O2—C15—C7	123.6 (3)
C4—C3—H3A	108.4	O3—C15—C7	114.6 (3)
C2—C3—H3B	108.4	C17—C16—O3	97.9 (7)
C4—C3—H3B	108.4	C17—C16—H16A	112.2
H3A—C3—H3B	107.4	O3—C16—H16A	112.2
N2—C4—C3	113.4 (2)	C17—C16—H16B	112.2
N2—C4—H4A	108.9	O3—C16—H16B	112.2
C3—C4—H4A	108.9	H16A—C16—H16B	109.8
N2—C4—H4B	108.9	C17'—C16'—O3	96.5 (10)
C3—C4—H4B	108.9	C17'—C16'—H16C	112.5
H4A—C4—H4B	107.7	O3—C16'—H16C	112.5
N2—C5—N1	116.0 (2)	C17'—C16'—H16D	112.5
N2—C5—S1	125.3 (2)	O3—C16'—H16D	112.5
N1—C5—S1	118.7 (2)	H16C—C16'—H16D	110.0
N2—C6—C7	110.4 (2)	C16'—C17'—H17D	109.5
N2—C6—C9	109.7 (2)	C16'—C17'—H17E	109.5
C7—C6—C9	113.2 (2)	H17D—C17'—H17E	109.5
N2—C6—H6	107.8	C16'—C17'—H17F	109.5
C7—C6—H6	107.8	H17D—C17'—H17F	109.5
C9—C6—H6	107.8	H17E—C17'—H17F	109.5
C8—C7—C15	126.8 (3)	C8—C18—H18A	109.5
C8—C7—C6	120.1 (3)	C8—C18—H18B	109.5
C15—C7—C6	113.1 (2)	H18A—C18—H18B	109.5
C7—C8—N1	118.3 (3)	C8—C18—H18C	109.5
C7—C8—C18	128.8 (3)	H18A—C18—H18C	109.5
N1—C8—C18	112.9 (3)	H18B—C18—H18C	109.5
O1—C2—C3—C4	-5.4 (4)	C7—C6—C9—C10	122.2 (3)
C1—C2—C3—C4	176.6 (3)	N2—C6—C9—C14	64.6 (3)
C5—N2—C4—C3	-79.4 (3)	C7—C6—C9—C14	-59.2 (3)
C6—N2—C4—C3	114.0 (3)	C14—C9—C10—C11	0.7 (4)
C2—C3—C4—N2	-67.4 (4)	C6—C9—C10—C11	179.3 (3)
C4—N2—C5—N1	177.4 (2)	C14—C9—C10—C11	-178.5 (2)

C6—N2—C5—N1	-17.0 (4)	C6—C9—C10—C11	0.1 (4)
C4—N2—C5—S1	-0.3 (4)	C9—C10—C11—C12	-0.8 (5)
C6—N2—C5—S1	165.4 (2)	C11—C10—C11—C12	178.4 (3)
C8—N1—C5—N2	-7.7 (4)	C10—C11—C12—C13	0.3 (5)
C8—N1—C5—S1	170.1 (2)	C11—C12—C13—C14	0.3 (5)
C5—N2—C6—C7	30.6 (4)	C12—C13—C14—C9	-0.4 (5)
C4—N2—C6—C7	-163.1 (2)	C10—C9—C14—C13	-0.1 (4)
C5—N2—C6—C9	-94.8 (3)	C6—C9—C14—C13	-178.7 (3)
C4—N2—C6—C9	71.4 (3)	C16—O3—C15—O2	-19.1 (9)
N2—C6—C7—C8	-22.2 (4)	C16'—O3—C15—O2	28.9 (11)
C9—C6—C7—C8	101.3 (3)	C16—O3—C15—C7	161.1 (6)
N2—C6—C7—C15	156.9 (3)	C16'—O3—C15—C7	-150.9 (10)
C9—C6—C7—C15	-79.7 (3)	C8—C7—C15—O2	173.4 (4)
C15—C7—C8—N1	-176.9 (3)	C6—C7—C15—O2	-5.5 (6)
C6—C7—C8—N1	2.0 (4)	C8—C7—C15—O3	-6.8 (6)
C15—C7—C8—C18	2.5 (6)	C6—C7—C15—O3	174.3 (4)
C6—C7—C8—C18	-178.6 (3)	C15—O3—C16—C17	107.2 (7)
C5—N1—C8—C7	15.2 (4)	C16'—O3—C16—C17	16.9 (14)
C5—N1—C8—C18	-164.4 (3)	C15—O3—C16'—C17'	-120.7 (15)
N2—C6—C9—C10	-114.0 (3)	C16—O3—C16'—C17'	-10.6 (11)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O1 <sup>i</sup>	0.86	2.16	2.984 (3)	160

Symmetry code: (i)  $x-1, y, z$ .