

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Hydroxy-*N'*-[1-(3-methylpyrazin-2-yl)-ethylidene]benzohydrazide

Tai Xi-Shi and Feng Yi-Min*

Department of Chemistry and Chemical Engineering, Weifang University, Weifang 261061, People's Republic of China


Correspondence e-mail: taixishi@lzu.edu.cn

Received 10 March 2008; accepted 10 March 2008

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.045; wR factor = 0.156; data-to-parameter ratio = 12.7.

The molecule of the title compound, $\text{C}_{14}\text{H}_{14}\text{N}_4\text{O}_2$, is slightly twisted, with a dihedral angle of 10.06 (14) $^\circ$ between the aromatic rings. An intramolecular $\text{N}-\text{H}\cdots\text{O}$ and an intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond help to establish the crystal structure.

Related literature

 For related literature, see: Tai *et al.* (2003).


Experimental

Crystal data

$\text{C}_{14}\text{H}_{14}\text{N}_4\text{O}_2$
 $M_r = 270.29$
 Orthorhombic, *Pbcn*
 $a = 12.6326$ (14) Å

$b = 9.3346$ (10) Å
 $c = 22.119$ (3) Å
 $V = 2608.3$ (5) Å³
 $Z = 8$

Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹

$T = 298$ (2) K
 $0.49 \times 0.43 \times 0.22$ mm

Data collection

Bruker SMART CCD diffractometer
 Absorption correction: multi-scan (*SADABS*; Bruker, 2000)
 $T_{\min} = 0.954$, $T_{\max} = 0.979$

12690 measured reflections
 2303 independent reflections
 1380 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.052$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.156$
 $S = 1.07$
 2303 reflections

181 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.20$ e Å⁻³
 $\Delta\rho_{\min} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots O2	0.86	1.95	2.624 (3)	134
O2—H2 \cdots O1 ¹	0.82	1.82	2.631 (3)	168

 Symmetry code: (i) $x + \frac{1}{2}, -y + \frac{1}{2}, -z + 1$.

Data collection: *SMART* (Bruker, 2000); cell refinement: *SAINTE* (Bruker, 2000); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the National Natural Science Foundation of China (20671073), the National Natural Science Foundation of Shandong, the Science and Technology Foundation of Weifang and Weifang University for research grants.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2708).

References

- Bruker (2000). *SMART*, *SAINTE* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Tai, X.-S., Yin, X.-H., Tan, M.-Y. & Li, Y.-Z. (2003). *Acta Cryst.* **E59**, o681–o682.

supporting information

Acta Cryst. (2008). E64, o707 [doi:10.1107/S1600536808006697]

2-Hydroxy-*N'*-[1-(3-methylpyrazin-2-yl)ethylidene]benzohydrazide

Tai Xi-Shi and Feng Yi-Min

S1. Comment

As part of our ongoing studies of the chemistry of aroylhydrazone ligands (Tai *et al.*, 2003), we now report the synthesis and structure of the title compound, (I), (Fig. 1).

The molecule is slightly twisted, with a dihedral angle of 10.06 (14)^o between the aromatic rings. An intramolecular N—H \cdots O and an intermolecular O—H \cdots O hydrogen bond help to establish the structure (Table 1).

S2. Experimental

10 mmol of 2-acetyl-3-methylpyrazine (10 mmol) was added to a solution of salicyloyl hydrazine (10 mmol) in 10 ml of ethanol. The mixture was continuously stirred for 3 h at refluxing temperature, evaporating some ethanol. Upon cooling, the solid product was collected by filtration and dried *in vacuo* (yield 62%). Colourless blocks of (I) were obtained by evaporation from a methanol solution after one week.

S3. Refinement

The H atoms were placed geometrically (C—H = 0.93–0.96 Å, O—H = 0.82 Å, N—H = 0.86 Å) and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{carrier})$ or $1.5U_{\text{eq}}(\text{methyl C})$.




Figure 1

The molecular structure of (I) showing 50% displacement ellipsoids. The hydrogen bond is indicated by a double-dashed line.

2-Hydroxy-*N'*-[1-(3-methylpyrazin-2-yl)ethylidene]benzohydrazide

Crystal data

$C_{14}H_{14}N_4O_2$

$M_r = 270.29$

Orthorhombic, *Pbcn*

Hall symbol: -P 2n 2ab

$a = 12.6326$ (14) Å

$b = 9.3346$ (10) Å

$c = 22.119$ (3) Å

$V = 2608.3$ (5) Å³

$Z = 8$

$F(000) = 1136$

$D_x = 1.377$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2332 reflections

$\theta = 2.5$ – 23.7°

$\mu = 0.10$ mm⁻¹

$T = 298$ K

Block, colourless

$0.49 \times 0.43 \times 0.22$ mm

Data collection

Bruker SMART CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2000)

$T_{\min} = 0.954$, $T_{\max} = 0.979$

12690 measured reflections

2303 independent reflections

1380 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.052$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 1.8^\circ$

$h = -15 \rightarrow 13$

$k = -11 \rightarrow 10$

$l = -26 \rightarrow 24$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.156$
 $S = 1.07$
 2303 reflections
 181 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0649P)^2 + 1.2514P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.20 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.79886 (17)	0.1654 (2)	0.42259 (10)	0.0408 (6)
H1	0.8658	0.1773	0.4278	0.049*
N2	0.76327 (17)	0.0717 (2)	0.37945 (10)	0.0396 (6)
N3	0.8738 (2)	-0.1605 (3)	0.27123 (12)	0.0590 (8)
N4	0.6645 (2)	-0.2165 (3)	0.24385 (12)	0.0586 (8)
O1	0.63328 (14)	0.2289 (2)	0.44838 (9)	0.0553 (6)
O2	0.95222 (14)	0.2544 (2)	0.49322 (9)	0.0574 (6)
H2	1.0112	0.2679	0.5076	0.086*
C1	0.7298 (2)	0.2386 (3)	0.45667 (12)	0.0379 (7)
C2	0.77414 (19)	0.3353 (3)	0.50398 (11)	0.0356 (6)
C3	0.8809 (2)	0.3424 (3)	0.52082 (12)	0.0401 (7)
C4	0.9125 (2)	0.4359 (3)	0.56555 (13)	0.0478 (8)
H4	0.9834	0.4394	0.5767	0.057*
C5	0.8411 (2)	0.5234 (3)	0.59369 (13)	0.0498 (8)
H5	0.8638	0.5865	0.6235	0.060*
C6	0.7359 (2)	0.5186 (3)	0.57807 (13)	0.0465 (8)
H6	0.6872	0.5779	0.5973	0.056*
C7	0.7034 (2)	0.4253 (3)	0.53383 (12)	0.0425 (7)
H7	0.6321	0.4221	0.5235	0.051*
C8	0.9508 (2)	0.0190 (4)	0.35800 (17)	0.0733 (11)
H8A	0.9701	0.1181	0.3542	0.110*
H8B	0.9873	-0.0360	0.3278	0.110*
H8C	0.9701	-0.0150	0.3975	0.110*
C9	0.8340 (2)	0.0033 (3)	0.34940 (13)	0.0426 (7)
C10	0.7959 (2)	-0.0970 (3)	0.30201 (12)	0.0409 (7)

C11	0.6890 (2)	-0.1265 (3)	0.28871 (12)	0.0428 (7)
C12	0.5955 (2)	-0.0662 (4)	0.32156 (14)	0.0568 (9)
H12A	0.5317	-0.1066	0.3053	0.085*
H12B	0.5941	0.0360	0.3167	0.085*
H12C	0.6007	-0.0892	0.3638	0.085*
C13	0.7439 (3)	-0.2763 (4)	0.21323 (15)	0.0627 (9)
H13	0.7283	-0.3383	0.1815	0.075*
C14	0.8469 (3)	-0.2494 (4)	0.22696 (15)	0.0639 (10)
H14	0.8998	-0.2945	0.2048	0.077*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0318 (12)	0.0478 (15)	0.0428 (14)	-0.0032 (11)	-0.0015 (10)	-0.0074 (12)
N2	0.0381 (13)	0.0423 (14)	0.0382 (13)	-0.0022 (11)	0.0012 (10)	-0.0040 (11)
N3	0.0574 (17)	0.0631 (18)	0.0565 (17)	0.0068 (14)	0.0117 (13)	-0.0102 (15)
N4	0.0638 (18)	0.0608 (18)	0.0510 (16)	0.0056 (15)	-0.0092 (13)	-0.0135 (14)
O1	0.0312 (11)	0.0761 (16)	0.0585 (13)	-0.0021 (10)	-0.0003 (9)	-0.0172 (12)
O2	0.0328 (11)	0.0712 (15)	0.0684 (14)	0.0027 (10)	-0.0054 (9)	-0.0268 (12)
C1	0.0308 (15)	0.0439 (18)	0.0389 (15)	0.0007 (13)	0.0035 (12)	0.0038 (13)
C2	0.0336 (15)	0.0361 (16)	0.0372 (15)	-0.0018 (12)	0.0008 (11)	0.0032 (13)
C3	0.0343 (15)	0.0423 (17)	0.0437 (16)	-0.0006 (13)	0.0029 (12)	-0.0033 (14)
C4	0.0374 (16)	0.0539 (19)	0.0522 (18)	-0.0010 (15)	-0.0050 (13)	-0.0084 (16)
C5	0.0552 (19)	0.0469 (19)	0.0473 (17)	-0.0040 (15)	-0.0003 (15)	-0.0080 (15)
C6	0.0506 (18)	0.0431 (18)	0.0459 (17)	0.0047 (14)	0.0040 (14)	-0.0008 (14)
C7	0.0385 (15)	0.0449 (18)	0.0440 (16)	0.0039 (14)	0.0017 (13)	0.0027 (15)
C8	0.0386 (18)	0.087 (3)	0.094 (3)	0.0031 (18)	0.0004 (17)	-0.033 (2)
C9	0.0359 (15)	0.0455 (18)	0.0464 (17)	0.0021 (14)	0.0031 (13)	0.0001 (14)
C10	0.0453 (17)	0.0388 (17)	0.0386 (15)	0.0063 (13)	0.0041 (13)	0.0013 (13)
C11	0.0474 (17)	0.0437 (18)	0.0374 (16)	0.0061 (14)	-0.0021 (13)	0.0019 (14)
C12	0.0408 (17)	0.072 (2)	0.0576 (19)	0.0035 (16)	-0.0044 (14)	-0.0143 (17)
C13	0.078 (2)	0.061 (2)	0.0490 (19)	0.011 (2)	-0.0054 (18)	-0.0167 (17)
C14	0.068 (2)	0.069 (2)	0.055 (2)	0.013 (2)	0.0114 (17)	-0.0118 (19)

Geometric parameters (Å, °)

N1—C1	1.340 (3)	C5—H5	0.9300
N1—N2	1.370 (3)	C6—C7	1.373 (4)
N1—H1	0.8600	C6—H6	0.9300
N2—C9	1.284 (3)	C7—H7	0.9300
N3—C14	1.328 (4)	C8—C9	1.495 (4)
N3—C10	1.335 (3)	C8—H8A	0.9600
N4—C13	1.332 (4)	C8—H8B	0.9600
N4—C11	1.336 (4)	C8—H8C	0.9600
O1—C1	1.236 (3)	C9—C10	1.485 (4)
O2—C3	1.364 (3)	C10—C11	1.409 (4)
O2—H2	0.8200	C11—C12	1.497 (4)
C1—C2	1.492 (4)	C12—H12A	0.9600

C2—C7	1.393 (4)	C12—H12B	0.9600
C2—C3	1.400 (3)	C12—H12C	0.9600
C3—C4	1.378 (4)	C13—C14	1.359 (5)
C4—C5	1.367 (4)	C13—H13	0.9300
C4—H4	0.9300	C14—H14	0.9300
C5—C6	1.374 (4)		
C1—N1—N2	120.2 (2)	C9—C8—H8A	109.5
C1—N1—H1	119.9	C9—C8—H8B	109.5
N2—N1—H1	119.9	H8A—C8—H8B	109.5
C9—N2—N1	116.7 (2)	C9—C8—H8C	109.5
C14—N3—C10	117.7 (3)	H8A—C8—H8C	109.5
C13—N4—C11	117.8 (3)	H8B—C8—H8C	109.5
C3—O2—H2	109.5	N2—C9—C10	117.0 (2)
O1—C1—N1	121.5 (3)	N2—C9—C8	124.9 (3)
O1—C1—C2	121.2 (2)	C10—C9—C8	118.1 (2)
N1—C1—C2	117.3 (2)	N3—C10—C11	120.8 (3)
C7—C2—C3	117.6 (2)	N3—C10—C9	113.7 (3)
C7—C2—C1	117.2 (2)	C11—C10—C9	125.5 (2)
C3—C2—C1	125.2 (2)	N4—C11—C10	120.0 (3)
O2—C3—C4	120.7 (2)	N4—C11—C12	114.4 (3)
O2—C3—C2	119.3 (2)	C10—C11—C12	125.5 (3)
C4—C3—C2	120.0 (3)	C11—C12—H12A	109.5
C5—C4—C3	120.9 (3)	C11—C12—H12B	109.5
C5—C4—H4	119.6	H12A—C12—H12B	109.5
C3—C4—H4	119.6	C11—C12—H12C	109.5
C4—C5—C6	120.3 (3)	H12A—C12—H12C	109.5
C4—C5—H5	119.9	H12B—C12—H12C	109.5
C6—C5—H5	119.9	N4—C13—C14	121.9 (3)
C7—C6—C5	119.3 (3)	N4—C13—H13	119.0
C7—C6—H6	120.3	C14—C13—H13	119.0
C5—C6—H6	120.3	N3—C14—C13	121.7 (3)
C6—C7—C2	121.9 (3)	N3—C14—H14	119.2
C6—C7—H7	119.1	C13—C14—H14	119.2
C2—C7—H7	119.1		
C1—N1—N2—C9	-178.6 (3)	N1—N2—C9—C10	-179.2 (2)
N2—N1—C1—O1	-3.4 (4)	N1—N2—C9—C8	-0.2 (4)
N2—N1—C1—C2	178.2 (2)	C14—N3—C10—C11	1.6 (4)
O1—C1—C2—C7	-6.7 (4)	C14—N3—C10—C9	-178.4 (3)
N1—C1—C2—C7	171.7 (2)	N2—C9—C10—N3	177.8 (3)
O1—C1—C2—C3	173.1 (3)	C8—C9—C10—N3	-1.3 (4)
N1—C1—C2—C3	-8.6 (4)	N2—C9—C10—C11	-2.2 (4)
C7—C2—C3—O2	179.0 (2)	C8—C9—C10—C11	178.7 (3)
C1—C2—C3—O2	-0.7 (4)	C13—N4—C11—C10	0.4 (4)
C7—C2—C3—C4	0.1 (4)	C13—N4—C11—C12	-178.9 (3)
C1—C2—C3—C4	-179.6 (3)	N3—C10—C11—N4	-1.6 (4)
O2—C3—C4—C5	-179.4 (3)	C9—C10—C11—N4	178.4 (3)

C2—C3—C4—C5	-0.6 (4)	N3—C10—C11—C12	177.6 (3)
C3—C4—C5—C6	0.6 (5)	C9—C10—C11—C12	-2.4 (5)
C4—C5—C6—C7	-0.2 (4)	C11—N4—C13—C14	0.8 (5)
C5—C6—C7—C2	-0.2 (4)	C10—N3—C14—C13	-0.4 (5)
C3—C2—C7—C6	0.3 (4)	N4—C13—C14—N3	-0.9 (6)
C1—C2—C7—C6	-180.0 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O2	0.86	1.95	2.624 (3)	134
O2—H2...O1 ⁱ	0.82	1.82	2.631 (3)	168

Symmetry code: (i) $x+1/2, -y+1/2, -z+1$.