

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-(2,4-Dichlorophenyl)-3-[5-(4-methoxyphenyl)-1,3,4-thiadiazol-2-yl]-1,3-thiazolidin-4-one

Rong Wan,* Li-He Yin, Feng Han, Bin Wang and Jin-Tang Wang

 Department of Applied Chemistry, College of Science, Nanjing University of Technology, No. 5 Xinmofan Road, Nanjing 210009, People's Republic of China
 Correspondence e-mail: rwan@njut.edu.cn


Received 27 March 2008; accepted 29 March 2008

 Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.008$ Å; R factor = 0.068; wR factor = 0.212; data-to-parameter ratio = 13.6.

In the molecule of the title compound, $\text{C}_{18}\text{H}_{13}\text{Cl}_2\text{N}_3\text{O}_2\text{S}_2$, the thiazolidinone ring has an envelope conformation with the S atom displaced by 0.394 (3) Å from the plane of the other ring atoms. The thiadiazole ring is oriented at a dihedral angle of 7.40 (4)° with respect to the 4-methoxyphenyl ring. Intramolecular C—H···S, C—H···N and C—H···Cl hydrogen bonds result in the formation of two planar and two non-planar five-membered rings. The planar five-membered rings are oriented at a dihedral angle of 6.23 (3)°. The 2,4-dichlorophenyl ring is oriented at dihedral angles of 84.21 (4) and 83.55 (3)° with respect to the thiadiazole and 4-methoxyphenyl rings, respectively. In the crystal structure, intermolecular C—H···O hydrogen bonds link the molecules into centrosymmetric dimers.

Related literature

For general background, see: Chen *et al.* (2000); Kidwai *et al.* (2000); Vicentini *et al.* (1998); Arun *et al.* (1999); Wasfy *et al.* (1996).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{13}\text{Cl}_2\text{N}_3\text{O}_2\text{S}_2$
 $M_r = 438.33$
 Triclinic, $P\bar{1}$
 $a = 7.1310$ (14) Å
 $b = 8.1540$ (16) Å
 $c = 16.671$ (3) Å
 $\alpha = 93.19$ (3)°
 $\beta = 96.43$ (3)°
 $\gamma = 105.89$ (3)°
 $V = 922.7$ (3) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.60$ mm⁻¹
 $T = 298$ (2) K
 $0.30 \times 0.10 \times 0.10$ mm

Data collection

Enraf–Nonius CAD-4 diffractometer
 Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.841$, $T_{\max} = 0.943$
 3606 measured reflections
 3315 independent reflections
 2228 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.084$
 3 standard reflections every 200 reflections
 intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.067$
 $wR(F^2) = 0.211$
 $S = 1.02$
 3315 reflections
 244 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.47$ e Å⁻³
 $\Delta\rho_{\min} = -0.61$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C4—H4A···S1	0.93	2.79	3.180 (7)	106
C6—H6A···N1	0.93	2.55	2.856 (8)	100
C12—H12A···Cl2	0.98	2.63	3.063 (5)	107
C14—H14A···N3	0.93	2.54	2.863 (8)	101
C14—H14A···O1 ¹	0.93	2.41	3.219 (7)	146

 Symmetry code: (i) $-x + 2, -y + 1, -z$.

Data collection: *CAD-4 Software* (Enraf–Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Professor Hua-Qin Wang of Nanjing University for carrying out the X-ray crystallographic analysis.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2442).

References

- Arun, K. P., Nag, V. L. & Panda, C. S. (1999). *Indian J. Chem.* **38B**, 998–1001.
 Chen, H. S., Li, Z. M. & Han, Y. F. (2000). *J. Agric. Food Chem.* **48**, 5312–5315.
 Enraf–Nonius (1989). *CAD-4 Software*. Enraf–Nonius, Delft, The Netherlands.
 Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
 Kidwai, M., Negi, N. & Misra, P. (2000). *J. Indian Chem. Soc.* **77**, 46–48.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Vicentini, C. B., Manfrini, M., Veronese, A. C. & Guarnieri, M. (1998). *J. Heterocycl. Chem.* **35**, 29–36.
 Wasfy, A. A., Nassar, S. A. & Eissa, A. M. (1996). *Indian J. Chem.* **35B**, 1218–1220.

supporting information

Acta Cryst. (2008). E64, o795 [doi:10.1107/S1600536808008465]

2-(2,4-Dichlorophenyl)-3-[5-(4-methoxyphenyl)-1,3,4-thiadiazol-2-yl]-1,3-thiazolidin-4-one

Rong Wan, Li-He Yin, Feng Han, Bin Wang and Jin-Tang Wang

S1. Comment

1,3,4-Thiadiazole derivatives containing the thiazolidinone unit are of great interest because of their chemical and pharmaceutical properties. Some derivatives have fungicidal activities and exhibit certain herbicidal activities (Chen *et al.*, 2000; Kidwai *et al.*, 2000; Vicentini *et al.*, 1998). On the other hand, some of them show insecticidal activities (Arun *et al.*, 1999; Wasfy *et al.*, 1996). We report herein the crystal structure of the title compound, (I).

In the molecule of (I), (Fig. 1), rings A (C2-C7), B (S1/N1/N2/C8/C9) and D (C13-C18) are, of course, planar. The dihedral angles between them are A/B = 7.40 (4)°, A/D = 83.55 (3)° and B/D = 84.21 (4)°. So, rings A and B are nearly coplanar. Ring C (S2/N3/C10-C12) has envelope conformation with atom S2 displaced by 0.394 (3) Å from the plane of the other ring atoms. The intramolecular C-H...S, C-H...N and C-H...Cl hydrogen bonds (Table 1) result in the formation of two planar and two non-planar five-membered rings E (S1/C4/H4A/C5/C8), F (N1/C5/C6/H6A/C8) and G (N3/C12-C14/H14A), H (C12/C12/H12A/C13/C18). The dihedral angle between the planar rings E and F is E/F = 6.23 (3)°, and they are oriented with respect to the adjacent rings at dihedral angles of A/E = 3.26 (4)°, A/F = 4.55 (3)°, B/E = 5.03 (4)° and B/F = 7.27 (4)°. So, they are also nearly coplanar.

In the crystal structure, intermolecular C-H...O hydrogen bonds (Table 1) link the molecules into centrosymmetric dimers (Fig. 2), in which they may be effective in the stabilization of the structure.

S2. Experimental

N-(2,4-dichlorobenzylidene)-5-(4-methoxyphenyl)-1,3,4-thiadiazol-2-amine (5 mmol) and mercapto-acetic acid (5 mmol) were added in toluene (50 ml). The water was removed by distillation for 5 h. The reaction mixture was left to cool to room temperature, filtered, and the filter cake was crystallized from acetone to give pure compound (I) (m.p. 507-509 K). Crystals of (I) suitable for X-ray analysis were obtained by slow evaporation of an acetone solution.

S3. Refinement

H atoms were positioned geometrically, with C-H = 0.93, 0.98, 0.97 and 0.96 Å for aromatic, methine, methylene and methyl H, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl H and $x = 1.2$ for all other H atoms.




Figure 1

The molecular structure of the title molecule, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. Hydrogen bonds are shown as dashed lines




Figure 2

A partial packing diagram of (I). Hydrogen bonds are shown as dashed lines.

2-(2,4-Dichlorophenyl)-3-[5-(4-methoxyphenyl)-1,3,4-thiadiazol-2-yl]-1,3-thiazolidin-4-one

Crystal data

$C_{18}H_{13}Cl_2N_3O_2S_2$

$M_r = 438.33$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 7.1310(14)\ \text{\AA}$

$b = 8.1540(16)\ \text{\AA}$

$c = 16.671(3)\ \text{\AA}$

$\alpha = 93.19(3)^\circ$

$\beta = 96.43(3)^\circ$

$\gamma = 105.89(3)^\circ$

$V = 922.7(3)\ \text{\AA}^3$

$Z = 2$

$F(000) = 448$
 $D_x = 1.578 \text{ Mg m}^{-3}$
 Melting point = 507–509 K
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 25 reflections

$\theta = 9\text{--}12^\circ$
 $\mu = 0.60 \text{ mm}^{-1}$
 $T = 298 \text{ K}$
 Block, colorless
 $0.30 \times 0.10 \times 0.10 \text{ mm}$

Data collection

Enraf–Nonius CAD-4
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 $\omega/2\theta$ scans
 Absorption correction: ψ scan
 (North *et al.*, 1968)
 $T_{\min} = 0.841$, $T_{\max} = 0.943$
 3606 measured reflections

3315 independent reflections
 2228 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.084$
 $\theta_{\max} = 25.2^\circ$, $\theta_{\min} = 1.2^\circ$
 $h = 0 \rightarrow 8$
 $k = -9 \rightarrow 9$
 $l = -19 \rightarrow 19$
 3 standard reflections every 200 reflections
 intensity decay: none

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.067$
 $wR(F^2) = 0.211$
 $S = 1.02$
 3315 reflections
 244 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 2P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.47 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.61 \text{ e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	1.3375 (3)	0.6645 (2)	0.50289 (13)	0.0801 (6)
Cl2	1.1228 (2)	1.20318 (19)	0.40205 (10)	0.0606 (5)
S1	0.78625 (18)	0.7174 (2)	0.06058 (9)	0.0492 (4)
S2	0.5812 (2)	0.9562 (2)	0.33166 (11)	0.0659 (5)
O1	1.3277 (6)	0.6538 (6)	-0.2389 (2)	0.0616 (11)
O2	0.4564 (5)	0.6490 (6)	0.1380 (3)	0.0653 (12)
N1	1.1375 (6)	0.9039 (7)	0.1050 (3)	0.0535 (13)
N2	1.0345 (6)	0.9262 (7)	0.1690 (3)	0.0557 (13)
N3	0.7229 (6)	0.8433 (6)	0.2094 (3)	0.0488 (12)
C1	1.5346 (9)	0.7003 (9)	-0.2423 (4)	0.0677 (18)

H1B	1.5586	0.6672	-0.2955	0.102*
H1C	1.5962	0.6431	-0.2030	0.102*
H1D	1.5880	0.8220	-0.2306	0.102*
C2	1.2658 (8)	0.6915 (7)	-0.1674 (3)	0.0481 (13)
C3	1.0653 (8)	0.6308 (9)	-0.1639 (4)	0.0606 (17)
H3A	0.9826	0.5675	-0.2088	0.073*
C4	0.9884 (8)	0.6632 (8)	-0.0956 (4)	0.0556 (16)
H4A	0.8538	0.6203	-0.0944	0.067*
C5	1.1052 (7)	0.7578 (7)	-0.0282 (3)	0.0444 (13)
C6	1.3061 (8)	0.8127 (8)	-0.0323 (4)	0.0561 (16)
H6A	1.3900	0.8721	0.0131	0.067*
C7	1.3842 (8)	0.7822 (8)	-0.1009 (4)	0.0556 (16)
H7A	1.5189	0.8237	-0.1021	0.067*
C8	1.0288 (7)	0.8015 (7)	0.0450 (3)	0.0451 (13)
C9	0.8518 (7)	0.8390 (7)	0.1524 (3)	0.0457 (13)
C10	0.5300 (7)	0.7452 (8)	0.1973 (4)	0.0491 (14)
C11	0.4280 (9)	0.7736 (9)	0.2684 (4)	0.0670 (18)
H11A	0.4038	0.6734	0.2987	0.080*
H11B	0.3025	0.7928	0.2499	0.080*
C12	0.7992 (7)	0.9486 (7)	0.2846 (3)	0.0463 (13)
H12A	0.8693	1.0643	0.2731	0.056*
C13	0.9363 (7)	0.8774 (7)	0.3396 (3)	0.0414 (12)
C14	0.9118 (8)	0.7054 (7)	0.3390 (4)	0.0501 (14)
H14A	0.8099	0.6322	0.3033	0.060*
C15	1.0296 (8)	0.6354 (7)	0.3883 (4)	0.0518 (14)
H15A	1.0095	0.5175	0.3865	0.062*
C16	1.1798 (8)	0.7468 (8)	0.4411 (3)	0.0473 (13)
C17	1.2064 (7)	0.9194 (7)	0.4448 (3)	0.0464 (13)
H17A	1.3064	0.9927	0.4813	0.056*
C18	1.0853 (7)	0.9832 (7)	0.3946 (3)	0.0425 (12)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0617 (10)	0.0839 (13)	0.0961 (14)	0.0270 (9)	-0.0084 (9)	0.0253 (10)
C12	0.0494 (8)	0.0463 (8)	0.0828 (11)	0.0091 (6)	0.0096 (7)	-0.0025 (7)
S1	0.0242 (6)	0.0653 (10)	0.0511 (9)	0.0018 (6)	0.0043 (6)	0.0040 (7)
S2	0.0417 (8)	0.0898 (13)	0.0728 (11)	0.0289 (8)	0.0138 (8)	-0.0028 (9)
O1	0.044 (2)	0.072 (3)	0.058 (3)	-0.001 (2)	0.0138 (19)	-0.011 (2)
O2	0.028 (2)	0.088 (3)	0.068 (3)	-0.001 (2)	0.0030 (19)	-0.002 (3)
N1	0.028 (2)	0.076 (3)	0.051 (3)	0.004 (2)	0.011 (2)	-0.005 (2)
N2	0.030 (2)	0.075 (3)	0.054 (3)	0.002 (2)	0.008 (2)	-0.003 (3)
N3	0.026 (2)	0.072 (3)	0.049 (3)	0.012 (2)	0.012 (2)	0.007 (2)
C1	0.045 (3)	0.090 (5)	0.071 (4)	0.018 (3)	0.026 (3)	0.002 (4)
C2	0.038 (3)	0.049 (3)	0.053 (3)	0.004 (2)	0.009 (3)	0.006 (3)
C3	0.033 (3)	0.084 (5)	0.053 (4)	0.004 (3)	0.000 (3)	-0.010 (3)
C4	0.025 (3)	0.076 (4)	0.057 (4)	0.003 (3)	-0.001 (2)	-0.002 (3)
C5	0.029 (3)	0.048 (3)	0.052 (3)	0.003 (2)	0.004 (2)	0.005 (3)

C6	0.027 (3)	0.076 (4)	0.050 (3)	-0.005 (3)	0.001 (2)	-0.009 (3)
C7	0.029 (3)	0.075 (4)	0.055 (4)	0.001 (3)	0.009 (3)	0.001 (3)
C8	0.025 (2)	0.052 (3)	0.056 (3)	0.004 (2)	0.007 (2)	0.011 (3)
C9	0.030 (3)	0.056 (3)	0.052 (3)	0.012 (2)	0.007 (2)	0.011 (3)
C10	0.025 (3)	0.061 (4)	0.062 (4)	0.010 (2)	0.008 (3)	0.012 (3)
C11	0.039 (3)	0.092 (5)	0.072 (4)	0.018 (3)	0.020 (3)	0.008 (4)
C12	0.034 (3)	0.054 (3)	0.054 (3)	0.015 (2)	0.010 (2)	0.006 (3)
C13	0.028 (2)	0.048 (3)	0.049 (3)	0.011 (2)	0.011 (2)	0.000 (2)
C14	0.037 (3)	0.052 (3)	0.058 (4)	0.009 (3)	0.005 (3)	-0.003 (3)
C15	0.041 (3)	0.043 (3)	0.071 (4)	0.011 (3)	0.014 (3)	0.004 (3)
C16	0.035 (3)	0.058 (4)	0.053 (3)	0.016 (3)	0.010 (2)	0.011 (3)
C17	0.028 (3)	0.057 (4)	0.050 (3)	0.006 (2)	0.006 (2)	0.000 (3)
C18	0.034 (3)	0.042 (3)	0.054 (3)	0.010 (2)	0.019 (2)	0.003 (2)

Geometric parameters (Å, °)

S1—C9	1.721 (6)	C4—C5	1.375 (8)
S1—C8	1.732 (5)	C4—H4A	0.9300
C11—C16	1.734 (6)	C5—C6	1.390 (7)
O1—C2	1.366 (7)	C5—C8	1.456 (8)
O1—C1	1.427 (7)	C6—C7	1.366 (8)
N1—C8	1.295 (7)	C6—H6A	0.9300
N1—N2	1.393 (6)	C7—H7A	0.9300
C1—H1B	0.9600	C10—C11	1.498 (8)
C1—H1C	0.9600	C11—H11A	0.9700
C1—H1D	0.9600	C11—H11B	0.9700
C12—C18	1.735 (5)	C12—C13	1.517 (7)
S2—C11	1.788 (7)	C12—H12A	0.9800
S2—C12	1.832 (5)	C13—C14	1.364 (8)
O2—C10	1.200 (7)	C13—C18	1.381 (7)
C2—C7	1.358 (8)	C14—C15	1.367 (8)
C2—C3	1.388 (7)	C14—H14A	0.9300
N2—C9	1.292 (7)	C15—C16	1.383 (8)
N3—C10	1.375 (7)	C15—H15A	0.9300
N3—C9	1.399 (7)	C16—C17	1.365 (8)
N3—C12	1.441 (7)	C17—C18	1.360 (8)
C3—C4	1.362 (8)	C17—H17A	0.9300
C3—H3A	0.9300		
C9—S1—C8	86.0 (3)	N2—C9—S1	116.2 (4)
C2—O1—C1	117.6 (5)	N3—C9—S1	124.3 (4)
C8—N1—N2	113.1 (4)	O2—C10—N3	124.0 (5)
O1—C1—H1B	109.5	O2—C10—C11	125.4 (5)
O1—C1—H1C	109.5	N3—C10—C11	110.6 (5)
H1B—C1—H1C	109.5	C10—C11—S2	108.7 (4)
O1—C1—H1D	109.5	C10—C11—H11A	110.0
H1B—C1—H1D	109.5	S2—C11—H11A	110.0
H1C—C1—H1D	109.5	C10—C11—H11B	110.0

C11—S2—C12	92.5 (3)	S2—C11—H11B	110.0
C7—C2—O1	125.2 (5)	H11A—C11—H11B	108.3
C7—C2—C3	118.7 (5)	N3—C12—C13	112.7 (4)
O1—C2—C3	116.1 (5)	N3—C12—S2	104.8 (3)
C9—N2—N1	110.6 (5)	C13—C12—S2	111.2 (4)
C10—N3—C9	122.2 (5)	N3—C12—H12A	109.3
C10—N3—C12	119.5 (4)	C13—C12—H12A	109.3
C9—N3—C12	118.3 (4)	S2—C12—H12A	109.3
C4—C3—C2	120.7 (5)	C14—C13—C18	117.3 (5)
C4—C3—H3A	119.7	C14—C13—C12	121.2 (5)
C2—C3—H3A	119.7	C18—C13—C12	121.5 (5)
C3—C4—C5	121.5 (5)	C13—C14—C15	123.3 (5)
C3—C4—H4A	119.2	C13—C14—H14A	118.4
C5—C4—H4A	119.2	C15—C14—H14A	118.4
C4—C5—C6	116.7 (5)	C14—C15—C16	117.3 (5)
C4—C5—C8	123.5 (5)	C14—C15—H15A	121.4
C6—C5—C8	119.8 (5)	C16—C15—H15A	121.4
C7—C6—C5	122.1 (5)	C17—C16—C15	121.2 (5)
C7—C6—H6A	119.0	C17—C16—C11	119.7 (4)
C5—C6—H6A	119.0	C15—C16—C11	119.0 (5)
C2—C7—C6	120.2 (5)	C18—C17—C16	119.4 (5)
C2—C7—H7A	119.9	C18—C17—H17A	120.3
C6—C7—H7A	119.9	C16—C17—H17A	120.3
N1—C8—C5	122.9 (4)	C17—C18—C13	121.5 (5)
N1—C8—S1	114.1 (4)	C17—C18—C12	117.8 (4)
C5—C8—S1	123.0 (4)	C13—C18—C12	120.6 (4)
N2—C9—N3	119.5 (5)		
C1—O1—C2—C7	5.7 (9)	C12—N3—C10—O2	-177.0 (6)
C1—O1—C2—C3	-174.3 (6)	C9—N3—C10—C11	179.2 (5)
C8—N1—N2—C9	-0.6 (7)	C12—N3—C10—C11	2.1 (7)
C7—C2—C3—C4	0.7 (10)	O2—C10—C11—S2	-169.0 (5)
O1—C2—C3—C4	-179.3 (6)	N3—C10—C11—S2	12.0 (7)
C2—C3—C4—C5	0.6 (10)	C12—S2—C11—C10	-17.2 (5)
C3—C4—C5—C6	-2.5 (9)	C10—N3—C12—C13	106.6 (5)
C3—C4—C5—C8	177.4 (6)	C9—N3—C12—C13	-70.6 (6)
C4—C5—C6—C7	3.2 (10)	C10—N3—C12—S2	-14.5 (6)
C8—C5—C6—C7	-176.8 (6)	C9—N3—C12—S2	168.3 (4)
O1—C2—C7—C6	179.9 (6)	C11—S2—C12—N3	17.6 (4)
C3—C2—C7—C6	-0.1 (10)	C11—S2—C12—C13	-104.5 (4)
C5—C6—C7—C2	-1.9 (10)	N3—C12—C13—C14	-31.0 (7)
N2—N1—C8—C5	-179.4 (5)	S2—C12—C13—C14	86.3 (6)
N2—N1—C8—S1	-0.9 (7)	N3—C12—C13—C18	151.9 (5)
C4—C5—C8—N1	-173.3 (6)	S2—C12—C13—C18	-90.8 (5)
C6—C5—C8—N1	6.6 (9)	C18—C13—C14—C15	-1.7 (8)
C4—C5—C8—S1	8.3 (8)	C12—C13—C14—C15	-178.9 (5)
C6—C5—C8—S1	-171.8 (5)	C13—C14—C15—C16	0.1 (9)
C9—S1—C8—N1	1.5 (5)	C14—C15—C16—C17	1.4 (8)

C9—S1—C8—C5	-179.9 (5)	C14—C15—C16—C11	-178.2 (4)
N1—N2—C9—N3	179.6 (5)	C15—C16—C17—C18	-1.3 (8)
N1—N2—C9—S1	1.8 (7)	C11—C16—C17—C18	178.3 (4)
C10—N3—C9—N2	-175.9 (5)	C16—C17—C18—C13	-0.3 (8)
C12—N3—C9—N2	1.1 (8)	C16—C17—C18—C12	179.4 (4)
C10—N3—C9—S1	1.7 (8)	C14—C13—C18—C17	1.7 (8)
C12—N3—C9—S1	178.7 (4)	C12—C13—C18—C17	179.0 (5)
C8—S1—C9—N2	-1.9 (5)	C14—C13—C18—C12	-178.0 (4)
C8—S1—C9—N3	-179.6 (5)	C12—C13—C18—C12	-0.7 (7)
C9—N3—C10—O2	0.1 (9)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C4—H4 <i>A</i> ...S1	0.93	2.79	3.180 (7)	106
C6—H6 <i>A</i> ...N1	0.93	2.55	2.856 (8)	100
C12—H12 <i>A</i> ...C12	0.98	2.63	3.063 (5)	107
C14—H14 <i>A</i> ...N3	0.93	2.54	2.863 (8)	101
C14—H14 <i>A</i> ...O1 ⁱ	0.93	2.41	3.219 (7)	146

Symmetry code: (i) $-x+2, -y+1, -z$.