

1-Methyl-3-phenylsulfonyl-2-piperidone

Julio Zukerman-Schpector,^{a*} Paulo R. Olivato,^b Carlos R. Cerqueira Jr,^b Elisângela Vinható^b and Edward R. T. Tiekink^c

^aDepartment of Chemistry, Universidade Federal de São Carlos, 13565-905 São Carlos, SP, Brazil, ^bChemistry Institute, University of São Paulo, 05508-000 São Paulo, SP, Brazil, and ^cDepartment of Chemistry, The University of Texas at San Antonio, One UTSA Circle, San Antonio, Texas 78249-0698, USA

Correspondence e-mail: julio@power.ufscar.br


Received 4 April 2008; accepted 5 April 2008

Key indicators: single-crystal X-ray study; $T = 98$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.047; wR factor = 0.119; data-to-parameter ratio = 17.7.

The piperidone ring in the title compound, $\text{C}_{12}\text{H}_{15}\text{NO}_3\text{S}$, has a slightly distorted half-chair conformation with the methyl, carbonyl and phenylsulfonyl ring substituents occupying equatorial, equatorial and axial positions, respectively. Molecules are connected into centrosymmetric dimers *via* $\text{C}-\text{H}\cdots\text{O}$ interactions and these associate into layers *via* $\text{C}-\text{H}\cdots\text{O}-\text{S}$ contacts. Further $\text{C}-\text{H}\cdots\text{O}$ interactions involving both the carbonyl and sulfonyl O atoms consolidate the crystal packing by providing connections between the layers.

Related literature

For related structures, see: Zukerman-Schpector *et al.* (1999, 2006). For related literature, see: Distefano *et al.* (1991); Olivato *et al.* (1992, 1997, 2003, 2004); Dal Colle *et al.* (1995). For ring conformational analysis, see: Cremer & Pople (1975). For the synthesis, see: Drabowicz *et al.* (1983); Zoretic & Soja (1976).



Experimental

Crystal data

$\text{C}_{12}\text{H}_{15}\text{NO}_3\text{S}$
 $M_r = 253.32$
 Monoclinic, $P2_1/n$
 $a = 9.0191$ (16) Å
 $b = 10.4920$ (18) Å
 $c = 13.446$ (3) Å
 $\beta = 107.861$ (3)°

$V = 1211.1$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.26$ mm⁻¹
 $T = 98$ (2) K
 $0.25 \times 0.18 \times 0.10$ mm

Data collection

Rigaku AFC12κ/SATURN724
 diffractometer
 Absorption correction: multi-scan
 (ABSCOR; Higashi, 1995)
 $T_{\min} = 0.945$, $T_{\max} = 0.974$

5193 measured reflections
 2729 independent reflections
 2549 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.118$
 $S = 1.12$
 2729 reflections

154 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.39$ e Å⁻³
 $\Delta\rho_{\min} = -0.45$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C2}-\text{H2}\cdots\text{O1}^i$	1.00	2.29	3.272 (2)	168
$\text{C6}-\text{H6B}\cdots\text{O2}^{ii}$	0.98	2.55	3.424 (3)	148
$\text{C11}-\text{H11}\cdots\text{O3}^{iii}$	0.95	2.62	3.224 (3)	122
$\text{C4}-\text{H4A}\cdots\text{O1}^{iv}$	0.99	2.48	3.328 (2)	144

Symmetry codes: (i) $-x + 2, -y, -z + 1$; (ii) $-x + 2, -y + 1, -z + 1$; (iii) $-x + 1, -y, -z + 1$; (iv) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1999); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEPII* (Johnson, 1976) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *SHELXL97*.

We thank FAPESP and CNPq for financial support. CRC and EV thank CNPq for doctoral fellowships; PRO and JZ-S thank CNPq for fellowships (Brazil). Support from UTSA, CNPq and FAPESP to allow JZ-S to spend a sabbatical at UTSA is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NG2443).

References

- Altomare, A., Burla, M. C., Camalli, M., Cascarano, G. L., Giacovazzo, C., Guagliardi, A., Moliterni, A. G. G., Polidori, G. & Spagna, R. (1999). *J. Appl. Cryst.* **32**, 115–119.
- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Dal Colle, M., Bertolasi, V., De Palo, M., Distefano, G., Jones, D., Modelli, A. & Olivato, P. R. (1995). *J. Chem. Phys.* **99**, 15011–15017.
- Distefano, G., Dal Colle, M., Bertolasi, V., Olivato, P. R., Bonfada, E. & Mondino, M. G. (1991). *J. Chem. Soc. Perkin Trans. 2*, pp. 1195–1199.
- Drabowicz, J., Lyzwa, P. & Mikolajczyk, M. (1983). *Phosphorus Sulfur Silicon*, **17**, 169–172.
- Higashi, T. (1995). *ABSCOR*. Rigaku Corporation, Tokyo, Japan.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Olivato, P. R., Bonfada, E. & Rittner, R. (1992). *Magn. Reson. Chem.* **30**, 81–84.
- Olivato, P. R., Guerrero, S. A. & Rittner, R. (1997). *Phosphorus Sulfur Silicon*, **130**, 155–174.
- Olivato, P. R., Hui, M. L. T., Rodrigues, A., Ruiz Filho, R., Rittner, R., Zukerman-Schpector, J., Distefano, G. & Dal Colle, M. (2003). *J. Mol. Struct.* **645**, 259–271.
- Olivato, P. R., Reis, A. K. C. A., Ruiz Filho, R., Zukerman-Schpector, J. & Rittner, R. (2004). *J. Mol. Struct. THEOCHEM*, **677**, 199–210.

Rigaku (2005). *CrystalClear*. Rigaku Americas Corporation, The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Zoretic, P. A. & Soja, P. (1976). *J. Org. Chem.* **41**, 3587–3589.

Zukerman-Schpector, J., Maganhi, S., Olivato, P. R., Vinhato, E. & Cerqueira, C. R. (2006). *Z. Kristallogr. New Cryst. Struct.* **221**, 165–166.
Zukerman-Schpector, J., Olivato, P. R., Bueno, E. & Guerrero, S. A. (1999). *Z. Kristallogr. New Cryst. Struct.* **214**, 563–564.

supporting information

Acta Cryst. (2008). E64, o835–o836 [doi:10.1107/S1600536808009288]

1-Methyl-3-phenylsulfonyl-2-piperidone

Julio Zukerman-Schpector, Paulo R. Olivato, Carlos R. Cerqueira Jr, Elisângela Vinhato and Edward R. T. Tiekink

S1. Comment

The title compound (I), Fig. 1, was studied as a part of an on-going investigation of conformational and electronic aspects of different classes of β -keto-sulfones, *i.e.* α -phenylsulfonyl -acetones, -acetophenones and -cyclohexanones, utilizing spectroscopic, theoretical and X-ray diffraction methods (Dal Colle *et al.*, 1995; Zukerman-Schpector *et al.*, 1999; 2006).

The piperidone ring has a slightly distorted half-chair conformation with a tendency towards a half-boat conformation: the ring-puckering parameters are $q_2 = 0.340(2) \text{ \AA}$, $q_3 = 0.332(2) \text{ \AA}$, $QT = 0.476(2)^\circ$, $\varphi_2 = -145.0(3)^\circ$ (Cremer & Pople, 1975). The ring substituents, *i.e.* *N*-methyl, *C*-carbonyl and *C*-phenylsulfonyl, occupy equatorial, equatorial and axial positions, respectively.

The crystal packing is dominated by $C-H\cdots O$ interactions, Table 1. Centrosymmetrically related molecules of (I) are connected into dimeric aggregates *via* $C2-H\cdots O1$ contacts and these are linked into layers stacked along (1 0 1) *via* $C6-H\cdots O2$ contacts. Connections between layers are also of the type $C-H\cdots O$ and serve to consolidate the crystal packing.

S2. Experimental

Initially, the 3-phenylsulfonyl-1-methyl-2-piperidone was obtained from the reaction of 1-methyl-2-piperidinone and diphenyl disulfide with LDA in THF as described in the literature (Zoretic and Soja, 1976). The product was oxidized with H_2O_2 and SeO_2 (as catalyst) in methanol (Drabowicz *et al.* 1983) to give compound (I). After extraction with chloroform and subsequent evaporation, a crude solid was obtained. This product was subjected to flash chromatography with a solution of ethyl acetate and acetone in a 7:3 ratio. Suitable crystals were obtained by vapor diffusion from chloroform/n-hexane at 283 K.; m.p. 414–415 K. IR (cm^{-1}): $\nu(C=O)$ 1652, $\nu(SO_2)(as)$ 1307, $\nu(SO_2)(s)$ 1148. NMR ($CDCl_3$, p.p.m.): δ 1.79–2.74 (4H, m), 2.95 (3H, s), 3.30–3.48 (2H, m), 3.97 (1H, triplet, $J = 6.1$ Hz), 7.53–7.57 (2H, m, aryl-H), 7.62–7.67 (1H, m, aryl-H), 7.92–7.94 (2H, m, aryl-H). Analysis found: C 56.86, H 6.04, N 5.58; $C_{12}H_{15}O_3NS$ requires: C 56.89, H 5.97, N 5.53%.

S3. Refinement

All H atoms were included in the riding-model approximation with $C-H = 0.95 - 1.00 \text{ \AA}$, and with $U_{iso}(H) = 1.5U_{eq}(\text{methyl-C})$ or $1.2U_{eq}(\text{remaining-C})$.





Figure 1

Molecular structure of (I) showing atom labelling and displacement ellipsoids at the 50% probability level.

**Figure 2**

Crystal packing in (I) highlighting the C—H...O hydrogen bonding contacts (orange dashed lines) leading to the formation of dimeric aggregates and the overall layer arrangement.

1-Methyl-3-phenylsulfonyl-2-piperidone

Crystal data

$C_{12}H_{15}NO_3S$

$M_r = 253.32$

Monoclinic, $P2_1/n$

Hall symbol: $-P 2_1n$

$a = 9.0191 (16) \text{ \AA}$

$b = 10.4920 (18) \text{ \AA}$

$c = 13.446 (3) \text{ \AA}$

$\beta = 107.861 (3)^\circ$

$V = 1211.1 (4) \text{ \AA}^3$

$Z = 4$

$F(000) = 536$

$D_x = 1.389 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71070 \text{ \AA}$

Cell parameters from 4417 reflections

$\theta = 2.4\text{--}40.6^\circ$

$\mu = 0.26 \text{ mm}^{-1}$

$T = 98 \text{ K}$

Block, colourless

$0.25 \times 0.18 \times 0.10 \text{ mm}$

Data collection

Rigaku AFC12κ/SATURN724
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 ω scans
Absorption correction: multi-scan
(*ABSCOR*; Higashi, 1995)
 $T_{\min} = 0.945$, $T_{\max} = 0.974$

5193 measured reflections
2729 independent reflections
2549 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 2.5^\circ$
 $h = -10 \rightarrow 11$
 $k = -13 \rightarrow 11$
 $l = -17 \rightarrow 7$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.047$
 $wR(F^2) = 0.118$
 $S = 1.12$
2729 reflections
154 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0539P)^2 + 0.7818P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.39 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.45 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.76305 (5)	0.20597 (4)	0.49677 (3)	0.01768 (15)
O1	1.13033 (15)	0.14856 (13)	0.56636 (9)	0.0186 (3)
O2	0.79732 (18)	0.33520 (13)	0.53443 (11)	0.0265 (3)
O3	0.60745 (15)	0.17739 (15)	0.43018 (11)	0.0275 (3)
N1	1.11807 (17)	0.30781 (15)	0.45077 (11)	0.0170 (3)
C1	1.0585 (2)	0.20646 (17)	0.48622 (13)	0.0147 (3)
C2	0.8952 (2)	0.16205 (17)	0.42479 (13)	0.0153 (3)
H2	0.8969	0.0669	0.4210	0.018*
C3	0.8348 (2)	0.21313 (18)	0.31267 (14)	0.0196 (4)
H3A	0.8843	0.1657	0.2676	0.024*
H3B	0.7208	0.1998	0.2850	0.024*
C4	0.8711 (2)	0.35463 (19)	0.30992 (14)	0.0214 (4)
H4A	0.8313	0.3865	0.2372	0.026*
H4B	0.8188	0.4027	0.3530	0.026*
C5	1.0457 (2)	0.37553 (19)	0.35160 (14)	0.0208 (4)
H5A	1.0942	0.3460	0.2989	0.025*

H5B	1.0668	0.4679	0.3625	0.025*
C6	1.2777 (2)	0.3460 (2)	0.50751 (14)	0.0213 (4)
H6A	1.3156	0.2956	0.5717	0.032*
H6B	1.2794	0.4367	0.5254	0.032*
H6C	1.3448	0.3314	0.4635	0.032*
C7	0.8027 (2)	0.10460 (17)	0.60682 (13)	0.0166 (3)
C8	0.9117 (2)	0.14160 (19)	0.70016 (14)	0.0199 (4)
H8	0.9681	0.2189	0.7041	0.024*
C9	0.9368 (2)	0.0641 (2)	0.78751 (14)	0.0211 (4)
H9	1.0114	0.0879	0.8516	0.025*
C10	0.8528 (2)	-0.04843 (19)	0.78116 (14)	0.0215 (4)
H10	0.8704	-0.1013	0.8411	0.026*
C11	0.7431 (2)	-0.08410 (19)	0.68753 (15)	0.0209 (4)
H11	0.6856	-0.1607	0.6839	0.025*
C12	0.7177 (2)	-0.00801 (18)	0.59943 (14)	0.0185 (4)
H12	0.6437	-0.0322	0.5352	0.022*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0168 (2)	0.0151 (3)	0.0225 (2)	0.00200 (16)	0.00796 (18)	0.00263 (16)
O1	0.0179 (6)	0.0205 (7)	0.0155 (6)	0.0007 (5)	0.0026 (5)	0.0026 (5)
O2	0.0382 (8)	0.0134 (7)	0.0353 (8)	0.0021 (6)	0.0223 (7)	0.0004 (6)
O3	0.0144 (6)	0.0339 (8)	0.0324 (7)	0.0022 (6)	0.0044 (6)	0.0120 (6)
N1	0.0153 (7)	0.0180 (8)	0.0159 (7)	-0.0018 (6)	0.0023 (6)	0.0013 (6)
C1	0.0153 (8)	0.0149 (9)	0.0147 (7)	-0.0002 (6)	0.0056 (6)	-0.0023 (6)
C2	0.0153 (8)	0.0141 (8)	0.0163 (8)	0.0000 (6)	0.0046 (6)	-0.0010 (6)
C3	0.0181 (9)	0.0224 (10)	0.0154 (8)	0.0001 (7)	0.0008 (7)	-0.0005 (7)
C4	0.0218 (9)	0.0215 (9)	0.0178 (8)	0.0016 (7)	0.0015 (7)	0.0027 (7)
C5	0.0242 (9)	0.0193 (9)	0.0174 (8)	-0.0015 (7)	0.0043 (7)	0.0044 (7)
C6	0.0187 (9)	0.0233 (10)	0.0205 (8)	-0.0054 (7)	0.0039 (7)	0.0003 (7)
C7	0.0180 (8)	0.0145 (8)	0.0194 (8)	0.0015 (7)	0.0090 (7)	0.0004 (7)
C8	0.0192 (9)	0.0192 (9)	0.0232 (9)	-0.0026 (7)	0.0096 (7)	-0.0035 (7)
C9	0.0197 (9)	0.0254 (10)	0.0191 (8)	-0.0010 (7)	0.0072 (7)	-0.0037 (7)
C10	0.0230 (9)	0.0243 (10)	0.0205 (8)	0.0027 (8)	0.0113 (7)	0.0037 (7)
C11	0.0204 (9)	0.0180 (9)	0.0274 (9)	-0.0019 (7)	0.0120 (7)	0.0002 (7)
C12	0.0173 (8)	0.0170 (9)	0.0214 (8)	-0.0014 (7)	0.0061 (7)	-0.0018 (7)

Geometric parameters (Å, °)

S1—O3	1.4457 (15)	C5—H5A	0.9900
S1—O2	1.4472 (15)	C5—H5B	0.9900
S1—C7	1.7674 (18)	C6—H6A	0.9800
S1—C2	1.8101 (18)	C6—H6B	0.9800
O1—C1	1.233 (2)	C6—H6C	0.9800
N1—C1	1.343 (2)	C7—C8	1.391 (3)
N1—C6	1.463 (2)	C7—C12	1.395 (3)
N1—C5	1.475 (2)	C8—C9	1.389 (3)

C1—C2	1.524 (2)	C8—H8	0.9500
C2—C3	1.534 (2)	C9—C10	1.391 (3)
C2—H2	1.0000	C9—H9	0.9500
C3—C4	1.523 (3)	C10—C11	1.393 (3)
C3—H3A	0.9900	C10—H10	0.9500
C3—H3B	0.9900	C11—C12	1.388 (3)
C4—C5	1.517 (3)	C11—H11	0.9500
C4—H4A	0.9900	C12—H12	0.9500
C4—H4B	0.9900		
O3—S1—O2	118.26 (9)	N1—C5—C4	112.68 (15)
O3—S1—C7	107.61 (9)	N1—C5—H5A	109.1
O2—S1—C7	107.71 (9)	C4—C5—H5A	109.1
O3—S1—C2	106.81 (9)	N1—C5—H5B	109.1
O2—S1—C2	108.70 (8)	C4—C5—H5B	109.1
C7—S1—C2	107.28 (8)	H5A—C5—H5B	107.8
C1—N1—C6	117.93 (15)	N1—C6—H6A	109.5
C1—N1—C5	126.04 (15)	N1—C6—H6B	109.5
C6—N1—C5	115.53 (15)	H6A—C6—H6B	109.5
O1—C1—N1	122.70 (16)	N1—C6—H6C	109.5
O1—C1—C2	118.95 (16)	H6A—C6—H6C	109.5
N1—C1—C2	118.35 (15)	H6B—C6—H6C	109.5
C1—C2—C3	114.75 (15)	C8—C7—C12	121.42 (17)
C1—C2—S1	108.53 (11)	C8—C7—S1	119.54 (14)
C3—C2—S1	110.06 (12)	C12—C7—S1	118.96 (14)
C1—C2—H2	107.8	C9—C8—C7	119.10 (18)
C3—C2—H2	107.8	C9—C8—H8	120.4
S1—C2—H2	107.8	C7—C8—H8	120.4
C4—C3—C2	110.50 (15)	C8—C9—C10	120.03 (17)
C4—C3—H3A	109.6	C8—C9—H9	120.0
C2—C3—H3A	109.6	C10—C9—H9	120.0
C4—C3—H3B	109.6	C9—C10—C11	120.40 (17)
C2—C3—H3B	109.6	C9—C10—H10	119.8
H3A—C3—H3B	108.1	C11—C10—H10	119.8
C5—C4—C3	109.79 (16)	C12—C11—C10	120.17 (18)
C5—C4—H4A	109.7	C12—C11—H11	119.9
C3—C4—H4A	109.7	C10—C11—H11	119.9
C5—C4—H4B	109.7	C11—C12—C7	118.87 (17)
C3—C4—H4B	109.7	C11—C12—H12	120.6
H4A—C4—H4B	108.2	C7—C12—H12	120.6
C6—N1—C1—O1	3.1 (3)	C1—N1—C5—C4	21.7 (3)
C5—N1—C1—O1	174.61 (17)	C6—N1—C5—C4	-166.58 (16)
C6—N1—C1—C2	-177.19 (15)	C3—C4—C5—N1	-47.8 (2)
C5—N1—C1—C2	-5.7 (3)	O3—S1—C7—C8	-154.80 (15)
O1—C1—C2—C3	-163.27 (15)	O2—S1—C7—C8	-26.27 (17)
N1—C1—C2—C3	17.0 (2)	C2—S1—C7—C8	90.58 (15)
O1—C1—C2—S1	73.16 (18)	O3—S1—C7—C12	21.86 (17)

N1—C1—C2—S1	-106.55 (15)	O2—S1—C7—C12	150.39 (14)
O3—S1—C2—C1	172.73 (12)	C2—S1—C7—C12	-92.76 (15)
O2—S1—C2—C1	44.09 (14)	C12—C7—C8—C9	0.5 (3)
C7—S1—C2—C1	-72.12 (14)	S1—C7—C8—C9	177.05 (14)
O3—S1—C2—C3	46.40 (15)	C7—C8—C9—C10	-0.5 (3)
O2—S1—C2—C3	-82.24 (14)	C8—C9—C10—C11	0.0 (3)
C7—S1—C2—C3	161.55 (12)	C9—C10—C11—C12	0.5 (3)
C1—C2—C3—C4	-44.1 (2)	C10—C11—C12—C7	-0.5 (3)
S1—C2—C3—C4	78.63 (17)	C8—C7—C12—C11	0.0 (3)
C2—C3—C4—C5	59.6 (2)	S1—C7—C12—C11	-176.58 (14)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
C2—H2 \cdots O1 ⁱ	1.00	2.29	3.272 (2)	168
C6—H6B \cdots O2 ⁱⁱ	0.98	2.55	3.424 (3)	148
C11—H11 \cdots O3 ⁱⁱⁱ	0.95	2.62	3.224 (3)	122
C4—H4A \cdots O1 ^{iv}	0.99	2.48	3.328 (2)	144

Symmetry codes: (i) $-x+2, -y, -z+1$; (ii) $-x+2, -y+1, -z+1$; (iii) $-x+1, -y, -z+1$; (iv) $x-1/2, -y+1/2, z-1/2$.