

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

catena-Poly[[dichloridonickel(II)]- $\mu$ -1,3-di-4-pyridylpropane]

Chun-Sheng Zhou\* and Guo-Chun Zhang

Department of Chemistry, ShangLuo University, ShangLuo, Shaanxi 726000, People's Republic of China

Correspondence e-mail: slzhousc@126.com


Received 19 June 2008; accepted 22 June 2008

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.039;  $wR$  factor = 0.068; data-to-parameter ratio = 15.1.

The title compound,  $[\text{NiCl}_2(\text{C}_{13}\text{H}_{14}\text{N}_2)]_n$ , is a one-dimensional polymer built up from alternating  $\text{NiCl}_2$  units and bridging 1,3-di-4-pyridylpropane ligands. The Ni atom has a distorted tetrahedral coordination formed by the Cl atoms and two N atoms from two ligands. A mirror plane passes through the central methylene group of the propyl chain.

## Related literature

For a closely related structure, see: Zhang & Yu (2007). For related literature, see: Carlucci *et al.* (2002); Hennigar *et al.* (1997); Yaghi *et al.* (1998); Dalbavie *et al.* (2002); Ghosh *et al.* (2006); Marshall & Grushin (2005); Masood *et al.* (1994); McConnell & Nuttall (1978); Wu *et al.* (1999).



## Experimental

## Crystal data

 $[\text{NiCl}_2(\text{C}_{13}\text{H}_{14}\text{N}_2)]$  $M_r = 327.87$ 

Monoclinic,  $P2_1/m$   
 $a = 5.1928$  (17) Å  
 $b = 12.972$  (4) Å  
 $c = 10.492$  (3) Å  
 $\beta = 93.588$  (6)°  
 $V = 705.3$  (4) Å<sup>3</sup>

$Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 1.74$  mm<sup>-1</sup>  
 $T = 298$  (2) K  
 $0.25 \times 0.20 \times 0.16$  mm

## Data collection

Bruker APEX area-detector diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.671$ ,  $T_{\max} = 0.769$

3581 measured reflections  
 1328 independent reflections  
 763 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.046$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$   
 $wR(F^2) = 0.067$   
 $S = 0.87$   
 1328 reflections

88 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.44$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.44$  e Å<sup>-3</sup>

Data collection: SMART (Bruker, 1998); cell refinement: SAINT (Bruker, 1999); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXTL.

The authors are grateful to ShangLuo University for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2360).

## References

- Bruker (1998). SMART. Bruker AXS Inc, Madison, Wisconsin, USA.  
 Bruker (1999). SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Carlucci, L., Ciani, G., Proserpio, D. M. & Rizzato, S. (2002). *CrystEngComm*, **4**, 121–129.  
 Dalbavie, J. O., Regnouf-de-Vains, J. B., Lamartine, R., Perrin, R., Lecocq, L. & Fenet, B. (2002). *Eur. J. Inorg. Chem.* pp. 901–909.  
 Ghosh, T., Maiya, B. G. & Samanta, A. (2006). *Dalton Trans.* pp. 795–802.  
 Hennigar, T. L., MacQuarrie, D. C., Losier, P., Gogers, R. D. & Zaworotko, M. J. (1997). *Angew. Chem. Int. Ed. Engl.* **36**, 972–973.  
 Marshall, W. J. & Grushin, V. V. (2005). *Can. J. Chem.* **83**, 640–645.  
 Masood, Md. A., Hodgson, D. J. & Zacharias, P. S. (1994). *Inorg. Chim. Acta*, **221**, 99–103.  
 McConnell, A. A. & Nuttall, R. H. (1978). *J. Mol. Struct.* **49**, 207–213.  
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Wu, H. P., Janiak, C., Rheinwald, G. & Lang, H. (1999). *J. Chem. Soc. Dalton Trans.* pp. 183–190.  
 Yaghi, O. M., Li, H., Davis, C., Richardson, D. & Groy, T. L. (1998). *Acc. Chem. Res.* **31**, 474–484.  
 Zhang, L. & Yu, J. (2007). *Acta Cryst.* **E63**, m1759.

## supporting information

*Acta Cryst.* (2008). E64, m966 [doi:10.1107/S1600536808018862]

**catena-Poly[[dichloridonickel(II)]- $\mu$ -1,3-di-4-pyridylpropane]****Chun-Sheng Zhou and Guo-Chun Zhang****S1. Comment**

Recent years have seen the evolution of a new class of coordination polymers known collectively as metal organic framework materials (Yaghi *et al.*, 1998). The most common approach for producing coordination polymers and metal organic framework materials is through the self-assembly of metal centers with appropriate organic linker species to promote extended topologies (Hennigar *et al.*, 1997). Conformationally flexible ligands are typical building elements in the molecular interlocked/intertwined species. Some work on the self-assembly of coordination networks have been reported in the presence of 1,3-di-4-pyridylpropane (bpp) ligand (Carlucci *et al.*, 2002). In this paper, we report here the synthesis and crystal structure of the title compound (I).


The Ni atom in the title complex has a distorted tetrahedral coordination formed by the chlorine atoms and two nitrogen from two separate bpp ligands (Fig. 1). The distances of Ni1—Cl and Ni1—Cl are 2.2533 (17) and 2.2382 (16) Å, respectively. Figure 1 show that this one-dimensional polymer built up from alternating (NiCl<sub>2</sub>) units and bridging 1,3-di-4-pyridylpropane ligands. Some other NiCl<sub>2</sub> complexes with tetrahedral coordination geometries have been reported (Wu *et al.*, 1999; Dalbavie *et al.*, 2002; Masood *et al.*, 1994; McConnell & Nuttall, 1978; Ghosh *et al.*, 2006; Marshall & Grushin, 2005).

**S2. Experimental**

Bpp (0.21, 0.1 mmol), NiCl<sub>2</sub> (0.22 g, 0.012 mmol), were added in a mixed solvent of methanol and acetonitrile, the mixture was heated for six hours under reflux. during the process stirring and influx were required. The resultant was then filtered to give a pure solution which was infiltrated by diethyl ether freely in a closed vessel, a weeks later some single crystals of the size suitable for X-Ray diffraction analysis.

**S3. Refinement**

All H atoms were fixed geometrically and treated as riding with C—H = 0.93 Å (aromatic) and U<sub>iso</sub>(H) = 1.2U<sub>eq</sub>(C).

**Figure 1**

The structure of (I), showing the atom labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii.

### catena-Poly[[dichloridonickel(II)]- $\mu$ -1,3-di-4-pyridylpropane]

#### Crystal data

[NiCl<sub>2</sub>(C<sub>13</sub>H<sub>14</sub>N<sub>2</sub>)]

$M_r = 327.87$

Monoclinic,  $P2_1/m$

Hall symbol: -P 2yb

$a = 5.1928$  (17) Å

$b = 12.972$  (4) Å

$c = 10.492$  (3) Å

$\beta = 93.588$  (6)°

$V = 705.3$  (4) Å<sup>3</sup>

$Z = 2$

$F(000) = 336$

$D_x = 1.544$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1328 reflections

$\theta = 2.5$ – $25.2$ °

$\mu = 1.74$  mm<sup>-1</sup>

$T = 298$  K

Bloc, green

$0.25 \times 0.20 \times 0.16$  mm

#### Data collection

Bruker APEX area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.671$ ,  $T_{\max} = 0.769$

3581 measured reflections

1328 independent reflections

763 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.046$

$\theta_{\max} = 25.2$ °,  $\theta_{\min} = 2.5$ °

$h = -6 \rightarrow 5$

$k = -13 \rightarrow 15$

$l = -12 \rightarrow 12$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.067$

$S = 0.87$

1328 reflections

88 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.009P)^2 + 0.821P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.44$  e Å<sup>-3</sup>

$\Delta\rho_{\min} = -0.44$  e Å<sup>-3</sup>

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Ni1	0.45214 (16)	0.7500	0.22234 (6)	0.0554 (3)
Cl1	0.7422 (3)	0.7500	0.39091 (12)	0.0566 (4)
Cl2	0.5444 (3)	0.7500	0.01676 (12)	0.0654 (5)
N1	0.2418 (6)	0.6191 (2)	0.2414 (3)	0.0443 (8)
C1	0.2869 (7)	0.5560 (3)	0.3416 (3)	0.0501 (10)
H1	0.4209	0.5719	0.4013	0.060*
C3	-0.0579 (7)	0.4430 (3)	0.2738 (3)	0.0445 (10)
C5	0.0501 (8)	0.5930 (3)	0.1581 (3)	0.0555 (11)
H5	0.0171	0.6349	0.0871	0.067*
C4	-0.1015 (8)	0.5079 (3)	0.1707 (3)	0.0546 (11)
H4	-0.2344	0.4939	0.1097	0.066*
C2	0.1436 (8)	0.4690 (3)	0.3594 (3)	0.0521 (11)
H2	0.1827	0.4271	0.4300	0.063*
C7	-0.0598 (10)	0.2500	0.2643 (4)	0.0462 (14)
H7A	-0.0175	0.2500	0.1755	0.055*
H7B	0.1005	0.2500	0.3169	0.055*
C6	-0.2131 (7)	0.3473 (3)	0.2915 (3)	0.0521 (11)
H6A	-0.3677	0.3495	0.2347	0.063*
H6B	-0.2658	0.3448	0.3786	0.063*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ni1	0.0726 (7)	0.0426 (4)	0.0503 (4)	0.000	-0.0005 (4)	0.000
Cl1	0.0671 (12)	0.0479 (8)	0.0534 (8)	0.000	-0.0083 (7)	0.000
Cl2	0.0733 (13)	0.0780 (10)	0.0444 (8)	0.000	-0.0006 (7)	0.000
N1	0.052 (2)	0.0332 (17)	0.0466 (17)	0.0034 (16)	-0.0023 (16)	-0.0024 (14)
C1	0.050 (3)	0.047 (2)	0.052 (2)	0.003 (2)	-0.0080 (19)	0.0003 (19)
C3	0.049 (3)	0.030 (2)	0.055 (2)	0.006 (2)	0.003 (2)	-0.0070 (17)
C5	0.070 (3)	0.042 (2)	0.053 (2)	0.003 (2)	-0.009 (2)	0.0027 (19)
C4	0.062 (3)	0.047 (2)	0.052 (2)	0.001 (2)	-0.014 (2)	-0.0052 (19)
C2	0.066 (3)	0.039 (2)	0.050 (2)	0.005 (2)	-0.007 (2)	0.0072 (18)
C7	0.054 (4)	0.034 (3)	0.051 (3)	0.000	0.000 (3)	0.000
C6	0.046 (3)	0.043 (2)	0.068 (3)	-0.001 (2)	0.004 (2)	-0.0085 (19)

*Geometric parameters (Å, °)*

Ni1—N1	2.036 (3)	C5—C4	1.366 (5)
Ni1—N1 <sup>i</sup>	2.036 (3)	C5—H5	0.9300
Ni1—C12	2.2384 (16)	C4—H4	0.9300
Ni1—C11	2.2503 (16)	C2—H2	0.9300
N1—C5	1.327 (4)	C7—C6 <sup>ii</sup>	1.529 (4)
N1—C1	1.341 (4)	C7—C6	1.529 (4)
C1—C2	1.372 (5)	C7—H7A	0.9700
C1—H1	0.9300	C7—H7B	0.9700
C3—C2	1.377 (5)	C6—H6A	0.9700
C3—C4	1.378 (4)	C6—H6B	0.9700
C3—C6	1.499 (5)		
N1—Ni1—N1 <sup>i</sup>	113.06 (17)	C5—C4—C3	120.1 (4)
N1—Ni1—C12	104.07 (8)	C5—C4—H4	119.9
N1 <sup>i</sup> —Ni1—C12	104.07 (8)	C3—C4—H4	119.9
N1—Ni1—C11	105.05 (9)	C1—C2—C3	120.5 (3)
N1 <sup>i</sup> —Ni1—C11	105.05 (9)	C1—C2—H2	119.7
C12—Ni1—C11	125.74 (7)	C3—C2—H2	119.7
C5—N1—C1	116.6 (3)	C6 <sup>ii</sup> —C7—C6	111.3 (4)
C5—N1—Ni1	122.2 (2)	C6 <sup>ii</sup> —C7—H7A	109.4
C1—N1—Ni1	121.1 (3)	C6—C7—H7A	109.4
N1—C1—C2	122.6 (3)	C6 <sup>ii</sup> —C7—H7B	109.4
N1—C1—H1	118.7	C6—C7—H7B	109.4
C2—C1—H1	118.7	H7A—C7—H7B	108.0
C2—C3—C4	116.3 (4)	C3—C6—C7	111.7 (3)
C2—C3—C6	120.9 (3)	C3—C6—H6A	109.3
C4—C3—C6	122.7 (4)	C7—C6—H6A	109.3
N1—C5—C4	123.7 (3)	C3—C6—H6B	109.3
N1—C5—H5	118.1	C7—C6—H6B	109.3
C4—C5—H5	118.1	H6A—C6—H6B	107.9

Symmetry codes: (i)  $x, -y+3/2, z$ ; (ii)  $x, -y+1/2, z$ .