

2-[(*E*)-(5-Amino-2,3-diphenylquinoxalin-6-yl)iminomethyl]-4-chlorophenol

Hoong-Kun Fun,^{a*} Reza Kia^{a‡} and Paul R. Raithby^{b§}

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bChemistry Department, University of Bath, Claverton Down, Bath BA2 7AY, England

Correspondence e-mail: hkfun@usm.my


Received 28 June 2008; accepted 1 July 2008

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.053; wR factor = 0.159; data-to-parameter ratio = 19.9.

The title Schiff base compound, $C_{27}\text{H}_{19}\text{ClN}_4\text{O}$, features two intramolecular $\text{O}-\text{H}\cdots\text{N}$ and $\text{N}-\text{H}\cdots\text{N}$ hydrogen bonds involving the hydroxy and amino groups to generate $S(6)$ and $S(5)$ ring motifs, respectively. In the crystal structure, weak intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{N}$ interactions, together with $\pi-\pi$ contacts [centroid–centroid distances = 3.6294 (11)–3.6881 (11) \AA], link neighboring molecules.

Related literature

For related literature on hydrogen-bond motifs, see: Bernstein *et al.* (1995). For related structures, see: Corden *et al.* (1996); Fun *et al.* (2008); Govindasamy *et al.* (1999). For applications and bioactivities, see: Anderson *et al.* (1997); Cohen & Schmidt (1964); Granovski *et al.* (1993); Li & Chang (1991); Shahrokhian *et al.* (2000); Unaleroglu & Hökelek (2002).



‡ Address of first Postdoctoral position: Chemistry Department, University of Bath, Claverton Down, Bath BA2 7AY, England.

§ Additional correspondence author, e-mail: p.r.raithby@bath.ac.uk.

Experimental

Crystal data

$C_{27}\text{H}_{19}\text{ClN}_4\text{O}$	$V = 2098.32\text{ (18) \AA}^3$
$M_r = 450.91$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 22.8728\text{ (11) \AA}$	$\mu = 0.21\text{ mm}^{-1}$
$b = 7.3068\text{ (4) \AA}$	$T = 100.0\text{ (1) K}$
$c = 12.5632\text{ (6) \AA}$	$0.55 \times 0.09 \times 0.07\text{ mm}$
$\beta = 92.037\text{ (2)}^\circ$	

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	25593 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2005)	6180 independent reflections
$T_{\min} = 0.843$, $T_{\max} = 0.985$	4118 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.062$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.159$	$\Delta\rho_{\text{max}} = 0.40\text{ e \AA}^{-3}$
$S = 1.01$	$\Delta\rho_{\text{min}} = -0.41\text{ e \AA}^{-3}$
6180 reflections	
310 parameters	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1O1 \cdots N1	0.93 (3)	1.72 (3)	2.584 (2)	153 (2)
N4—H1N4 \cdots O1 ⁱ	1.01 (3)	2.47 (3)	3.099 (2)	120.4 (18)
N4—H2N4 \cdots N2	1.04 (3)	2.31 (3)	2.750 (2)	104.3 (17)
C27—H27A \cdots N3 ⁱⁱ	0.93	2.62	3.373 (2)	138

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $x, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *APEX2*; data reduction: *SAINT* (Bruker, 2005); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2003).

HKF and RK thank the Malaysian Government and Universiti Sains Malaysia for Science Fund grant No. 305/PFIZIK/613312. RK thanks the Universiti Sains Malaysia and the University of Bath for post-doctoral research fellowships.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2279).

References

- Anderson, O. P., Cour, A. L., Findeisen, M., Hennig, L., Simonsen, O., Taylor, L. & Toftlund, H. L. (1997). *J. Chem. Soc. Dalton Trans.* pp. 111–120.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cohen, M. D. & Schmidt, G. M. J. (1964). *J. Chem. Soc.* pp. 2041–2051.
- Corden, J. P., Bishop, P. R., Errington, W. & Wallbridge, M. G. H. (1996). *Acta Cryst. C52*, 2777–2779.
- Fun, H.-K., Kia, R. & Raithby, P. R. (2008). *Acta Cryst. E64*, o1271–o1272.
- Govindasamy, L., Velmurugan, D. & Rajendran, T. M. (1999). *Acta Cryst. C55*, 1368–1369.
- Granovski, A. D., Nivorozhkin, A. L. & Minkin, V. I. (1993). *Coord. Chem. Rev.* **126**, 1–69.

organic compounds

- Li, C. H. & Chang, T. C. (1991). *Eur. Polym. J.* **27**, 35–39.
Shahrokhan, S., Amini, M. K., Kia, R. & Tangestaninejad, S. (2000). *Anal. Chem.* **72**, 956–962.
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
Unaleroğlu, C. & Hökelek, T. (2002). *Spectrosc. Lett.* **32**, 317–326.

supporting information

Acta Cryst. (2008). E64, o1411–o1412 [doi:10.1107/S1600536808020023]

2-[*(E*)-(5-Amino-2,3-diphenylquinoxalin-6-yl)iminomethyl]-4-chlorophenol

Hoong-Kun Fun, Reza Kia and Paul R. Raithby

S1. Comment

Schiff bases are one of the most prevalent mixed-donor ligand types in coordination chemistry. The biologically activities of their complexes have been studied extensively over the last decades (Anderson *et al.*, 1997; Corden *et al.*, 1996; Govindasamy *et al.*, 1999; Granovski *et al.*, 1993; Li & Chang, 1991; Shahrokhan *et al.*, 2000). 2-Hydroxy Schiff base ligands are of additional interest mainly due to the existence of O—H···N and O···H—N type hydrogen bonds and tautomerization between the phenol-imine and keto-amine forms (Unaleroğlu & Hokelek, 2002; Fun *et al.*, 2008). This type of tautomerism plays an important role for distinguishing their photochromic and thermochromic properties (Cohen & Schmidt, 1964). Knowing the solution and solid-state structures of free Schiff bases is important in view of the intramolecular hydrogen bonding and comparing conformation with that in the structures of Schiff base complexes. In view of the importance of these organic ligands, the title compound (I) was synthesized and its crystal structure is reported herein.


Compound (I, Fig. 1), features intramolecular O—H···N and N—H···N hydrogen bonds to form six- and five-membered rings, producing S(6) and S(5) ring motifs, respectively (Bernstein *et al.*, 1995). The two phenyl substituents on the quinoxaline unit are inclined at an angle of 61.14 (9)° to one another. They also form dihedral angles of 43.38 (9) and 39.50 (9)°, respectively, with the ten-membered quinoxaline ring. In the crystal packing (Table 1 & Fig. 2), molecules are stacked when viewed down the *b* axis, being consolidated by π – π interactions with $Cg_2 \cdots Cg_3$ distances ranging from 3.6294 (11) – 3.6881 (11) Å; symmetry codes 1 - *x*, -1/2 + *y*, 3/2 - *z* and 1 - *x*, 1/2 + *y*, 3/2 - *z*, and Cg_2 and Cg_3 are the centroids of the C1–C6 and C8/C9/C10/C11/C14/C15 phenyl rings, respectively. The crystal structure is also stabilized by intramolecular O—H···N and N—H···N contacts.

S2. Experimental


The synthetic method used for the preparation of (I) has been described earlier (Fun *et al.*, 2008). Single crystals suitable for X-ray diffraction were obtained by evaporation of a mixed dichloromethane-ethanol (3/1) solution of (I), held at room temperature.

S3. Refinement

The H-atoms attached to O1 and N4 were located in a difference Fourier map and refined freely; see Table 1 for bond distances. The remaining H atoms were included in the riding model approximation with C—H = 0.93 Å, and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

**Figure 1**

The molecular structure of (I), showing 50% probability displacement ellipsoids and atomic numbering. Intramolecular interactions are drawn as dashed lines.

**Figure 2**

The crystal packing in (I), viewed down the *b*-axis showing stacking arrangement. Intramolecular and intermolecular interactions are shown as dashed lines.

2-[(*E*)-(5-Amino-2,3-diphenylquinoxalin-6-yl)iminomethyl]-4-chlorophenol

Crystal data

$C_{27}H_{19}ClN_4O$
 $M_r = 450.91$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 22.8728 (11)$ Å
 $b = 7.3068 (4)$ Å
 $c = 12.5632 (6)$ Å
 $\beta = 92.037 (2)^\circ$
 $V = 2098.32 (18)$ Å³
 $Z = 4$

$F(000) = 936$
 $D_x = 1.427$ Mg m⁻³
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 2969 reflections
 $\theta = 2.9\text{--}28.1^\circ$
 $\mu = 0.21$ mm⁻¹
 $T = 100$ K
Block, yellow
 $0.55 \times 0.09 \times 0.07$ mm

Data collection

Bruker SMART APEXII CCD area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2005)
 $T_{\min} = 0.843$, $T_{\max} = 0.985$

25593 measured reflections
6180 independent reflections
4118 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.062$
 $\theta_{\max} = 30.1^\circ$, $\theta_{\min} = 0.9^\circ$
 $h = -32 \rightarrow 31$
 $k = -10 \rightarrow 10$
 $l = -17 \rightarrow 17$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.052$$

$$wR(F^2) = 0.159$$

$$S = 1.01$$

6180 reflections

310 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sitesH atoms treated by a mixture of independent
and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0879P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.40 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.41 \text{ e } \text{\AA}^{-3}$$

*Special details***Experimental.** The low-temperature data was collected with the Oxford Cyrosystem Cobra low-temperature attachment.**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.**Refinement.** Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.26328 (2)	0.46708 (7)	0.59559 (4)	0.02813 (14)
O1	0.44768 (6)	0.2687 (2)	0.91204 (10)	0.0252 (3)
N1	0.52358 (6)	0.3641 (2)	0.77495 (12)	0.0200 (3)
N2	0.72297 (7)	0.4713 (2)	0.88154 (11)	0.0174 (3)
N3	0.75971 (6)	0.3333 (2)	0.68619 (11)	0.0185 (3)
N4	0.60625 (8)	0.4640 (2)	0.92698 (13)	0.0263 (4)
C1	0.40599 (8)	0.3116 (3)	0.83649 (14)	0.0203 (4)
C2	0.34744 (8)	0.2879 (3)	0.85952 (15)	0.0236 (4)
H2A	0.3375	0.2402	0.9251	0.028*
C3	0.30418 (8)	0.3350 (3)	0.78542 (15)	0.0233 (4)
H3A	0.2651	0.3199	0.8013	0.028*
C4	0.31874 (8)	0.4049 (3)	0.68704 (15)	0.0214 (4)
C5	0.37624 (8)	0.4249 (3)	0.66116 (14)	0.0211 (4)
H5A	0.3855	0.4700	0.5946	0.025*
C6	0.42106 (8)	0.3773 (2)	0.73536 (14)	0.0197 (4)
C7	0.48149 (8)	0.3948 (3)	0.70633 (14)	0.0206 (4)
H7A	0.4901	0.4287	0.6373	0.025*
C8	0.58278 (7)	0.3677 (2)	0.74723 (14)	0.0183 (3)
C9	0.60160 (8)	0.3128 (3)	0.64598 (14)	0.0206 (4)
H9A	0.5739	0.2813	0.5932	0.025*
C10	0.65962 (8)	0.3050 (3)	0.62430 (14)	0.0205 (4)
H10A	0.6713	0.2684	0.5574	0.025*
C11	0.70168 (7)	0.3530 (2)	0.70445 (13)	0.0179 (3)

C12	0.79775 (7)	0.3790 (2)	0.76358 (13)	0.0175 (3)
C13	0.77919 (7)	0.4620 (2)	0.86047 (13)	0.0169 (3)
C14	0.68362 (7)	0.4111 (2)	0.80547 (13)	0.0169 (3)
C15	0.62300 (8)	0.4161 (2)	0.82744 (13)	0.0173 (3)
C16	0.85933 (7)	0.3244 (2)	0.74618 (13)	0.0186 (4)
C17	0.88232 (8)	0.3426 (3)	0.64549 (14)	0.0211 (4)
H17A	0.8603	0.3993	0.5913	0.025*
C18	0.93766 (8)	0.2771 (3)	0.62536 (14)	0.0232 (4)
H18A	0.9529	0.2925	0.5583	0.028*
C19	0.97043 (8)	0.1884 (3)	0.70519 (15)	0.0241 (4)
H19A	1.0072	0.1419	0.6912	0.029*
C20	0.94797 (8)	0.1695 (3)	0.80613 (14)	0.0221 (4)
H20A	0.9698	0.1112	0.8600	0.027*
C21	0.89277 (8)	0.2380 (3)	0.82614 (13)	0.0217 (4)
H21A	0.8780	0.2260	0.8938	0.026*
C22	0.82037 (8)	0.5458 (2)	0.93956 (13)	0.0181 (4)
C23	0.86747 (8)	0.6521 (3)	0.90850 (14)	0.0218 (4)
H23A	0.8743	0.6675	0.8365	0.026*
C24	0.90422 (8)	0.7350 (3)	0.98433 (15)	0.0250 (4)
H24A	0.9350	0.8081	0.9630	0.030*
C25	0.89514 (8)	0.7091 (3)	1.09180 (15)	0.0244 (4)
H25A	0.9205	0.7617	1.1426	0.029*
C26	0.84845 (8)	0.6051 (3)	1.12332 (14)	0.0227 (4)
H26A	0.8422	0.5887	1.1954	0.027*
C27	0.81078 (8)	0.5246 (2)	1.04769 (13)	0.0194 (4)
H27A	0.7790	0.4563	1.0694	0.023*
H1O1	0.4828 (12)	0.299 (4)	0.881 (2)	0.062 (9)*
H1N4	0.5681 (12)	0.531 (4)	0.9337 (19)	0.052 (8)*
H2N4	0.6396 (13)	0.516 (4)	0.976 (2)	0.060 (8)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.0190 (2)	0.0284 (3)	0.0367 (3)	0.0003 (2)	-0.00334 (18)	-0.0011 (2)
O1	0.0226 (7)	0.0291 (8)	0.0241 (6)	0.0010 (6)	0.0023 (5)	0.0007 (6)
N1	0.0176 (7)	0.0168 (7)	0.0256 (8)	-0.0017 (6)	0.0014 (6)	-0.0025 (6)
N2	0.0188 (7)	0.0151 (7)	0.0184 (7)	0.0010 (6)	0.0009 (5)	0.0010 (6)
N3	0.0183 (7)	0.0180 (7)	0.0194 (7)	0.0015 (6)	0.0001 (5)	0.0003 (6)
N4	0.0242 (8)	0.0304 (9)	0.0242 (8)	0.0028 (8)	0.0009 (6)	-0.0002 (7)
C1	0.0223 (9)	0.0162 (8)	0.0223 (8)	-0.0010 (7)	0.0013 (7)	-0.0036 (7)
C2	0.0245 (9)	0.0206 (9)	0.0262 (9)	-0.0020 (8)	0.0060 (7)	-0.0033 (7)
C3	0.0182 (8)	0.0210 (9)	0.0308 (9)	-0.0009 (8)	0.0039 (7)	-0.0060 (8)
C4	0.0178 (8)	0.0161 (8)	0.0302 (9)	0.0003 (7)	-0.0017 (7)	-0.0042 (7)
C5	0.0212 (9)	0.0173 (9)	0.0251 (9)	-0.0020 (8)	0.0022 (7)	-0.0012 (7)
C6	0.0191 (8)	0.0158 (8)	0.0244 (9)	-0.0004 (7)	0.0017 (7)	-0.0026 (7)
C7	0.0212 (9)	0.0174 (9)	0.0234 (8)	-0.0006 (8)	0.0027 (7)	-0.0012 (7)
C8	0.0170 (8)	0.0142 (8)	0.0237 (8)	-0.0002 (7)	0.0021 (6)	0.0005 (7)
C9	0.0196 (8)	0.0191 (9)	0.0230 (8)	-0.0005 (7)	-0.0012 (7)	-0.0031 (7)

C10	0.0225 (9)	0.0197 (9)	0.0193 (8)	0.0021 (8)	0.0002 (7)	-0.0018 (7)
C11	0.0178 (8)	0.0164 (8)	0.0194 (8)	0.0004 (7)	0.0007 (6)	-0.0009 (7)
C12	0.0181 (8)	0.0164 (8)	0.0179 (8)	0.0002 (7)	0.0022 (6)	0.0024 (6)
C13	0.0168 (8)	0.0161 (8)	0.0179 (8)	0.0007 (7)	0.0011 (6)	0.0019 (6)
C14	0.0176 (8)	0.0142 (8)	0.0190 (8)	0.0011 (7)	0.0014 (6)	0.0012 (6)
C15	0.0185 (8)	0.0135 (8)	0.0199 (8)	0.0017 (7)	0.0024 (6)	0.0016 (7)
C16	0.0164 (8)	0.0178 (8)	0.0218 (8)	-0.0005 (7)	0.0009 (6)	-0.0016 (7)
C17	0.0198 (8)	0.0230 (9)	0.0206 (8)	-0.0022 (8)	0.0006 (7)	0.0013 (7)
C18	0.0200 (9)	0.0296 (10)	0.0202 (8)	-0.0031 (8)	0.0032 (7)	-0.0018 (7)
C19	0.0154 (8)	0.0278 (10)	0.0292 (9)	0.0005 (8)	0.0034 (7)	-0.0014 (8)
C20	0.0178 (8)	0.0251 (10)	0.0234 (9)	0.0030 (8)	-0.0014 (7)	0.0012 (8)
C21	0.0212 (9)	0.0260 (10)	0.0182 (8)	0.0013 (8)	0.0030 (7)	-0.0001 (7)
C22	0.0171 (8)	0.0165 (8)	0.0204 (8)	0.0032 (7)	-0.0008 (6)	-0.0015 (7)
C23	0.0205 (8)	0.0220 (9)	0.0231 (8)	0.0011 (8)	0.0032 (7)	-0.0005 (7)
C24	0.0200 (9)	0.0237 (10)	0.0314 (10)	-0.0031 (8)	0.0035 (7)	-0.0025 (8)
C25	0.0198 (9)	0.0250 (10)	0.0280 (9)	0.0008 (8)	-0.0040 (7)	-0.0052 (8)
C26	0.0254 (9)	0.0224 (9)	0.0202 (8)	0.0025 (8)	-0.0003 (7)	-0.0006 (7)
C27	0.0170 (8)	0.0202 (9)	0.0210 (8)	-0.0001 (7)	0.0023 (6)	-0.0011 (7)

Geometric parameters (\AA , $^\circ$)

C11—C4	1.7414 (19)	C10—H10A	0.9300
O1—C1	1.358 (2)	C11—C14	1.414 (2)
O1—H1O1	0.93 (3)	C12—C13	1.437 (2)
N1—C7	1.289 (2)	C12—C16	1.488 (2)
N1—C8	1.410 (2)	C13—C22	1.478 (2)
N2—C13	1.324 (2)	C14—C15	1.424 (2)
N2—C14	1.362 (2)	C16—C21	1.392 (2)
N3—C12	1.324 (2)	C16—C17	1.393 (2)
N3—C11	1.363 (2)	C17—C18	1.385 (2)
N4—C15	1.366 (2)	C17—H17A	0.9300
N4—H1N4	1.01 (3)	C18—C19	1.391 (3)
N4—H2N4	1.03 (3)	C18—H18A	0.9300
C1—C2	1.391 (3)	C19—C20	1.392 (3)
C1—C6	1.412 (2)	C19—H19A	0.9300
C2—C3	1.378 (3)	C20—C21	1.389 (2)
C2—H2A	0.9300	C20—H20A	0.9300
C3—C4	1.389 (3)	C21—H21A	0.9300
C3—H3A	0.9300	C22—C27	1.392 (2)
C4—C5	1.374 (2)	C22—C23	1.395 (2)
C5—C6	1.405 (2)	C23—C24	1.387 (3)
C5—H5A	0.9300	C23—H23A	0.9300
C6—C7	1.448 (2)	C24—C25	1.386 (3)
C7—H7A	0.9300	C24—H24A	0.9300
C8—C15	1.386 (2)	C25—C26	1.380 (3)
C8—C9	1.415 (2)	C25—H25A	0.9300
C9—C10	1.365 (2)	C26—C27	1.390 (2)
C9—H9A	0.9300	C26—H26A	0.9300

C10—C11	1.412 (2)	C27—H27A	0.9300
C1—O1—H1O1	104.0 (16)	N2—C13—C22	116.42 (15)
C7—N1—C8	122.17 (15)	C12—C13—C22	122.95 (15)
C13—N2—C14	117.74 (14)	N2—C14—C11	121.32 (15)
C12—N3—C11	117.91 (15)	N2—C14—C15	118.65 (15)
C15—N4—H1N4	118.3 (14)	C11—C14—C15	119.96 (16)
C15—N4—H2N4	114.3 (16)	N4—C15—C8	122.02 (16)
H1N4—N4—H2N4	113 (2)	N4—C15—C14	119.38 (16)
O1—C1—C2	118.90 (16)	C8—C15—C14	118.58 (15)
O1—C1—C6	121.29 (16)	C21—C16—C17	118.89 (16)
C2—C1—C6	119.81 (17)	C21—C16—C12	120.97 (15)
C3—C2—C1	120.12 (17)	C17—C16—C12	119.83 (15)
C3—C2—H2A	119.9	C18—C17—C16	120.63 (17)
C1—C2—H2A	119.9	C18—C17—H17A	119.7
C2—C3—C4	120.27 (17)	C16—C17—H17A	119.7
C2—C3—H3A	119.9	C17—C18—C19	120.12 (16)
C4—C3—H3A	119.9	C17—C18—H18A	119.9
C5—C4—C3	120.78 (17)	C19—C18—H18A	119.9
C5—C4—C11	119.81 (14)	C18—C19—C20	119.77 (17)
C3—C4—C11	119.40 (14)	C18—C19—H19A	120.1
C4—C5—C6	119.91 (17)	C20—C19—H19A	120.1
C4—C5—H5A	120.0	C21—C20—C19	119.75 (17)
C6—C5—H5A	120.0	C21—C20—H20A	120.1
C5—C6—C1	119.04 (16)	C19—C20—H20A	120.1
C5—C6—C7	119.45 (16)	C20—C21—C16	120.81 (16)
C1—C6—C7	121.51 (16)	C20—C21—H21A	119.6
N1—C7—C6	120.88 (16)	C16—C21—H21A	119.6
N1—C7—H7A	119.6	C27—C22—C23	119.01 (16)
C6—C7—H7A	119.6	C27—C22—C13	119.40 (16)
C15—C8—N1	116.33 (15)	C23—C22—C13	121.53 (15)
C15—C8—C9	120.70 (16)	C24—C23—C22	120.41 (16)
N1—C8—C9	122.82 (15)	C24—C23—H23A	119.8
C10—C9—C8	121.32 (16)	C22—C23—H23A	119.8
C10—C9—H9A	119.3	C25—C24—C23	120.07 (18)
C8—C9—H9A	119.3	C25—C24—H24A	120.0
C9—C10—C11	119.34 (16)	C23—C24—H24A	120.0
C9—C10—H10A	120.3	C26—C25—C24	119.92 (17)
C11—C10—H10A	120.3	C26—C25—H25A	120.0
N3—C11—C10	119.78 (15)	C24—C25—H25A	120.0
N3—C11—C14	120.02 (15)	C25—C26—C27	120.26 (16)
C10—C11—C14	120.09 (16)	C25—C26—H26A	119.9
N3—C12—C13	121.47 (15)	C27—C26—H26A	119.9
N3—C12—C16	115.16 (15)	C26—C27—C22	120.28 (17)
C13—C12—C16	123.18 (15)	C26—C27—H27A	119.9
N2—C13—C12	120.61 (15)	C22—C27—H27A	119.9
O1—C1—C2—C3	178.24 (16)	C10—C11—C14—N2	-175.02 (16)

C6—C1—C2—C3	-2.5 (3)	N3—C11—C14—C15	-174.42 (16)
C1—C2—C3—C4	0.5 (3)	C10—C11—C14—C15	1.8 (3)
C2—C3—C4—C5	1.3 (3)	N1—C8—C15—N4	-2.2 (3)
C2—C3—C4—C11	-178.91 (14)	C9—C8—C15—N4	-177.84 (17)
C3—C4—C5—C6	-1.1 (3)	N1—C8—C15—C14	176.25 (16)
C11—C4—C5—C6	179.12 (14)	C9—C8—C15—C14	0.6 (3)
C4—C5—C6—C1	-0.9 (3)	N2—C14—C15—N4	-6.2 (3)
C4—C5—C6—C7	178.53 (17)	C11—C14—C15—N4	176.85 (17)
O1—C1—C6—C5	-178.08 (16)	N2—C14—C15—C8	175.28 (16)
C2—C1—C6—C5	2.7 (3)	C11—C14—C15—C8	-1.6 (3)
O1—C1—C6—C7	2.5 (3)	N3—C12—C16—C21	-132.71 (18)
C2—C1—C6—C7	-176.75 (17)	C13—C12—C16—C21	42.4 (3)
C8—N1—C7—C6	175.52 (16)	N3—C12—C16—C17	40.8 (2)
C5—C6—C7—N1	175.24 (17)	C13—C12—C16—C17	-144.14 (18)
C1—C6—C7—N1	-5.3 (3)	C21—C16—C17—C18	-0.4 (3)
C7—N1—C8—C15	151.46 (17)	C12—C16—C17—C18	-174.06 (17)
C7—N1—C8—C9	-33.0 (3)	C16—C17—C18—C19	1.4 (3)
C15—C8—C9—C10	0.3 (3)	C17—C18—C19—C20	-1.5 (3)
N1—C8—C9—C10	-175.08 (17)	C18—C19—C20—C21	0.5 (3)
C8—C9—C10—C11	-0.1 (3)	C19—C20—C21—C16	0.5 (3)
C12—N3—C11—C10	-179.68 (16)	C17—C16—C21—C20	-0.6 (3)
C12—N3—C11—C14	-3.4 (3)	C12—C16—C21—C20	173.01 (17)
C9—C10—C11—N3	175.32 (17)	N2—C13—C22—C27	40.2 (2)
C9—C10—C11—C14	-0.9 (3)	C12—C13—C22—C27	-141.69 (18)
C11—N3—C12—C13	-5.4 (3)	N2—C13—C22—C23	-137.21 (17)
C11—N3—C12—C16	169.78 (15)	C12—C13—C22—C23	40.9 (3)
C14—N2—C13—C12	-4.3 (2)	C27—C22—C23—C24	0.1 (3)
C14—N2—C13—C22	173.81 (15)	C13—C22—C23—C24	177.55 (17)
N3—C12—C13—N2	9.7 (3)	C22—C23—C24—C25	1.5 (3)
C16—C12—C13—N2	-165.05 (17)	C23—C24—C25—C26	-1.9 (3)
N3—C12—C13—C22	-168.31 (16)	C24—C25—C26—C27	0.7 (3)
C16—C12—C13—C22	16.9 (3)	C25—C26—C27—C22	1.0 (3)
C13—N2—C14—C11	-4.5 (3)	C23—C22—C27—C26	-1.4 (3)
C13—N2—C14—C15	178.63 (16)	C13—C22—C27—C26	-178.89 (17)
N3—C11—C14—N2	8.7 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

D—H···A	D—H	H···A	D···A	D—H···A
O1—H1O1···N1	0.93 (3)	1.72 (3)	2.584 (2)	153 (2)
N4—H1N4···O1 ⁱ	1.01 (3)	2.47 (3)	3.099 (2)	120.4 (18)
N4—H2N4···N2	1.04 (3)	2.31 (3)	2.750 (2)	104.3 (17)
C27—H27A···N3 ⁱⁱ	0.93	2.62	3.373 (2)	138

Symmetry codes: (i) $-x+1, -y+1, -z+2$; (ii) $x, -y+1/2, z+1/2$.