

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Tetrakis(1,3,4,6,7,9-hexaaza-1*H*-phenalen-6-ium) sodium(I) pentakis(tetrafluoridoborate)

Yuqin Jiang, Yanhui Hou and Yongsheng Chen*

Key Laboratory for Functional Polymer Materials and Center for Nanoscale Science and Technology, Institute of Polymer Chemistry, College of Chemistry, Nankai University, Tianjin, Weijin Road No. 94, Tianjin, People's Republic of China
Correspondence e-mail: jiangyuqin800520@mail.nankai.edu.cn


Received 1 September 2008; accepted 20 September 2008

Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.055; wR factor = 0.152; data-to-parameter ratio = 13.7.

In the title compound, $\text{Na}^+ \cdot 4\text{C}_7\text{H}_5\text{N}_6^+ \cdot 5\text{BF}_4^-$, the Na^+ ion lies on a fourfold rotation axis and one of the tetrafluoridoborate ions lies on a site of symmetry $\bar{4}$. Each Na^+ ion is surrounded by four symmetry-related tetrafluoridoborate ions, and is eight-coordinated by F atoms, the $\text{Na} \cdots \text{F}$ separation being 2.3956 (15) or 2.4347 (17) Å. The hexaazaphenalenium ring system is essentially planar. In the crystal structure, the cations and anions are linked into a three-dimensional network by $\text{N}-\text{H} \cdots \text{N}$ and $\text{C}-\text{H} \cdots \text{F}$ hydrogen bonds.

Related literature

For general background, see: Goto *et al.* (1999); Haddon (1975); Koutentis *et al.* (2001). For related structures, see: Morita *et al.* (2002, 2003); Tamaki *et al.* (1997); Zheng *et al.* (2003, 2005). For related preparation, see: Suzuki *et al.* (2005).



Experimental

Crystal data

$\text{Na}^+ \cdot 4\text{C}_7\text{H}_5\text{N}_6^+ \cdot 5\text{BF}_4^-$
 $M_r = 1149.72$
Tetragonal, $P4/n$
 $a = 15.0665$ (6) Å
 $c = 8.9322$ (5) Å
 $V = 2027.60$ (16) Å³

$Z = 2$
Mo $K\alpha$ radiation
 $\mu = 0.20$ mm⁻¹
 $T = 113$ (2) K
 $0.14 \times 0.12 \times 0.12$ mm

Data collection

Rigaku Saturn CCD area-detector diffractometer
Absorption correction: multi-scan (*CrystalClear*, Rigaku/MSC, 2005)
 $T_{\min} = 0.877$, $T_{\max} = 0.977$
21698 measured reflections
2423 independent reflections
2247 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.044$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.152$
 $S = 1.19$
2423 reflections
177 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.74$ e Å⁻³
 $\Delta\rho_{\min} = -0.79$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N3}-\text{H3N} \cdots \text{N4}^{\text{i}}$	0.88	2.05	2.930 (2)	176
$\text{N6}-\text{H6N} \cdots \text{N5}^{\text{ii}}$	0.88	2.09	2.866 (2)	146
$\text{C1}-\text{H1} \cdots \text{F3}^{\text{iii}}$	0.95	2.47	3.179 (2)	132
$\text{C1}-\text{H1} \cdots \text{F4}^{\text{ii}}$	0.95	2.55	2.940 (3)	105
$\text{C3}-\text{H3} \cdots \text{F2}^{\text{i}}$	0.95	2.22	3.124 (2)	159
$\text{C5}-\text{H5} \cdots \text{F3}^{\text{iv}}$	0.95	2.40	3.097 (2)	130

Symmetry codes: (i) $-x + 1, -y + 1, -z + 2$; (ii) $-y + 1, x + \frac{1}{2}, -z + 1$; (iii) $y, -x + \frac{1}{2}, z$; (iv) $-y + \frac{1}{2}, x, z$.

Data collection: *CrystalClear* (Rigaku/MSC, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *CrystalStructure* (Rigaku/MSC, 2005).

This work was supported by the National Natural Science Foundation of China (NSFC; grant Nos. 20644004 and 20774047), the Ministry of Science and Technology of China (MOST; grant No. 2006CB932702) and the NSF of Tianjin City (grant No. 07JCYBJC03000).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2666).

References

- Goto, K., Kubo, T., Yamamoto, K., Nakasuji, K., Sato, K., Shiomi, D., Takui, T., Kubota, M., Kobayashi, T., Yakushi, K. & Ouyang, J. (1999). *J. Am. Chem. Soc.* **121**, 1619–1620.
Haddon, R. C. (1975). *Nature (London)*, **256**, 394–396.
Koutentis, P. A., Chen, Y., Cao, Y., Best, T. P., Itkis, M. E., Beer, L., Oakley, R. T., Cordes, A. M., Brock, C. P. & Haddon, R. C. (2001). *J. Am. Chem. Soc.* **123**, 3864–3871.
Morita, Y., Aoki, T., Fukui, K., Nakazawa, S., Tamaki, K., Suzuki, S., Fuyuhiro, A., Yamamoto, K., Sato, K., Shiomi, D., Naito, A., Takui, T. & Nakasuji, K. (2002). *Angew. Chem. Int. Ed.* **41**, 1793–1796.
Morita, Y., Suzuki, S., Fukui, K., Nakazawa, S., Sato, K., Shiomi, D., Takui, T. & Nakasuji, K. (2003). *Polyhedron*, **22**, 2215–2218.
Rigaku/MSC (2005). *CrystalClear* and *CrystalStructure*. Rigaku/MSC, The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Suzuki, S., Morita, Y., Fukui, K., Sato, K., Shiomi, D., Takui, T. & Nakasuji, K. (2005). *Inorg. Chem.* **44**, 8197–8199.
Tamaki, K., Morita, Y., Toyoda, J., Yamochi, H., Saito, G. & Nakasuji, K. (1997). *Tetrahedron Lett.* **38**, 4583–4586.
Zheng, S., Lan, J., Kahn, S. I. & Rubin, Y. (2003). *J. Am. Chem. Soc.* **125**, 5786–5791.
Zheng, S., Thompson, J. D., Tontcheva, A., Khan, S. I. & Rubin, Y. (2005). *Org. Lett.* **7**, 1861–1863.

supporting information

Acta Cryst. (2008). E64, m1318 [doi:10.1107/S1600536808030286]

Tetrakis(1,3,4,6,7,9-hexaaza-1*H*-phenalen-6-ium) sodium(I) pentakis-(tetrafluoridoborate)

Yuqin Jiang, Yanhui Hou and Yongsheng Chen

S1. Comment

Phenalenyl, as a planar π system with D_{3h} symmetry, has received much attention and still played an important role as a building block for spin-mediated molecular functional materials (Haddon 1975; Koutentis *et al.*, 2001; Goto *et al.*, 1999). Hexaazaphenalene (HAP) is a highly symmetric heterocycle with full nitrogen substitution in all of the α sites of phenalene (Morita *et al.*, 2002; Morita *et al.*, 2003; Suzuki *et al.*, 2005; Tamaki *et al.*, 1997; Zheng *et al.*, 2003; Zheng *et al.*, 2005). Because of the directionality of lone-pair electrons at the nitrogen sites, incorporation of nitrogen atoms into the phenalenyl skeleton was found to give substantial effects on its electronic structure. Suzuki *et al.* have reported the preparation and single crystal structure of the anion of hexaazaphenalene. Herein, we report the preparation and crystal structure of a hexaazaphenalene fluoroboric salt, the title compound.

In the title compound, the Na^+ ion lies on a four-fold rotation axis and one of the tetrafluoridoborate ions lies on a $\bar{4}$ site symmetry. One of the remaining four BF_4^- ions and one of the four hexaazaphenalenium ions are symmetry independent, and they lie on general positions. Each Na^+ ion is surrounded by four symmetry related tetrafluoridoborate ions, with the $\text{Na}\cdots\text{F}$ separation being 2.3956 (15) or 2.4347 (17) Å (Fig.1). The hexaazaphenalenium ring system is essentially planar, with a maximum deviation of 0.042 (2) Å for atom N6. The C—N distances lie in the range 1.310 (3)-1.363 (3) Å.


Two types of hydrogen bonds viz. $\text{N—H}\cdots\text{N}$ and $\text{C—H}\cdots\text{F}$ exist in the solid state of the title compound. These hydrogen bonds link the cationic and anionic units into a three-dimensional network (Fig. 2). There are no face to face π - π staking involving the hexaazaphenalenium units, which is a common packing mode in other phenalene compounds.

S2. Experimental


The sodium salt of hexaazaphenalene was prepared according to the literature method (Suzuki *et al.*, 2005). To a water (5 ml) solution of sodium salt of hexaazaphenalene (97 mg, 0.5 mmol) was added 40% HBF_4 (0.5 ml, 3.0 mmol). The reaction mixture was evaporated slowly for 7 d to afford colourless crystals of the title compound.

S3. Refinement

H atoms were initially located in difference maps and then refined using the riding-model approximation, with C-H = 0.95 Å, N-H = 0.88 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C},\text{N})$.

**Figure 1**

A view of the environment around the Na⁺ ion in the title compound. Displacement ellipsoids are drawn at the 50% probability level. Atoms labelled with the suffixes A, B, C, D, E and F are generated by the symmetry operations $(y, 1/2-x, z)$, $(1/2-x, 1/2-y, z)$, $(1/2-y, x, z)$, $(1-y, x-1/2, -z)$, $(y+1/2, 1-x, -z)$ and $(3/2-x, 1/2-y, z)$, respectively.

**Figure 2**

The crystal packing of the title compound.

Tetrakis(1,3,4,6,7,9-hexaaza-1H-phenalen-6-ium) sodium(I) pentakis(tetrafluoridoborate)

Crystal data

$\text{Na}^+ \cdot 4\text{C}_7\text{H}_5\text{N}_6^+ \cdot 5\text{BF}_4^-$

$M_r = 1149.72$

Tetragonal, $P4/n$

Hall symbol: $-P\ 4a$

$a = 15.0665\ (6)\ \text{\AA}$

$c = 8.9322\ (5)\ \text{\AA}$

$V = 2027.60\ (16)\ \text{\AA}^3$

$Z = 2$

$F(000) = 1144$

$D_x = 1.883\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71070\ \text{\AA}$

Cell parameters from 4230 reflections

$\theta = 1.9\text{--}27.9^\circ$

$\mu = 0.20\ \text{mm}^{-1}$

$T = 113\ \text{K}$

Block, colourless

$0.14 \times 0.12 \times 0.12\ \text{mm}$

Data collection

Rigaku Saturn CCD area-detector
diffractometer

Radiation source: rotating anode

Confocal monochromator

Detector resolution: $7.31\ \text{pixels mm}^{-1}$

ω and φ scans

Absorption correction: multi-scan

(*CrystalClear*, Rigaku/MSC, 2005)

$T_{\min} = 0.878$, $T_{\max} = 0.977$

21698 measured reflections

2423 independent reflections

2247 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.044$

$\theta_{\max} = 27.9^\circ$, $\theta_{\min} = 1.9^\circ$

$h = -19 \rightarrow 19$

$k = -19 \rightarrow 19$

$l = -11 \rightarrow 11$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.055$
 $wR(F^2) = 0.152$
 $S = 1.19$
 2423 reflections
 177 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0759P)^2 + 1.4095P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.74 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.79 \text{ e } \text{Å}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Na1	0.2500	0.2500	0.4845 (2)	0.0275 (4)
B1	0.42320 (16)	0.33819 (16)	0.5211 (3)	0.0204 (5)
F1	0.39625 (9)	0.28695 (10)	0.39963 (15)	0.0293 (4)
F2	0.35132 (8)	0.34700 (9)	0.61981 (15)	0.0241 (3)
F3	0.49388 (9)	0.29764 (9)	0.59536 (16)	0.0274 (3)
F4	0.44983 (10)	0.42159 (9)	0.47314 (19)	0.0346 (4)
F5	0.73061 (11)	0.17735 (10)	0.08990 (17)	0.0371 (4)
B2	0.7500	0.2500	0.0000	0.0203 (9)
C1	0.43924 (15)	0.86438 (14)	0.7069 (2)	0.0202 (4)
H1	0.4317	0.9205	0.6602	0.024*
C2	0.50050 (13)	0.78212 (13)	0.8918 (2)	0.0150 (4)
C3	0.56501 (13)	0.69463 (13)	1.0679 (2)	0.0164 (4)
H3	0.6034	0.6886	1.1517	0.020*
C4	0.47115 (12)	0.62292 (12)	0.8956 (2)	0.0126 (4)
C5	0.38121 (13)	0.56354 (13)	0.7195 (2)	0.0157 (4)
H5	0.3522	0.5127	0.6798	0.019*
C6	0.40548 (13)	0.71345 (12)	0.7050 (2)	0.0135 (4)
C7	0.45850 (12)	0.70676 (12)	0.8328 (2)	0.0124 (4)
N1	0.48868 (12)	0.86262 (11)	0.82849 (19)	0.0188 (4)
N2	0.55508 (11)	0.77454 (11)	1.01274 (19)	0.0166 (4)
N3	0.52553 (11)	0.61957 (11)	1.01648 (18)	0.0138 (3)
H3N	0.5353	0.5686	1.0617	0.017*
N4	0.43264 (11)	0.54996 (11)	0.83918 (18)	0.0146 (4)
N5	0.36565 (11)	0.64098 (11)	0.64850 (19)	0.0155 (4)
N6	0.39805 (12)	0.79459 (11)	0.64176 (19)	0.0181 (4)

H6N 0.3669 0.8019 0.5594 0.022*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Na1	0.0214 (6)	0.0214 (6)	0.0396 (11)	0.000	0.000	0.000
B1	0.0199 (11)	0.0185 (11)	0.0227 (12)	-0.0013 (9)	0.0011 (9)	0.0025 (9)
F1	0.0278 (7)	0.0369 (8)	0.0231 (7)	-0.0037 (6)	0.0034 (5)	-0.0066 (6)
F2	0.0218 (7)	0.0290 (7)	0.0214 (7)	0.0028 (5)	-0.0002 (5)	-0.0023 (5)
F3	0.0219 (7)	0.0271 (7)	0.0332 (8)	0.0028 (5)	-0.0018 (5)	0.0047 (6)
F4	0.0324 (8)	0.0248 (7)	0.0466 (9)	-0.0042 (6)	-0.0024 (7)	0.0124 (6)
F5	0.0538 (10)	0.0295 (8)	0.0279 (8)	-0.0065 (7)	0.0028 (7)	0.0055 (6)
B2	0.0219 (14)	0.0219 (14)	0.017 (2)	0.000	0.000	0.000
C1	0.0289 (11)	0.0151 (9)	0.0167 (9)	0.0048 (8)	0.0014 (8)	0.0023 (8)
C2	0.0178 (9)	0.0139 (9)	0.0134 (9)	0.0013 (7)	0.0007 (7)	0.0003 (7)
C3	0.0188 (9)	0.0164 (9)	0.0139 (9)	0.0011 (7)	-0.0035 (7)	-0.0004 (7)
C4	0.0115 (8)	0.0136 (9)	0.0126 (8)	0.0013 (7)	0.0004 (7)	0.0005 (7)
C5	0.0131 (8)	0.0185 (9)	0.0155 (9)	-0.0013 (7)	-0.0005 (7)	-0.0013 (7)
C6	0.0159 (9)	0.0147 (9)	0.0099 (8)	0.0026 (7)	0.0003 (7)	-0.0005 (7)
C7	0.0141 (8)	0.0123 (9)	0.0110 (8)	0.0005 (7)	0.0009 (7)	0.0006 (7)
N1	0.0253 (9)	0.0131 (8)	0.0181 (8)	0.0011 (6)	-0.0028 (7)	0.0016 (6)
N2	0.0205 (9)	0.0139 (8)	0.0155 (8)	-0.0003 (6)	-0.0044 (6)	-0.0005 (6)
N3	0.0165 (8)	0.0120 (8)	0.0130 (8)	0.0010 (6)	-0.0029 (6)	0.0023 (6)
N4	0.0144 (8)	0.0153 (8)	0.0141 (8)	-0.0017 (6)	-0.0012 (6)	-0.0001 (6)
N5	0.0143 (8)	0.0177 (8)	0.0146 (8)	0.0006 (6)	-0.0016 (6)	-0.0017 (6)
N6	0.0249 (9)	0.0171 (8)	0.0124 (8)	0.0045 (7)	-0.0018 (7)	0.0033 (6)

Geometric parameters (Å, °)

Na1—F1 ⁱ	2.3956 (15)	C1—H1	0.95
Na1—F1 ⁱⁱ	2.3956 (15)	C2—N1	1.350 (3)
Na1—F1 ⁱⁱⁱ	2.3956 (15)	C2—N2	1.363 (3)
Na1—F1	2.3957 (15)	C2—C7	1.403 (3)
Na1—F2	2.4347 (17)	C3—N2	1.310 (3)
Na1—F2 ⁱ	2.4347 (17)	C3—N3	1.358 (3)
Na1—F2 ⁱⁱ	2.4347 (17)	C3—H3	0.95
Na1—F2 ⁱⁱⁱ	2.4347 (17)	C4—N4	1.341 (2)
B1—F4	1.387 (3)	C4—N3	1.356 (2)
B1—F1	1.392 (3)	C4—C7	1.395 (3)
B1—F3	1.395 (3)	C5—N4	1.336 (3)
B1—F2	1.403 (3)	C5—N5	1.349 (3)
F5—B2	1.3887 (14)	C5—H5	0.95
B2—F5 ^{iv}	1.3887 (14)	C6—N5	1.344 (3)
B2—F5 ^v	1.3887 (14)	C6—N6	1.351 (2)
B2—F5 ^{vi}	1.3887 (14)	C6—C7	1.397 (3)
C1—N1	1.317 (3)	N3—H3N	0.88
C1—N6	1.353 (3)	N6—H6N	0.88

F1 ⁱ —Na1—F1 ⁱⁱ	84.26 (3)	F1—Na1—B1 ⁱⁱⁱ	104.13 (6)
F1 ⁱ —Na1—F1 ⁱⁱⁱ	143.12 (11)	F2—Na1—B1 ⁱⁱⁱ	72.30 (6)
F1 ⁱⁱ —Na1—F1 ⁱⁱⁱ	84.26 (3)	F2 ⁱ —Na1—B1 ⁱⁱⁱ	140.43 (9)
F1 ⁱ —Na1—F1	84.26 (3)	F2 ⁱⁱ —Na1—B1 ⁱⁱⁱ	101.17 (7)
F1 ⁱⁱ —Na1—F1	143.12 (11)	F2 ⁱⁱⁱ —Na1—B1 ⁱⁱⁱ	28.22 (6)
F1 ⁱⁱⁱ —Na1—F1	84.26 (3)	B1—Na1—B1 ⁱⁱⁱ	89.293 (14)
F1 ⁱ —Na1—F2	124.30 (5)	B1 ⁱ —Na1—B1 ⁱⁱⁱ	167.24 (13)
F1 ⁱⁱ —Na1—F2	150.83 (6)	B1 ⁱⁱ —Na1—B1 ⁱⁱⁱ	89.292 (14)
F1 ⁱⁱⁱ —Na1—F2	75.56 (5)	F4—B1—F1	110.24 (19)
F1—Na1—F2	56.00 (4)	F4—B1—F3	108.83 (18)
F1 ⁱ —Na1—F2 ⁱ	56.00 (4)	F1—B1—F3	110.49 (19)
F1 ⁱⁱ —Na1—F2 ⁱ	124.30 (5)	F4—B1—F2	109.39 (19)
F1 ⁱⁱⁱ —Na1—F2 ⁱ	150.83 (6)	F1—B1—F2	108.49 (18)
F1—Na1—F2 ⁱ	75.56 (5)	F3—B1—F2	109.39 (18)
F2—Na1—F2 ⁱ	75.73 (5)	F4—B1—Na1	129.09 (15)
F1 ⁱ —Na1—F2 ⁱⁱ	75.56 (5)	F1—B1—Na1	53.49 (10)
F1 ⁱⁱ —Na1—F2 ⁱⁱ	56.00 (4)	F3—B1—Na1	122.09 (14)
F1 ⁱⁱⁱ —Na1—F2 ⁱⁱ	124.30 (5)	F2—B1—Na1	55.16 (10)
F1—Na1—F2 ⁱⁱ	150.84 (6)	B1—F1—Na1	98.66 (13)
F2—Na1—F2 ⁱⁱ	120.46 (11)	B1—F2—Na1	96.62 (12)
F2 ⁱ —Na1—F2 ⁱⁱ	75.73 (5)	F5—B2—F5 ^{iv}	109.54 (6)
F1 ⁱ —Na1—F2 ⁱⁱⁱ	150.84 (6)	F5—B2—F5 ^v	109.54 (6)
F1 ⁱⁱ —Na1—F2 ⁱⁱⁱ	75.56 (5)	F5 ^{iv} —B2—F5 ^v	109.34 (13)
F1 ⁱⁱⁱ —Na1—F2 ⁱⁱⁱ	56.00 (4)	F5—B2—F5 ^{vi}	109.34 (13)
F1—Na1—F2 ⁱⁱⁱ	124.29 (5)	F5 ^{iv} —B2—F5 ^{vi}	109.54 (6)
F2—Na1—F2 ⁱⁱⁱ	75.73 (5)	F5 ^v —B2—F5 ^{vi}	109.54 (6)
F2 ⁱ —Na1—F2 ⁱⁱⁱ	120.46 (11)	N1—C1—N6	126.77 (19)
F2 ⁱⁱ —Na1—F2 ⁱⁱⁱ	75.73 (5)	N1—C1—H1	116.6
F1 ⁱ —Na1—B1	104.13 (6)	N6—C1—H1	116.6
F1 ⁱⁱ —Na1—B1	162.64 (8)	N1—C2—N2	119.14 (18)
F1 ⁱⁱⁱ —Na1—B1	79.99 (6)	N1—C2—C7	120.70 (18)
F1—Na1—B1	27.85 (6)	N2—C2—C7	120.15 (18)
F2—Na1—B1	28.22 (6)	N2—C3—N3	126.03 (18)
F2 ⁱ —Na1—B1	72.30 (6)	N2—C3—H3	117.0
F2 ⁱⁱ —Na1—B1	140.43 (9)	N3—C3—H3	117.0
F2 ⁱⁱⁱ —Na1—B1	101.16 (7)	N4—C4—N3	122.01 (17)
F1 ⁱ —Na1—B1 ⁱ	27.85 (6)	N4—C4—C7	122.11 (17)
F1 ⁱⁱ —Na1—B1 ⁱ	104.13 (6)	N3—C4—C7	115.87 (17)
F1 ⁱⁱⁱ —Na1—B1 ⁱ	162.64 (8)	N4—C5—N5	127.57 (18)
F1—Na1—B1 ⁱ	79.99 (6)	N4—C5—H5	116.2
F2—Na1—B1 ⁱ	101.16 (7)	N5—C5—H5	116.2
F2 ⁱ —Na1—B1 ⁱ	28.22 (6)	N5—C6—N6	122.77 (18)
F2 ⁱⁱ —Na1—B1 ⁱ	72.30 (6)	N5—C6—C7	120.22 (17)
F2 ⁱⁱⁱ —Na1—B1 ⁱ	140.43 (9)	N6—C6—C7	117.00 (17)
B1—Na1—B1 ⁱ	89.292 (14)	C4—C7—C6	118.18 (17)
F1 ⁱ —Na1—B1 ⁱⁱ	79.99 (6)	C4—C7—C2	121.34 (18)
F1 ⁱⁱ —Na1—B1 ⁱⁱ	27.85 (6)	C6—C7—C2	120.43 (17)
F1 ⁱⁱⁱ —Na1—B1 ⁱⁱ	104.13 (6)	C1—N1—C2	115.98 (18)

F1—Na1—B1 ⁱⁱ	162.64 (8)	C3—N2—C2	116.39 (17)
F2—Na1—B1 ⁱⁱ	140.43 (9)	C4—N3—C3	120.21 (16)
F2 ⁱ —Na1—B1 ⁱⁱ	101.17 (7)	C4—N3—H3N	119.9
F2 ⁱⁱ —Na1—B1 ⁱⁱ	28.22 (6)	C3—N3—H3N	119.9
F2 ⁱⁱⁱ —Na1—B1 ⁱⁱ	72.30 (6)	C5—N4—C4	115.22 (17)
B1—Na1—B1 ⁱⁱ	167.24 (13)	C6—N5—C5	116.66 (17)
B1 ⁱ —Na1—B1 ⁱⁱ	89.293 (14)	C6—N6—C1	119.05 (18)
F1 ⁱ —Na1—B1 ⁱⁱⁱ	162.64 (8)	C6—N6—H6N	120.5
F1 ⁱⁱ —Na1—B1 ⁱⁱⁱ	79.99 (6)	C1—N6—H6N	120.5
F1 ⁱⁱⁱ —Na1—B1 ⁱⁱⁱ	27.85 (6)		
F1 ⁱ —Na1—B1—F4	133.1 (2)	F2 ⁱ —Na1—F1—B1	-79.36 (13)
F1 ⁱⁱ —Na1—B1—F4	15.7 (4)	F2 ⁱⁱ —Na1—F1—B1	-89.7 (2)
F1 ⁱⁱⁱ —Na1—B1—F4	-9.4 (2)	F2 ⁱⁱⁱ —Na1—F1—B1	37.90 (16)
F1—Na1—B1—F4	87.3 (2)	B1 ⁱ —Na1—F1—B1	-107.90 (14)
F2—Na1—B1—F4	-87.5 (2)	B1 ⁱⁱ —Na1—F1—B1	-160.5 (3)
F2 ⁱ —Na1—B1—F4	179.8 (2)	B1 ⁱⁱⁱ —Na1—F1—B1	59.75 (12)
F2 ⁱⁱ —Na1—B1—F4	-142.58 (17)	F4—B1—F2—Na1	124.71 (16)
F2 ⁱⁱⁱ —Na1—B1—F4	-61.5 (2)	F1—B1—F2—Na1	4.42 (17)
B1 ⁱ —Na1—B1—F4	156.90 (17)	F3—B1—F2—Na1	-116.18 (16)
B1 ⁱⁱ —Na1—B1—F4	-119.4 (2)	F1 ⁱ —Na1—F2—B1	49.76 (16)
B1 ⁱⁱⁱ —Na1—B1—F4	-35.8 (2)	F1 ⁱⁱ —Na1—F2—B1	-143.4 (2)
F1 ⁱ —Na1—B1—F1	45.78 (13)	F1 ⁱⁱⁱ —Na1—F2—B1	-95.75 (12)
F1 ⁱⁱ —Na1—B1—F1	-71.6 (4)	F1—Na1—F2—B1	-2.94 (12)
F1 ⁱⁱⁱ —Na1—B1—F1	-96.70 (14)	F2 ⁱ —Na1—F2—B1	79.09 (14)
F2—Na1—B1—F1	-174.8 (2)	F2 ⁱⁱ —Na1—F2—B1	142.68 (12)
F2 ⁱ —Na1—B1—F1	92.50 (13)	F2 ⁱⁱⁱ —Na1—F2—B1	-153.72 (12)
F2 ⁱⁱ —Na1—B1—F1	130.10 (12)	B1 ⁱ —Na1—F2—B1	66.77 (10)
F2 ⁱⁱⁱ —Na1—B1—F1	-148.85 (13)	B1 ⁱⁱ —Na1—F2—B1	169.42 (10)
B1 ⁱ —Na1—B1—F1	69.58 (15)	B1 ⁱⁱⁱ —Na1—F2—B1	-124.56 (13)
B1 ⁱⁱ —Na1—B1—F1	153.24 (12)	N4—C4—C7—C6	-1.5 (3)
B1 ⁱⁱⁱ —Na1—B1—F1	-123.10 (13)	N3—C4—C7—C6	177.86 (16)
F1 ⁱ —Na1—B1—F3	-47.1 (2)	N4—C4—C7—C2	-178.91 (18)
F1 ⁱⁱ —Na1—B1—F3	-164.5 (2)	N3—C4—C7—C2	0.4 (3)
F1 ⁱⁱⁱ —Na1—B1—F3	170.43 (18)	N5—C6—C7—C4	2.3 (3)
F1—Na1—B1—F3	-92.9 (2)	N6—C6—C7—C4	-176.63 (17)
F2—Na1—B1—F3	92.3 (2)	N5—C6—C7—C2	179.74 (18)
F2 ⁱ —Na1—B1—F3	-0.36 (17)	N6—C6—C7—C2	0.8 (3)
F2 ⁱⁱ —Na1—B1—F3	37.2 (2)	N1—C2—C7—C4	179.01 (18)
F2 ⁱⁱⁱ —Na1—B1—F3	118.28 (17)	N2—C2—C7—C4	0.1 (3)
B1 ⁱ —Na1—B1—F3	-23.3 (2)	N1—C2—C7—C6	1.7 (3)
B1 ⁱⁱ —Na1—B1—F3	60.38 (17)	N2—C2—C7—C6	-177.25 (18)
B1 ⁱⁱⁱ —Na1—B1—F3	144.04 (13)	N6—C1—N1—C2	1.5 (3)
F1 ⁱ —Na1—B1—F2	-139.43 (13)	N2—C2—N1—C1	176.21 (19)
F1 ⁱⁱ —Na1—B1—F2	103.2 (3)	C7—C2—N1—C1	-2.7 (3)
F1 ⁱⁱⁱ —Na1—B1—F2	78.08 (12)	N3—C3—N2—C2	-0.6 (3)
F1—Na1—B1—F2	174.8 (2)	N1—C2—N2—C3	-179.00 (18)
F2 ⁱ —Na1—B1—F2	-92.71 (14)	C7—C2—N2—C3	-0.1 (3)

F2 ⁱⁱ —Na1—B1—F2	-55.11 (18)	N4—C4—N3—C3	178.34 (18)
F2 ⁱⁱⁱ —Na1—B1—F2	25.93 (12)	C7—C4—N3—C3	-1.0 (3)
B1 ⁱ —Na1—B1—F2	-115.63 (11)	N2—C3—N3—C4	1.1 (3)
B1 ⁱⁱ —Na1—B1—F2	-31.97 (11)	N5—C5—N4—C4	1.6 (3)
B1 ⁱⁱⁱ —Na1—B1—F2	51.69 (14)	N3—C4—N4—C5	-179.66 (17)
F4—B1—F1—Na1	-124.27 (16)	C7—C4—N4—C5	-0.3 (3)
F3—B1—F1—Na1	115.40 (16)	N6—C6—N5—C5	177.64 (18)
F2—B1—F1—Na1	-4.51 (18)	C7—C6—N5—C5	-1.2 (3)
F1 ⁱ —Na1—F1—B1	-135.69 (13)	N4—C5—N5—C6	-0.8 (3)
F1 ⁱⁱ —Na1—F1—B1	151.86 (13)	N5—C6—N6—C1	179.05 (18)
F1 ⁱⁱⁱ —Na1—F1—B1	79.41 (15)	C7—C6—N6—C1	-2.1 (3)
F2—Na1—F1—B1	2.97 (12)	N1—C1—N6—C6	1.0 (3)

Symmetry codes: (i) $y, -x+1/2, z$; (ii) $-x+1/2, -y+1/2, z$; (iii) $-y+1/2, x, z$; (iv) $-y+1, x-1/2, -z$; (v) $y+1/2, -x+1, -z$; (vi) $-x+3/2, -y+1/2, z$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N3—H3N...N4 ^{vii}	0.88	2.05	2.930 (2)	176
N6—H6N...N5 ^{viii}	0.88	2.09	2.866 (2)	146
C1—H1...F3 ^{ix}	0.95	2.47	3.179 (2)	132
C1—H1...F4 ^{viii}	0.95	2.55	2.940 (3)	105
C3—H3...F2 ^{vii}	0.95	2.22	3.124 (2)	159
C5—H5...F3 ⁱⁱⁱ	0.95	2.40	3.097 (2)	130

Symmetry codes: (iii) $-y+1/2, x, z$; (vii) $-x+1, -y+1, -z+2$; (viii) $-y+1, x+1/2, -z+1$; (ix) $y, -x+3/2, z$.