

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,3,4,5,6-Pentabromophenol

Richard Betz, Peter Klüfers* and Peter Mayer

Department Chemie und Biochemie, Ludwig-Maximilians-Universität, Butenandtstrasse 5–13 (Haus D), 81377 München, Germany

Correspondence e-mail: kluef@cup.uni-muenchen.de


Received 3 September 2008; accepted 6 September 2008

 Key indicators: single-crystal X-ray study; $T = 200$ K; mean $\sigma(\text{C}–\text{C}) = 0.005$ Å; R factor = 0.030; wR factor = 0.075; data-to-parameter ratio = 20.0.

The title compound, $\text{C}_6\text{HBr}_5\text{O}$, is the perbrominated derivative of phenol. The molecule shows non-crystallographic mirror symmetry. Bond lengths between the C and Br atoms are normal. In the crystal structure, $\text{O}–\text{H}\cdots\text{O}$ hydrogen bonds connect the molecules into infinite strands. Dispersive $\text{Br}\cdots\text{Br}$ contacts are observed. No significant π – π stacking is obvious.

Related literature

For the structure of the perfluorinated derivative of phenol, see: Das *et al.* (2006); Gdaniec (2007). For the structure of 2,3,4,5,6-pentachlorophenol, see: Sakurai (1962).



Experimental

Crystal data

 $\text{C}_6\text{HBr}_5\text{O}$
 $M_r = 488.57$

 Monoclinic, $C2/c$
 $a = 32.3058$ (15) Å

 $b = 3.9957$ (2) Å
 $c = 16.1887$ (8) Å
 $\beta = 112.118$ (3)°
 $V = 1935.93$ (17) Å³
 $Z = 8$

 Mo $K\alpha$ radiation
 $\mu = 20.70$ mm⁻¹
 $T = 200$ (2) K
 $0.28 \times 0.08 \times 0.05$ mm

Data collection

 Nonius Kappa CCD diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 2001)
 $T_{\min} = 0.062$, $T_{\max} = 0.355$

 13465 measured reflections
 2219 independent reflections
 1930 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.054$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.074$
 $S = 1.03$
 2219 reflections

 111 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.88$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.02$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D–H\cdots A$	$D–H$	$H\cdots A$	$D\cdots A$	$D–H\cdots A$
$\text{O1}–\text{H1}\cdots\text{O1}^i$	0.84	2.19	2.844 (4)	134

 Symmetry code: (i) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: COLLECT (Nonius, 2004); cell refinement: SCALEPACK (Otwinowski & Minor 1997); data reduction: DENZO (Otwinowski & Minor 1997) and SCALEPACK; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and Mercury (Macrae *et al.*, 2006); software used to prepare material for publication: SHELXL97.

The authors thank Dr Peter Mayer for professional support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2109).

References

- Das, D., Banerjee, R., Mondal, R., Howard, J. A. K., Boese, R. & Desiraju, G. R. (2006). *Chem. Commun.* pp. 555–557.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Gdaniec, M. (2007). *CrystEngComm*, **9**, 286–288.
 Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
 Nonius (2004). COLLECT. Nonius BV, Delft, The Netherlands.
 Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography, Part A*, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
 Sakurai, T. (1962). *Acta Cryst.* **15**, 1164–1173.
 Sheldrick, G. M. (2001). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2008). E64, o1921 [doi:10.1107/S1600536808028602]

2,3,4,5,6-Pentabromophenol

Richard Betz, Peter Klüfers and Peter Mayer

S1. Comment

During efforts to obtain tetraaryloxy derivatives of orthocarbonic acid it was interesting to determine the influence of bonding to one central carbon atom on geometric parameters of the ligands. Thus the crystal structure of 2,3,4,5,6-pentabromophenol was determined.

In the molecule (Fig. 1), C—C—C angles adopt values covering a range from 119.1 (3)° on the C atom bonded to the hydroxy group to 120.7 (3)° on one of the C atoms in *ortho*-position to the hydroxy group. The alterations between the C—C—C angles thus are less pronounced than in the perfluorinated derivative of phenol, where the angle on the C atom bearing the hydroxy group was found at a value slightly above 116° (Gdaniec, 2007). The values more closely resemble the ones apparent in the molecular structure of the perchlorinated derivative, yet the smallest C—C—C angle is not present on the C atom bearing the hydroxy group in that compound (Sakurai, 1962).


In the crystal structure H-bonds connect the molecules to infinite strands along [010] (Fig. 2). A bifurcation of the hydrogen bond between oxygen and one of the halogen atoms in *ortho*-position was not observed. This is in contrast to 2,3,4,5,6-pentachlorophenol, where the presence of such a bifurcated hydrogen bond was substantiated upon nuclear quadrupole resonance spectra for the Cl atoms (Sakurai, 1962). Additionally, dispersive Br...Br interactions between the Br atoms in both *meta*-positions to the hydroxy group are observed. The range of these interactions falls by about 0.1 Å below the sum of van der Waals radii of the respective atoms. These connect the molecules to chains along [001]. No significant π -stacking is apparent in the crystal structure. The molecular packing is shown in Fig. 3.

S2. Experimental


The compound was obtained commercially from Aldrich. Crystals suitable for X-ray diffraction were obtained upon recrystallization of the compound from boiling toluene.

S3. Refinement

The H atom was located in a difference map and refined as riding on its parent O atom with an $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$.

**Figure 1**

The molecular structure of the title compound, with atom labels. The displacement ellipsoids are drawn at 50% probability level. H atom is presented as a small sphere of arbitrary radius.

**Figure 2**

The crystal packing diagram, viewed along [010].

2,3,4,5,6-Pentabromophenol

Crystal data

C₆HBr₅O $M_r = 488.57$

Monoclinic, C2/c

Hall symbol: -C 2yc

 $a = 32.3058 (15) \text{ \AA}$ $b = 3.9957 (2) \text{ \AA}$ $c = 16.1887 (8) \text{ \AA}$ $\beta = 112.118 (3)^\circ$ $V = 1935.93 (17) \text{ \AA}^3$ $Z = 8$ $F(000) = 1760$ $D_x = 3.353 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 8265 reflections

 $\theta = 3.1\text{--}27.5^\circ$ $\mu = 20.70 \text{ mm}^{-1}$ $T = 200 \text{ K}$

Rod, colourless

 $0.28 \times 0.08 \times 0.05 \text{ mm}$

Data collection

Nonius Kappa CCD

diffractometer

Radiation source: Rotating anode

MONTELE, graded multilayered X-ray optics

monochromator

Rotation images; thick slices scans

Absorption correction: multi-scan

(SADABS; Sheldrick, 2001)

 $T_{\min} = 0.062$, $T_{\max} = 0.355$

13465 measured reflections

2219 independent reflections

1930 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.054$ $\theta_{\text{max}} = 27.6^\circ$, $\theta_{\text{min}} = 3.5^\circ$ $h = -41 \rightarrow 41$ $k = -4 \rightarrow 5$ $l = -21 \rightarrow 21$

Refinement

Refinement on F^2

Least-squares matrix: Full

 $R[F^2 > 2\sigma(F^2)] = 0.030$ $wR(F^2) = 0.075$ $S = 1.03$

2219 reflections

111 parameters

0 restraints

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0374P)^2 + 5.8817P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\text{max}} = 0.001$ $\Delta\rho_{\text{max}} = 0.88 \text{ e \AA}^{-3}$ $\Delta\rho_{\text{min}} = -1.02 \text{ e \AA}^{-3}$ Extinction correction: SHELXL97 (Sheldrick,
2008)

Extinction coefficient: 0.00087 (8)

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.204848 (13)	0.51248 (10)	0.41077 (2)	0.03029 (14)
Br2	0.096769 (13)	0.38703 (10)	0.36125 (2)	0.02821 (13)

Br3	0.024427 (13)	0.61492 (11)	0.16493 (3)	0.03106 (14)
Br4	0.060474 (12)	0.97946 (11)	0.02144 (2)	0.02709 (13)
Br5	0.169251 (13)	1.09979 (10)	0.07733 (2)	0.02651 (13)
O1	0.22213 (8)	0.8423 (7)	0.26447 (18)	0.0289 (6)
H1	0.2271	0.9643	0.2270	0.043*
C1	0.17732 (11)	0.7978 (9)	0.2394 (2)	0.0213 (7)
C2	0.16196 (12)	0.6421 (9)	0.2996 (2)	0.0216 (7)
C3	0.11662 (12)	0.5889 (9)	0.2773 (2)	0.0211 (7)
C4	0.08601 (11)	0.6883 (9)	0.1945 (2)	0.0213 (7)
C5	0.10107 (12)	0.8416 (8)	0.1339 (2)	0.0209 (7)
C6	0.14663 (12)	0.8957 (8)	0.1567 (2)	0.0197 (7)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0248 (2)	0.0387 (3)	0.0229 (2)	0.00504 (16)	0.00385 (16)	0.00541 (16)
Br2	0.0300 (2)	0.0331 (2)	0.0244 (2)	-0.00243 (15)	0.01352 (16)	0.00349 (14)
Br3	0.0171 (2)	0.0443 (3)	0.0318 (2)	-0.00324 (15)	0.00924 (16)	0.00412 (16)
Br4	0.0201 (2)	0.0377 (2)	0.0212 (2)	0.00309 (15)	0.00524 (15)	0.00423 (14)
Br5	0.0238 (2)	0.0331 (2)	0.0255 (2)	-0.00301 (14)	0.01252 (16)	0.00230 (14)
O1	0.0160 (12)	0.0393 (16)	0.0315 (14)	0.0005 (11)	0.0091 (11)	0.0025 (12)
C1	0.0146 (16)	0.0222 (16)	0.0260 (17)	-0.0013 (14)	0.0066 (13)	-0.0031 (14)
C2	0.0188 (18)	0.0236 (18)	0.0201 (16)	0.0003 (14)	0.0048 (13)	-0.0015 (13)
C3	0.0237 (19)	0.0211 (16)	0.0208 (17)	-0.0001 (13)	0.0110 (14)	-0.0028 (13)
C4	0.0146 (16)	0.0261 (17)	0.0244 (17)	-0.0029 (14)	0.0087 (13)	-0.0038 (14)
C5	0.0196 (17)	0.0237 (17)	0.0184 (15)	-0.0001 (14)	0.0061 (13)	-0.0027 (13)
C6	0.0213 (17)	0.0215 (17)	0.0207 (16)	-0.0026 (14)	0.0129 (13)	-0.0009 (13)

Geometric parameters (Å, °)

Br1—C2	1.884 (3)	C1—C6	1.389 (5)
Br2—C3	1.888 (4)	C1—C2	1.395 (5)
Br3—C4	1.886 (3)	C2—C3	1.387 (5)
Br4—C5	1.882 (3)	C3—C4	1.391 (5)
Br5—C6	1.886 (4)	C4—C5	1.390 (5)
O1—C1	1.360 (4)	C5—C6	1.393 (5)
O1—H1	0.8400		
C1—O1—H1	109.5	C5—C4—C3	119.7 (3)
O1—C1—C6	122.9 (3)	C5—C4—Br3	120.4 (2)
O1—C1—C2	117.9 (3)	C3—C4—Br3	120.0 (3)
C6—C1—C2	119.1 (3)	C4—C5—C6	119.8 (3)
C3—C2—C1	120.3 (3)	C4—C5—Br4	120.7 (3)
C3—C2—Br1	122.1 (3)	C6—C5—Br4	119.5 (3)
C1—C2—Br1	117.6 (3)	C1—C6—C5	120.7 (3)
C2—C3—C4	120.4 (3)	C1—C6—Br5	117.4 (3)
C2—C3—Br2	119.3 (3)	C5—C6—Br5	121.9 (3)
C4—C3—Br2	120.3 (3)		

O1—C1—C2—C3	-179.7 (3)	C3—C4—C5—C6	-0.2 (5)
C6—C1—C2—C3	-0.6 (5)	Br3—C4—C5—C6	179.7 (3)
O1—C1—C2—Br1	0.7 (4)	C3—C4—C5—Br4	-179.9 (3)
C6—C1—C2—Br1	179.8 (3)	Br3—C4—C5—Br4	0.1 (4)
C1—C2—C3—C4	0.4 (5)	O1—C1—C6—C5	179.4 (3)
Br1—C2—C3—C4	180.0 (3)	C2—C1—C6—C5	0.4 (5)
C1—C2—C3—Br2	-178.5 (3)	O1—C1—C6—Br5	-0.2 (5)
Br1—C2—C3—Br2	1.1 (4)	C2—C1—C6—Br5	-179.2 (3)
C2—C3—C4—C5	0.0 (5)	C4—C5—C6—C1	0.0 (5)
Br2—C3—C4—C5	178.9 (3)	Br4—C5—C6—C1	179.7 (3)
C2—C3—C4—Br3	-179.9 (3)	C4—C5—C6—Br5	179.6 (3)
Br2—C3—C4—Br3	-1.1 (4)	Br4—C5—C6—Br5	-0.7 (4)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots O1 ⁱ	0.84	2.19	2.844 (4)	134

Symmetry code: (i) $-x+1/2, y+1/2, -z+1/2$.