

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

5-(2-Phenylethynyl)isobenzofuran-1,3-dione

 Mei-Jia Yang,^a Jian Men,^a Xiao-Yan Ma,^b Shi-Xu Yi^a and Guo-Wei Gao^{a*}
^aCollege of Chemistry, Sichuan University, Chengdu 610064, People's Republic of China, and ^bCollege of Materials and Chemical Engineering, Chengdu University of Technology, Chengdu 610059, People's Republic of China

Correspondence e-mail: gaosunday@yahoo.com.cn


Received 30 August 2008; accepted 4 September 2008

 Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.065; wR factor = 0.158; data-to-parameter ratio = 12.7.

The title compound, $\text{C}_{16}\text{H}_8\text{O}_3$, was synthesized by the Pd-coupling reaction of phenylacetylene with 4-bromophthalic anhydride. The phenyl and isobenzofuran rings are nearly coplanar, forming a dihedral angle of 6.70 (10)°. In the crystal structure, centrosymmetrically related molecules are linked into dimers by $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds.

Related literature

For a general background to the synthesis and applications of the title compound, see: Hergenrother & Smith (1994, 1996); Takekoshi & Terry (1994); Urazoe *et al.* (2005); Urazoe & Mori (2006). For the properties of polyimides, see: Feger *et al.* (1989); Ghosh & Mittal (1996). For the crystal structure of related compounds, see: Wright & Schorzman (2000).



Experimental

Crystal data

 $\text{C}_{16}\text{H}_8\text{O}_3$
 $M_r = 248.22$
 Triclinic, $P\bar{1}$
 $a = 6.998$ (3) Å
 $b = 7.518$ (3) Å
 $c = 11.683$ (4) Å

 $\alpha = 89.06$ (2)°
 $\beta = 79.31$ (3)°
 $\gamma = 81.04$ (2)°
 $V = 596.6$ (4) Å³
 $Z = 2$

 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 294$ (2) K
 $0.50 \times 0.40 \times 0.22$ mm

Data collection

 Enraf-Nonius CAD4 diffractometer
 Absorption correction: none
 2227 measured reflections
 2198 independent reflections

 1105 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.007$
 3 standard reflections every 100 reflections
 intensity decay: 1.8%

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.064$
 $wR(F^2) = 0.158$
 $S = 1.09$
 2198 reflections

 173 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.21$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.21$ e Å⁻³
Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C16}-\text{H16}\cdots\text{O2}^i$	0.93	2.56	3.436 (5)	158

 Symmetry code: (i) $-x, -y + 1, -z + 1$.

Data collection: *DIFRAC* (Gabe & White, 1993); cell refinement: *DIFRAC*; data reduction: *NRCVAX* (Gabe *et al.*, 1989); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

The authors thank Mr Zhi-Hua Mao of Sichuan University for the X-ray data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RZ2241).

References

- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Feger, C., Khohasteh, M. M. & McGrath, J. E. (1989). *Polyimides: Chemistry, Materials, and Characterization*. Amsterdam: Elsevier.
 Gabe, E. J., Le Page, Y., Charland, J.-P., Lee, F. L. & White, P. S. (1989). *J. Appl. Cryst.* **22**, 384–387.
 Gabe, E. J. & White, P. S. (1993). *DIFRAC*. American Crystallographic Association Pittsburgh Meeting, Abstract PA 104.
 Ghosh, M. K. & Mittal, K. L. (1996). *Polyimides: Fundamentals and Applications*. New York: Dekker.
 Hergenrother, P. M. & Smith, J. G. Jr (1994). *Polymer*, **35**, 4857–4864.
 Hergenrother, P. M. & Smith, J. G. Jr (1996). US Patent 5 567 800.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Takekoshi, T. & Terry, J. M. (1994). *Polymer*, **35**, 4874–4880.
 Urazoe, D. & Mori, H. (2006). Jpn Patent JP 2 006 151 904.
 Urazoe, D., Mori, H. & Yamakawa, K. (2005). US Patent 2 005 215 820.
 Wright, M. E. & Schorzman, D. A. (2000). *Macromolecules*, **33**, 8611–8617.

supporting information

Acta Cryst. (2008). E64, o2193 [doi:10.1107/S1600536808028407]

5-(2-Phenylethynyl)isobenzofuran-1,3-dione

Mei-Jia Yang, Jian Men, Xiao-Yan Ma, Shi-Xu Yi and Guo-Wei Gao

S1. Comment

Polyimides are well known for possessing excellent thermal and oxidative stability, as well as excellent mechanical properties (Ghosh & Mittal, 1996; Feger *et al.*, 1989). The title compound is used as terminal endcapping agent which imparts thermal curability, thermal resistance and solvent resistance to polyimide (Hergenrother & Smith, 1994, 1996; Takekoshi & Terry, 1994). Further, various types of method for producing 4-phenylethynylphthalic anhydride have been described (Urazoe *et al.*, 2005; Urazoe & Mori, 2006). We report here the crystal structure of the title compound.


In the molecule of the title compound (Fig. 1) the phenyl and isobenzofurane rings are nearly coplanar, making a dihedral angle of 6.70 (10)°. Bond distances of the ethyne chain show similar values to those in C₂₂H₁₃O₂N (Wright & Schorzman, 2000), with the C7—C9, C9—C10 and C10—C11 distances of 1.418 (4), 1.203 (4) and 1.429 (4) Å, respectively. The bond angles within the ethyne chain are slightly bent, with the C7—C9—C10 and C9—C10—C11 angles of 176.3 (4) and 173.6 (4)°, respectively. The isobenzofurane ring is flat, atoms C2 deviating only by 0.027 (5) Å from the mean plane. In the crystal structure, centrosymmetrically related molecules are linked into dimers by intermolecular C—H···O hydrogen interactions (Table 1).

S2. Experimental

4-Bromophthalic anhydride (5.00 g, 22.0 mmol), phenylacetylene (2.69 g, 26.4 mmol), PdCl₂(PPh₃)₂ (0.11 g, 0.157 mmol), and PPh₃ (0.22 g, 0.840 mmol) were dissolved in 40 ml of dry NEt₃ under argon, and the mixture was heated to 333 K. Then CuI (0.10 g, 0.524 mmol) was added and the solution was stirred at 353 K for 12 h. The precipitated triethylammonium bromide was separated after cooling and the solvent was evaporated. The residue was recrystallized from toluene/n-hexane (1:1 v/v) twice to give 4-phenylethynylphthalic anhydride as pale yellow crystals (yield 83.6%; m.p. 424–425 K). Colourless crystals suitable for X-ray analysis were obtained by slow evaporation of an acetic anhydride solution at room temperature.

S3. Refinement

H atoms were positioned geometrically (C—H = 0.93 Å) and refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

**Figure 1**

The molecular structure of the title compound, with displacement ellipsoids drawn at the 30% probability level.

5-(2-Phenylethynyl)isobenzofuran-1,3-dione

Crystal data

$C_{16}H_8O_3$
 $M_r = 248.22$
 Triclinic, $P\bar{1}$
 Hall symbol: $-P\ 1$
 $a = 6.998\ (3)\ \text{\AA}$
 $b = 7.518\ (3)\ \text{\AA}$
 $c = 11.683\ (4)\ \text{\AA}$
 $\alpha = 89.06\ (2)^\circ$
 $\beta = 79.31\ (3)^\circ$
 $\gamma = 81.04\ (2)^\circ$
 $V = 596.6\ (4)\ \text{\AA}^3$

$Z = 2$
 $F(000) = 256$
 $D_x = 1.382\ \text{Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$
 Cell parameters from 25 reflections
 $\theta = 4.5\text{--}11.8^\circ$
 $\mu = 0.10\ \text{mm}^{-1}$
 $T = 294\ \text{K}$
 Block, colourless
 $0.50 \times 0.40 \times 0.22\ \text{mm}$

Data collection

Enraf–Nonius CAD4
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 $\omega/2\theta$ scans
 2227 measured reflections
 2198 independent reflections
 1105 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.007$
 $\theta_{\text{max}} = 25.5^\circ$, $\theta_{\text{min}} = 2.7^\circ$
 $h = -8 \rightarrow 8$
 $k = -3 \rightarrow 9$
 $l = -14 \rightarrow 14$
 3 standard reflections every 100 reflections
 intensity decay: 1.8%

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.064$
 $wR(F^2) = 0.158$
 $S = 1.09$
 2198 reflections
 173 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0535P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.22\ \text{e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.21\ \text{e \AA}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick,
 1997), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.014 (4)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.5201 (3)	0.1623 (3)	0.3262 (2)	0.0689 (8)
O2	0.3130 (4)	0.3047 (4)	0.4814 (2)	0.0912 (10)
O3	0.6535 (3)	0.0590 (3)	0.1454 (2)	0.0814 (9)
C1	0.3406 (5)	0.2634 (5)	0.3814 (4)	0.0655 (11)
C2	0.5145 (5)	0.1372 (5)	0.2089 (4)	0.0621 (10)
C3	0.2138 (5)	0.2980 (4)	0.2931 (3)	0.0484 (9)
C4	0.3202 (4)	0.2205 (4)	0.1897 (3)	0.0489 (9)
C5	0.2391 (5)	0.2304 (4)	0.0913 (3)	0.0604 (10)
H5	0.3096	0.1777	0.0216	0.072*
C6	0.0505 (5)	0.3205 (4)	0.0986 (3)	0.0570 (10)
H6	-0.0073	0.3273	0.0329	0.068*
C7	-0.0562 (4)	0.4021 (4)	0.2025 (3)	0.0505 (9)
C8	0.0267 (5)	0.3885 (4)	0.3023 (3)	0.0521 (9)
H8	-0.0430	0.4392	0.3728	0.062*
C9	-0.2470 (5)	0.5020 (4)	0.2070 (3)	0.0607 (10)
C10	-0.4084 (5)	0.5899 (4)	0.2170 (3)	0.0603 (10)
C11	-0.6025 (4)	0.6902 (4)	0.2426 (3)	0.0534 (9)
C12	-0.7133 (5)	0.7369 (4)	0.1575 (3)	0.0594 (10)
H12	-0.6615	0.7056	0.0801	0.071*
C13	-0.9015 (5)	0.8304 (5)	0.1881 (4)	0.0667 (11)
H13	-0.9752	0.8638	0.1305	0.080*
C14	-0.9807 (5)	0.8744 (5)	0.3007 (4)	0.0697 (11)
H14	-1.1088	0.9354	0.3204	0.084*
C15	-0.8719 (5)	0.8288 (5)	0.3852 (3)	0.0726 (11)
H15	-0.9272	0.8589	0.4624	0.087*
C16	-0.6828 (5)	0.7396 (4)	0.3586 (3)	0.0647 (11)
H16	-0.6088	0.7123	0.4167	0.078*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0505 (16)	0.0776 (17)	0.076 (2)	0.0049 (13)	-0.0171 (14)	0.0015 (15)
O2	0.081 (2)	0.137 (3)	0.0513 (18)	0.0076 (17)	-0.0214 (16)	-0.0083 (17)
O3	0.0510 (16)	0.0859 (19)	0.095 (2)	0.0177 (14)	-0.0024 (15)	-0.0271 (16)
C1	0.054 (2)	0.075 (3)	0.065 (3)	0.000 (2)	-0.011 (2)	0.001 (2)
C2	0.056 (3)	0.058 (2)	0.070 (3)	-0.0026 (19)	-0.010 (2)	-0.004 (2)

C3	0.046 (2)	0.049 (2)	0.049 (2)	-0.0002 (16)	-0.0090 (18)	-0.0020 (17)
C4	0.0405 (19)	0.047 (2)	0.056 (2)	0.0001 (15)	-0.0068 (18)	-0.0085 (17)
C5	0.057 (2)	0.061 (2)	0.058 (2)	0.0036 (18)	-0.007 (2)	-0.0159 (19)
C6	0.053 (2)	0.064 (2)	0.051 (2)	0.0041 (18)	-0.0135 (18)	-0.0154 (18)
C7	0.041 (2)	0.050 (2)	0.057 (2)	0.0056 (16)	-0.0105 (18)	-0.0051 (17)
C8	0.044 (2)	0.058 (2)	0.048 (2)	0.0031 (17)	-0.0006 (18)	-0.0095 (17)
C9	0.054 (2)	0.059 (2)	0.068 (3)	-0.004 (2)	-0.013 (2)	-0.0032 (19)
C10	0.061 (2)	0.058 (2)	0.064 (3)	-0.006 (2)	-0.019 (2)	-0.0029 (19)
C11	0.036 (2)	0.0445 (19)	0.076 (3)	0.0035 (16)	-0.009 (2)	-0.0023 (18)
C12	0.058 (2)	0.058 (2)	0.061 (2)	-0.0038 (18)	-0.012 (2)	-0.0039 (18)
C13	0.057 (2)	0.063 (2)	0.085 (3)	-0.005 (2)	-0.028 (2)	-0.003 (2)
C14	0.055 (2)	0.063 (2)	0.083 (3)	0.0072 (19)	-0.006 (2)	-0.014 (2)
C15	0.074 (3)	0.070 (3)	0.066 (3)	0.011 (2)	-0.009 (2)	-0.016 (2)
C16	0.061 (2)	0.067 (3)	0.065 (3)	0.004 (2)	-0.020 (2)	-0.008 (2)

Geometric parameters (Å, °)

O1—C2	1.395 (4)	C8—H8	0.9300
O1—C1	1.411 (4)	C9—C10	1.203 (4)
O2—C1	1.186 (4)	C10—C11	1.429 (4)
O3—C2	1.187 (4)	C11—C12	1.379 (5)
C1—C3	1.476 (5)	C11—C16	1.399 (4)
C2—C4	1.461 (4)	C12—C13	1.379 (4)
C3—C8	1.365 (4)	C12—H12	0.9300
C3—C4	1.381 (4)	C13—C14	1.354 (4)
C4—C5	1.369 (4)	C13—H13	0.9300
C5—C6	1.375 (4)	C14—C15	1.363 (5)
C5—H5	0.9300	C14—H14	0.9300
C6—C7	1.397 (4)	C15—C16	1.370 (4)
C6—H6	0.9300	C15—H15	0.9300
C7—C8	1.391 (4)	C16—H16	0.9300
C7—C9	1.418 (4)		
C2—O1—C1	109.3 (3)	C3—C8—H8	121.2
O2—C1—O1	121.4 (4)	C7—C8—H8	121.2
O2—C1—C3	131.4 (4)	C10—C9—C7	176.3 (4)
O1—C1—C3	107.2 (3)	C9—C10—C11	173.6 (4)
O3—C2—O1	120.3 (3)	C12—C11—C16	119.3 (3)
O3—C2—C4	132.2 (4)	C12—C11—C10	122.3 (3)
O1—C2—C4	107.5 (3)	C16—C11—C10	118.3 (3)
C8—C3—C4	122.4 (3)	C11—C12—C13	119.6 (3)
C8—C3—C1	130.3 (3)	C11—C12—H12	120.2
C4—C3—C1	107.3 (3)	C13—C12—H12	120.2
C5—C4—C3	120.6 (3)	C14—C13—C12	121.0 (4)
C5—C4—C2	130.8 (3)	C14—C13—H13	119.5
C3—C4—C2	108.7 (3)	C12—C13—H13	119.5
C4—C5—C6	118.1 (3)	C13—C14—C15	119.7 (4)
C4—C5—H5	121.0	C13—C14—H14	120.2

C6—C5—H5	121.0	C15—C14—H14	120.2
C5—C6—C7	121.6 (3)	C14—C15—C16	121.3 (4)
C5—C6—H6	119.2	C14—C15—H15	119.3
C7—C6—H6	119.2	C16—C15—H15	119.3
C8—C7—C6	119.8 (3)	C15—C16—C11	119.0 (3)
C8—C7—C9	119.4 (3)	C15—C16—H16	120.5
C6—C7—C9	120.8 (3)	C11—C16—H16	120.5
C3—C8—C7	117.6 (3)		
C2—O1—C1—O2	-178.4 (4)	C2—C4—C5—C6	-179.3 (3)
C2—O1—C1—C3	1.8 (4)	C4—C5—C6—C7	0.7 (5)
C1—O1—C2—O3	178.1 (3)	C5—C6—C7—C8	-1.8 (5)
C1—O1—C2—C4	-2.1 (4)	C5—C6—C7—C9	176.9 (3)
O2—C1—C3—C8	0.0 (7)	C4—C3—C8—C7	-0.4 (5)
O1—C1—C3—C8	179.7 (3)	C1—C3—C8—C7	179.1 (3)
O2—C1—C3—C4	179.5 (4)	C6—C7—C8—C3	1.6 (5)
O1—C1—C3—C4	-0.8 (4)	C9—C7—C8—C3	-177.2 (3)
C8—C3—C4—C5	-0.7 (5)	C16—C11—C12—C13	-0.4 (5)
C1—C3—C4—C5	179.7 (3)	C10—C11—C12—C13	178.6 (3)
C8—C3—C4—C2	179.1 (3)	C11—C12—C13—C14	-1.2 (5)
C1—C3—C4—C2	-0.4 (4)	C12—C13—C14—C15	1.3 (6)
O3—C2—C4—C5	1.1 (7)	C13—C14—C15—C16	0.3 (6)
O1—C2—C4—C5	-178.6 (3)	C14—C15—C16—C11	-1.8 (6)
O3—C2—C4—C3	-178.7 (4)	C12—C11—C16—C15	1.9 (5)
O1—C2—C4—C3	1.5 (4)	C10—C11—C16—C15	-177.2 (3)
C3—C4—C5—C6	0.5 (5)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C16—H16 \cdots O2 ⁱ	0.93	2.56	3.436 (5)	158

Symmetry code: (i) $-x, -y+1, -z+1$.