

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

5-(4-Chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazole-3-carboxylic acid

Wei Wang and Zheng Fang*

College of Life Sciences and Pharmaceutical Engineering, Nanjing University of Technology, Ximofan Road No. 5 Nanjing, Nanjing 210009, People's Republic of China

Correspondence e-mail: fzcpcu@163.com


Received 13 November 2008; accepted 16 November 2008

 Key indicators: single-crystal X-ray study; $T = 294$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; R factor = 0.072; wR factor = 0.181; data-to-parameter ratio = 14.3.

The asymmetric unit of the title compound, $\text{C}_{17}\text{H}_{11}\text{Cl}_3\text{N}_2\text{O}_2$, contains two independent molecules; the pyrazole rings are oriented with respect to the chlorophenyl and dichlorophenyl rings at dihedral angles of 43.00 (3) and 65.06 (4)°, respectively, in one molecule, and 51.17 (3) and 69.99 (3)°, respectively, in the other. Pairs of intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into dimers. In the crystal structure, there are $\pi-\pi$ contacts between the pyrazole rings and dichlorophenyl rings [centroid-centroid distances = 3.859 (3) and 3.835 (3) Å].

Related literature

For bond-length data, see: Allen *et al.* (1987). For the chemical background, see: Tang *et al.* (2007).



Experimental

Crystal data

$\text{C}_{17}\text{H}_{11}\text{Cl}_3\text{N}_2\text{O}_2$
 $M_r = 381.63$
 Monoclinic, $P2_1/c$
 $a = 13.192$ (3) Å
 $b = 8.8170$ (18) Å
 $c = 30.012$ (6) Å
 $\beta = 102.42$ (3)°
 $V = 3409.1$ (13) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.55$ mm⁻¹
 $T = 294$ (2) K
 $0.30 \times 0.20 \times 0.10$ mm

Data collection

Enraf-Nonius CAD-4 diffractometer
 Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.853$, $T_{\max} = 0.947$
 6479 measured reflections
 6190 independent reflections
 2893 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$
 3 standard reflections
 frequency: 120 min
 intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.072$
 $wR(F^2) = 0.181$
 $S = 0.99$
 6190 reflections
 433 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.33$ e Å⁻³
 $\Delta\rho_{\min} = -0.28$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O}2-\text{H}2A\cdots\text{O}4$	0.85	1.74	2.564 (7)	163
$\text{O}3-\text{H}3B\cdots\text{O}1$	0.85	1.89	2.723 (6)	165

 Symmetry codes: (i) $-x + 1, -y, -z$; (ii) $-x + 1, -y + 1, -z$.

Data collection: *CAD-4 Software* (Enraf-Nonius, 1989); cell refinement: *CAD-4 Software*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2003); software used to prepare material for publication: *SHELXL97*.

The authors thank the Center for Testing and Analysis, Nanjing University, for the support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HK2574).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
 Enraf-Nonius (1989). *CAD-4 Software*. Enraf-Nonius, Delft, The Netherlands.
 Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
 Tang, L. H., Tao, L., Chen, H. B. & Zhong, B. H. (2007). *Chin. J. Pharm.* **38**, 252–254.

supporting information

Acta Cryst. (2008). E64, o2413 [doi:10.1107/S1600536808038105]

5-(4-Chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxylic acid

Wei Wang and Zheng Fang

S1. Comment

Some derivatives of benzoic acid are important chemical materials. We report herein the crystal structure of the title compound.

The asymmetric unit of the title compound contains two crystallographically independent molecules (Fig. 1). The bond lengths (Allen *et al.*, 1987) and angles are within normal ranges. Rings A (C1-C6), B (C7-C12), C (N1/N2/C13-C15) and D (C18-C23), E (C24-C29), F (N3/N4/C30-C32) are, of course, planar and they are oriented at dihedral angles of A/B = 58.42 (3)°, A/C = 65.06 (4), B/C = 43.00 (3)° and D/E = 57.07 (4)°, D/F = 69.99 (3)°, E/F = 51.17 (3)°. The intramolecular O-H...O hydrogen bonds (Table 1) link the molecules (Fig. 1), in which they may be effective in the stabilization of the structure.

In the crystal structure, there are π - π contacts between the pyrazole rings and dichlorophenyl rings, Cg1—Cg2ⁱ and Cg3—Cg5ⁱⁱ [symmetry codes: (i) $x + 1, y, z$; (ii) $1 - x, y, 1/2 - z$, where Cg1, Cg2, Cg3 and Cg5 are centroids of the rings C (N1/N2/C13-C15), F (N3/N4/C30-C32), A (C3-C8) and D (C18-C23), respectively] may stabilize the structure, with centroid-centroid distances of 3.859 (3) Å and 3.835 (3) Å, respectively. There also exist C—H... π contacts (Table 1) between the methyl groups and the chlorophenyl rings.

S2. Experimental

For the preparation of the title compound, 2,4-dichlorophenylhydrazine hydrochloride (13.3 g) diluted in ethanol (20 ml) is added to ethyl 4-(4-chlorophenyl)-3-methyl-2,4-dioxobutanoate (17.6 g) diluted in toluene (50 ml) and the mixture is stirred for 18 h at room temperature. Without isolating the hydrazone, paratoluenesulfonic acid (0.56 g) is added, and the ternary azeotrope (water, ethanol, toluene) is distilled. Toluene reflux is continued for 1 h and the reaction mixture is cooled to room temperature. The insoluble material is filtered off. The solvents are removed under reduced pressure to give an oil. KOH (8.1 g) in pellets are added to a solution of the oil obtained in the previous step in MeOH (100 ml). The mixture is left for 1 h at room temperature and the solvents are decanted into water (200 ml) at 333 K. Hydrochloric acid is then added to the aqueous phase until pH = 1.5. The colorless crystals formed are filtered off, washed with water and dried under vacuum to give the expected product (yield; 9.9 g). Crystals suitable for X-ray analysis were obtained by slow evaporation of an acetic acid solution.

S3. Refinement

H atoms were positioned geometrically, with O-H = 0.85 Å (for OH) and C-H = 0.93 and 0.96 Å for aromatic and methyl H, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C}, \text{O})$, where $x = 1.2$ for aromatic H and $x = 1.5$ for all other H atoms.




Figure 1

The molecular structure of the title molecule, with the atom-numbering scheme.

5-(4-Chlorophenyl)-1-(2,4-dichlorophenyl)-4-methyl-1H-pyrazole-3-carboxylic acid

Crystal data

$C_{17}H_{11}Cl_3N_2O_2$
 $M_r = 381.63$
 Monoclinic, $P2_1/c$
 Hall symbol: $-P\ 2_1/c$
 $a = 13.192(3)\ \text{\AA}$
 $b = 8.8170(18)\ \text{\AA}$
 $c = 30.012(6)\ \text{\AA}$
 $\beta = 102.42(3)^\circ$
 $V = 3409.1(13)\ \text{\AA}^3$
 $Z = 8$

$F(000) = 1552$
 $D_x = 1.487\ \text{Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$
 Cell parameters from 25 reflections
 $\theta = 10\text{--}13^\circ$
 $\mu = 0.55\ \text{mm}^{-1}$
 $T = 294\ \text{K}$
 Block, colorless
 $0.30 \times 0.20 \times 0.10\ \text{mm}$

Data collection

Enraf–Nonius CAD-4
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 $\omega/2\theta$ scans
 Absorption correction: ψ scan
 (North *et al.*, 1968)
 $T_{\min} = 0.853$, $T_{\max} = 0.947$
 6479 measured reflections

6190 independent reflections
 2893 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.038$
 $\theta_{\max} = 25.3^\circ$, $\theta_{\min} = 1.4^\circ$
 $h = 0 \rightarrow 15$
 $k = 0 \rightarrow 10$
 $l = -36 \rightarrow 36$
 3 standard reflections every 120 min
 intensity decay: 1%

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.072$
 $wR(F^2) = 0.181$
 $S = 0.99$
 6190 reflections
 433 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.06P)^2 + 5P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.33\ \text{e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.28\ \text{e \AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.3235 (2)	0.6137 (2)	0.22847 (7)	0.1112 (9)
C12	0.41462 (12)	0.2754 (2)	0.37992 (6)	0.0718 (5)
C13	0.42824 (16)	0.8503 (2)	0.49709 (6)	0.0871 (7)
C14	-0.31722 (17)	-1.1411 (2)	0.16706 (6)	0.0858 (6)
C15	-0.38786 (12)	-0.7789 (2)	0.29976 (6)	0.0696 (5)
C16	-0.45426 (14)	-1.3607 (2)	0.39707 (6)	0.0814 (6)
O1	0.0459 (3)	-0.1244 (5)	0.43492 (13)	0.0571 (11)
O2	0.0428 (3)	-0.1317 (5)	0.35970 (14)	0.0613 (12)
H2A	0.0228	-0.2219	0.3629	0.092*
O3	-0.0346 (3)	-0.4099 (5)	0.42547 (14)	0.0648 (13)
H3B	-0.0097	-0.3220	0.4233	0.097*
O4	-0.0241 (3)	-0.4055 (5)	0.35213 (14)	0.0636 (12)
N1	0.1830 (3)	0.2661 (5)	0.37220 (15)	0.0398 (11)
N2	0.1338 (3)	0.1348 (5)	0.36084 (15)	0.0442 (12)
N3	-0.1099 (3)	-0.6749 (5)	0.33613 (15)	0.0414 (11)
N4	-0.1607 (3)	-0.8069 (5)	0.33785 (14)	0.0387 (11)
C1	0.2820 (7)	0.5083 (7)	0.2692 (2)	0.065 (2)
C2	0.1789 (7)	0.4874 (8)	0.2669 (2)	0.071 (2)
H2B	0.1303	0.5256	0.2423	0.085*
C3	0.1464 (5)	0.4081 (7)	0.3017 (2)	0.0626 (18)
H3A	0.0759	0.3967	0.3009	0.075*
C4	0.2182 (4)	0.3469 (6)	0.33725 (18)	0.0435 (14)
C5	0.3238 (5)	0.3634 (7)	0.3377 (2)	0.0519 (16)
C6	0.3555 (6)	0.4459 (7)	0.3041 (2)	0.0642 (19)
H6A	0.4258	0.4592	0.3049	0.077*
C7	0.3576 (5)	0.6908 (7)	0.4752 (2)	0.0504 (16)
C8	0.2851 (5)	0.7026 (7)	0.4350 (2)	0.0496 (16)
H8A	0.2715	0.7959	0.4205	0.060*
C9	0.2321 (4)	0.5733 (6)	0.4163 (2)	0.0446 (14)
H9A	0.1836	0.5801	0.3888	0.053*
C10	0.2507 (4)	0.4355 (6)	0.43793 (19)	0.0393 (13)
C11	0.3227 (4)	0.4274 (7)	0.4789 (2)	0.0525 (16)
H11A	0.3356	0.3349	0.4940	0.063*
C12	0.3753 (5)	0.5553 (8)	0.4976 (2)	0.0563 (17)
H12A	0.4226	0.5494	0.5254	0.068*

C13	0.1975 (4)	0.2945 (6)	0.41780 (18)	0.0399 (13)
C14	0.1523 (4)	0.1770 (6)	0.43595 (19)	0.0427 (14)
C15	0.1141 (4)	0.0811 (6)	0.39951 (18)	0.0412 (14)
C16	0.1457 (5)	0.1614 (7)	0.4862 (2)	0.0612 (18)
H16A	0.1098	0.0694	0.4902	0.092*
H16B	0.1086	0.2465	0.4948	0.092*
H16C	0.2144	0.1587	0.5050	0.092*
C17	0.0649 (4)	-0.0666 (7)	0.3986 (2)	0.0462 (15)
C18	-0.2676 (5)	-1.0395 (7)	0.21641 (19)	0.0508 (16)
C19	-0.3380 (5)	-0.9630 (7)	0.2361 (2)	0.0519 (16)
H19A	-0.4084	-0.9654	0.2227	0.062*
C20	-0.3033 (4)	-0.8837 (6)	0.27553 (19)	0.0435 (14)
C21	-0.1980 (4)	-0.8850 (6)	0.29602 (18)	0.0405 (14)
C22	-0.1292 (5)	-0.9617 (7)	0.2757 (2)	0.0549 (17)
H22A	-0.0588	-0.9621	0.2892	0.066*
C23	-0.1643 (5)	-1.0374 (7)	0.2354 (2)	0.0570 (17)
H23A	-0.1177	-1.0871	0.2212	0.068*
C24	-0.3712 (5)	-1.2088 (8)	0.3953 (2)	0.0524 (16)
C25	-0.3943 (5)	-1.0715 (8)	0.4093 (2)	0.0643 (19)
H25A	-0.4533	-1.0587	0.4212	0.077*
C26	-0.3309 (5)	-0.9481 (7)	0.4061 (2)	0.0578 (17)
H26A	-0.3466	-0.8535	0.4165	0.069*
C27	-0.2438 (4)	-0.9654 (7)	0.38723 (19)	0.0430 (14)
C28	-0.2201 (4)	-1.1102 (7)	0.37496 (19)	0.0500 (16)
H28A	-0.1604	-1.1255	0.3637	0.060*
C29	-0.2827 (5)	-1.2322 (7)	0.3790 (2)	0.0554 (17)
H29A	-0.2654	-1.3290	0.3707	0.067*
C30	-0.1821 (4)	-0.8338 (6)	0.38025 (18)	0.0380 (13)
C31	-0.1414 (4)	-0.7112 (6)	0.40758 (17)	0.0373 (13)
C32	-0.0975 (4)	-0.6191 (6)	0.37874 (18)	0.0386 (13)
C33	-0.1414 (5)	-0.6905 (7)	0.45721 (18)	0.0589 (18)
H33A	-0.1783	-0.7728	0.4674	0.088*
H33B	-0.1748	-0.5964	0.4614	0.088*
H33C	-0.0712	-0.6891	0.4746	0.088*
C34	-0.0488 (4)	-0.4708 (7)	0.3863 (2)	0.0466 (15)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.190 (3)	0.0800 (15)	0.0846 (15)	-0.0236 (16)	0.0770 (16)	0.0149 (12)
Cl2	0.0473 (10)	0.0823 (13)	0.0861 (13)	0.0030 (9)	0.0155 (9)	0.0118 (10)
Cl3	0.1092 (16)	0.0820 (14)	0.0773 (13)	-0.0542 (12)	0.0357 (11)	-0.0305 (11)
Cl4	0.1134 (16)	0.0851 (14)	0.0521 (11)	-0.0172 (12)	0.0026 (10)	-0.0250 (10)
Cl5	0.0472 (10)	0.0915 (13)	0.0668 (11)	0.0158 (10)	0.0052 (8)	-0.0112 (10)
Cl6	0.0785 (13)	0.0780 (13)	0.0838 (13)	-0.0437 (11)	0.0093 (10)	0.0142 (11)
O1	0.070 (3)	0.057 (3)	0.044 (2)	-0.018 (2)	0.013 (2)	0.008 (2)
O2	0.076 (3)	0.048 (3)	0.056 (3)	-0.028 (2)	0.007 (2)	-0.002 (2)
O3	0.074 (3)	0.064 (3)	0.059 (3)	-0.028 (2)	0.018 (2)	-0.022 (2)

O4	0.077 (3)	0.055 (3)	0.060 (3)	-0.023 (2)	0.019 (2)	-0.010 (2)
N1	0.043 (3)	0.032 (3)	0.046 (3)	-0.012 (2)	0.012 (2)	0.006 (2)
N2	0.041 (3)	0.046 (3)	0.045 (3)	-0.007 (2)	0.008 (2)	-0.009 (2)
N3	0.036 (3)	0.038 (3)	0.048 (3)	-0.015 (2)	0.004 (2)	-0.005 (2)
N4	0.035 (3)	0.041 (3)	0.039 (3)	-0.009 (2)	0.007 (2)	-0.002 (2)
C1	0.107 (6)	0.043 (4)	0.055 (4)	-0.005 (4)	0.037 (4)	0.003 (3)
C2	0.099 (6)	0.069 (5)	0.042 (4)	-0.004 (5)	0.008 (4)	0.006 (3)
C3	0.072 (5)	0.067 (5)	0.050 (4)	-0.002 (4)	0.016 (4)	0.008 (4)
C4	0.049 (4)	0.036 (3)	0.043 (3)	-0.009 (3)	0.005 (3)	0.002 (3)
C5	0.047 (4)	0.048 (4)	0.061 (4)	-0.006 (3)	0.013 (3)	0.003 (3)
C6	0.072 (5)	0.055 (4)	0.078 (5)	-0.017 (4)	0.044 (4)	-0.013 (4)
C7	0.052 (4)	0.057 (4)	0.047 (4)	-0.027 (3)	0.021 (3)	-0.014 (3)
C8	0.062 (4)	0.035 (3)	0.060 (4)	-0.002 (3)	0.031 (3)	-0.001 (3)
C9	0.037 (3)	0.044 (4)	0.052 (4)	0.001 (3)	0.009 (3)	0.006 (3)
C10	0.036 (3)	0.039 (3)	0.046 (3)	-0.007 (3)	0.015 (3)	-0.002 (3)
C11	0.046 (4)	0.048 (4)	0.061 (4)	-0.013 (3)	0.005 (3)	0.002 (3)
C12	0.045 (4)	0.073 (5)	0.049 (4)	-0.019 (4)	0.006 (3)	-0.004 (4)
C13	0.029 (3)	0.052 (4)	0.039 (3)	-0.004 (3)	0.007 (2)	0.006 (3)
C14	0.034 (3)	0.049 (4)	0.046 (3)	-0.001 (3)	0.010 (3)	0.003 (3)
C15	0.039 (3)	0.044 (3)	0.039 (3)	-0.006 (3)	0.006 (3)	-0.008 (3)
C16	0.078 (5)	0.054 (4)	0.057 (4)	-0.013 (4)	0.027 (3)	0.002 (3)
C17	0.034 (3)	0.049 (4)	0.050 (4)	-0.003 (3)	-0.004 (3)	0.004 (3)
C18	0.070 (5)	0.053 (4)	0.031 (3)	-0.007 (4)	0.013 (3)	-0.007 (3)
C19	0.044 (4)	0.060 (4)	0.045 (4)	-0.007 (3)	-0.006 (3)	-0.001 (3)
C20	0.038 (3)	0.049 (4)	0.044 (3)	0.000 (3)	0.008 (3)	0.012 (3)
C21	0.038 (3)	0.047 (3)	0.036 (3)	-0.010 (3)	0.007 (3)	-0.003 (3)
C22	0.044 (4)	0.077 (5)	0.045 (4)	-0.008 (3)	0.012 (3)	-0.016 (3)
C23	0.057 (5)	0.065 (5)	0.054 (4)	0.000 (4)	0.024 (3)	-0.009 (3)
C24	0.052 (4)	0.059 (4)	0.044 (4)	-0.020 (3)	0.004 (3)	0.020 (3)
C25	0.053 (4)	0.064 (5)	0.083 (5)	-0.016 (4)	0.030 (4)	0.000 (4)
C26	0.053 (4)	0.056 (4)	0.074 (5)	-0.010 (3)	0.033 (4)	-0.005 (3)
C27	0.033 (3)	0.048 (4)	0.044 (3)	-0.005 (3)	0.001 (3)	0.006 (3)
C28	0.038 (4)	0.056 (4)	0.056 (4)	-0.007 (3)	0.009 (3)	0.005 (3)
C29	0.063 (4)	0.046 (4)	0.058 (4)	-0.013 (3)	0.013 (3)	0.003 (3)
C30	0.032 (3)	0.042 (3)	0.040 (3)	0.000 (3)	0.006 (2)	-0.002 (3)
C31	0.037 (3)	0.043 (3)	0.032 (3)	-0.004 (3)	0.007 (2)	-0.007 (3)
C32	0.031 (3)	0.040 (3)	0.044 (3)	-0.010 (3)	0.007 (3)	-0.002 (3)
C33	0.072 (4)	0.064 (4)	0.045 (4)	-0.011 (4)	0.022 (3)	-0.008 (3)
C34	0.038 (4)	0.056 (4)	0.046 (4)	-0.003 (3)	0.010 (3)	-0.001 (3)

Geometric parameters (Å, °)

C11—C1	1.717 (7)	C11—C12	1.378 (8)
C12—C5	1.729 (6)	C11—H11A	0.9300
C13—C7	1.737 (6)	C12—H12A	0.9300
C14—C18	1.735 (6)	C13—C14	1.366 (7)
C15—C20	1.725 (6)	C14—C15	1.388 (7)
C16—C24	1.739 (6)	C14—C16	1.536 (7)

O1—C17	1.277 (6)	C15—C17	1.453 (8)
O2—C17	1.276 (6)	C16—H16A	0.9600
O2—H2A	0.8500	C16—H16B	0.9600
O3—C34	1.269 (6)	C16—H16C	0.9600
O3—H3B	0.8500	C18—C23	1.360 (8)
O4—C34	1.279 (6)	C18—C19	1.379 (8)
N1—N2	1.336 (6)	C19—C20	1.365 (8)
N1—C13	1.364 (6)	C19—H19A	0.9300
N1—C4	1.426 (6)	C20—C21	1.393 (7)
N2—C15	1.329 (6)	C21—C22	1.376 (7)
N3—C32	1.347 (6)	C22—C23	1.370 (8)
N3—N4	1.349 (6)	C22—H22A	0.9300
N4—C30	1.382 (6)	C23—H23A	0.9300
N4—C21	1.424 (6)	C24—C25	1.337 (9)
C1—C2	1.360 (9)	C24—C29	1.376 (8)
C1—C6	1.379 (9)	C25—C26	1.388 (8)
C2—C3	1.401 (8)	C25—H25A	0.9300
C2—H2B	0.9300	C26—C27	1.394 (7)
C3—C4	1.374 (8)	C26—H26A	0.9300
C3—H3A	0.9300	C27—C28	1.383 (8)
C4—C5	1.397 (8)	C27—C30	1.458 (7)
C5—C6	1.380 (8)	C28—C29	1.377 (8)
C6—H6A	0.9300	C28—H28A	0.9300
C7—C12	1.366 (8)	C29—H29A	0.9300
C7—C8	1.372 (8)	C30—C31	1.393 (7)
C8—C9	1.391 (8)	C31—C32	1.400 (7)
C8—H8A	0.9300	C31—C33	1.501 (7)
C9—C10	1.375 (7)	C32—C34	1.452 (8)
C9—H9A	0.9300	C33—H33A	0.9600
C10—C11	1.384 (7)	C33—H33B	0.9600
C10—C13	1.490 (7)	C33—H33C	0.9600
C17—O2—H2A	109.6	H16A—C16—H16C	109.5
C34—O3—H3B	107.3	H16B—C16—H16C	109.5
N2—N1—C13	111.8 (4)	O1—C17—O2	123.6 (5)
N2—N1—C4	117.6 (4)	O1—C17—C15	120.7 (6)
C13—N1—C4	130.5 (5)	O2—C17—C15	115.7 (5)
C15—N2—N1	105.1 (4)	C23—C18—C19	121.3 (6)
C32—N3—N4	103.9 (4)	C23—C18—C14	121.7 (5)
N3—N4—C30	112.6 (4)	C19—C18—C14	117.0 (5)
N3—N4—C21	117.6 (4)	C20—C19—C18	119.4 (6)
C30—N4—C21	129.0 (5)	C20—C19—H19A	120.3
C2—C1—C6	121.2 (6)	C18—C19—H19A	120.3
C2—C1—C11	120.3 (6)	C19—C20—C21	119.7 (5)
C6—C1—C11	118.5 (6)	C19—C20—C15	120.8 (5)
C1—C2—C3	119.5 (7)	C21—C20—C15	119.5 (4)
C1—C2—H2B	120.3	C22—C21—C20	119.9 (5)
C3—C2—H2B	120.3	C22—C21—N4	119.6 (5)

C4—C3—C2	120.3 (7)	C20—C21—N4	120.5 (5)
C4—C3—H3A	119.8	C23—C22—C21	120.0 (6)
C2—C3—H3A	119.8	C23—C22—H22A	120.0
C3—C4—C5	119.1 (5)	C21—C22—H22A	120.0
C3—C4—N1	119.2 (5)	C18—C23—C22	119.7 (6)
C5—C4—N1	121.7 (5)	C18—C23—H23A	120.2
C6—C5—C4	120.4 (6)	C22—C23—H23A	120.2
C6—C5—C12	120.0 (5)	C25—C24—C29	121.1 (6)
C4—C5—C12	119.5 (5)	C25—C24—C16	120.0 (5)
C1—C6—C5	119.4 (6)	C29—C24—C16	118.9 (6)
C1—C6—H6A	120.3	C24—C25—C26	120.4 (6)
C5—C6—H6A	120.3	C24—C25—H25A	119.8
C12—C7—C8	120.8 (5)	C26—C25—H25A	119.8
C12—C7—C13	120.1 (5)	C25—C26—C27	120.2 (6)
C8—C7—C13	119.1 (5)	C25—C26—H26A	119.9
C7—C8—C9	119.1 (6)	C27—C26—H26A	119.9
C7—C8—H8A	120.4	C28—C27—C26	117.6 (6)
C9—C8—H8A	120.4	C28—C27—C30	122.1 (5)
C10—C9—C8	120.7 (5)	C26—C27—C30	120.4 (6)
C10—C9—H9A	119.7	C29—C28—C27	121.6 (6)
C8—C9—H9A	119.7	C29—C28—H28A	119.2
C9—C10—C11	118.9 (5)	C27—C28—H28A	119.2
C9—C10—C13	121.6 (5)	C28—C29—C24	119.0 (6)
C11—C10—C13	119.4 (5)	C28—C29—H29A	120.5
C12—C11—C10	120.5 (6)	C24—C29—H29A	120.5
C12—C11—H11A	119.7	N4—C30—C31	106.3 (5)
C10—C11—H11A	119.7	N4—C30—C27	120.0 (5)
C7—C12—C11	119.8 (6)	C31—C30—C27	133.6 (5)
C7—C12—H12A	120.1	C30—C31—C32	104.2 (4)
C11—C12—H12A	120.1	C30—C31—C33	126.6 (5)
N1—C13—C14	106.4 (5)	C32—C31—C33	129.2 (5)
N1—C13—C10	120.8 (5)	N3—C32—C31	113.1 (5)
C14—C13—C10	132.8 (5)	N3—C32—C34	115.9 (5)
C13—C14—C15	105.3 (5)	C31—C32—C34	130.8 (5)
C13—C14—C16	125.6 (5)	C31—C33—H33A	109.5
C15—C14—C16	129.2 (5)	C31—C33—H33B	109.5
N2—C15—C14	111.4 (5)	H33A—C33—H33B	109.5
N2—C15—C17	118.2 (5)	C31—C33—H33C	109.5
C14—C15—C17	130.3 (5)	H33A—C33—H33C	109.5
C14—C16—H16A	109.5	H33B—C33—H33C	109.5
C14—C16—H16B	109.5	O3—C34—O4	122.7 (6)
H16A—C16—H16B	109.5	O3—C34—C32	119.9 (5)
C14—C16—H16C	109.5	O4—C34—C32	117.4 (5)
C13—N1—N2—C15	1.7 (6)	C32—N3—N4—C30	1.1 (6)
C4—N1—N2—C15	177.3 (5)	C32—N3—N4—C21	171.9 (4)
C6—C1—C2—C3	3.5 (10)	C23—C18—C19—C20	-0.1 (9)
C11—C1—C2—C3	-176.6 (5)	C14—C18—C19—C20	178.7 (5)

C1—C2—C3—C4	-2.4 (10)	C18—C19—C20—C21	-2.0 (9)
C2—C3—C4—C5	-0.8 (9)	C18—C19—C20—C15	176.8 (4)
C2—C3—C4—N1	-179.7 (6)	C19—C20—C21—C22	2.2 (9)
N2—N1—C4—C3	67.3 (7)	C15—C20—C21—C22	-176.6 (5)
C13—N1—C4—C3	-118.2 (6)	C19—C20—C21—N4	-178.1 (5)
N2—N1—C4—C5	-111.7 (6)	C15—C20—C21—N4	3.2 (7)
C13—N1—C4—C5	62.9 (8)	N3—N4—C21—C22	74.7 (7)
C3—C4—C5—C6	2.9 (9)	C30—N4—C21—C22	-116.2 (6)
N1—C4—C5—C6	-178.2 (5)	N3—N4—C21—C20	-105.0 (6)
C3—C4—C5—C12	-175.3 (5)	C30—N4—C21—C20	64.1 (8)
N1—C4—C5—C12	3.6 (8)	C20—C21—C22—C23	-0.4 (9)
C2—C1—C6—C5	-1.4 (10)	N4—C21—C22—C23	179.9 (5)
C11—C1—C6—C5	178.6 (5)	C19—C18—C23—C22	1.9 (10)
C4—C5—C6—C1	-1.8 (10)	C14—C18—C23—C22	-176.8 (5)
C12—C5—C6—C1	176.3 (5)	C21—C22—C23—C18	-1.7 (10)
C12—C7—C8—C9	-2.5 (9)	C29—C24—C25—C26	2.2 (10)
C13—C7—C8—C9	176.9 (4)	C16—C24—C25—C26	-177.1 (5)
C7—C8—C9—C10	1.0 (8)	C24—C25—C26—C27	1.6 (10)
C8—C9—C10—C11	0.4 (8)	C25—C26—C27—C28	-4.2 (9)
C8—C9—C10—C13	-178.0 (5)	C25—C26—C27—C30	174.8 (6)
C9—C10—C11—C12	-0.4 (9)	C26—C27—C28—C29	3.2 (9)
C13—C10—C11—C12	178.1 (5)	C30—C27—C28—C29	-175.7 (5)
C8—C7—C12—C11	2.6 (9)	C27—C28—C29—C24	0.4 (9)
C13—C7—C12—C11	-176.8 (5)	C25—C24—C29—C28	-3.1 (9)
C10—C11—C12—C7	-1.1 (9)	C16—C24—C29—C28	176.1 (4)
N2—N1—C13—C14	-1.8 (6)	N3—N4—C30—C31	-0.5 (6)
C4—N1—C13—C14	-176.6 (5)	C21—N4—C30—C31	-170.0 (5)
N2—N1—C13—C10	-179.9 (4)	N3—N4—C30—C27	175.2 (4)
C4—N1—C13—C10	5.3 (8)	C21—N4—C30—C27	5.7 (8)
C9—C10—C13—N1	40.9 (8)	C28—C27—C30—N4	51.7 (7)
C11—C10—C13—N1	-137.5 (5)	C26—C27—C30—N4	-127.2 (6)
C9—C10—C13—C14	-136.6 (6)	C28—C27—C30—C31	-134.0 (7)
C11—C10—C13—C14	45.0 (9)	C26—C27—C30—C31	47.1 (9)
N1—C13—C14—C15	1.1 (6)	N4—C30—C31—C32	-0.3 (6)
C10—C13—C14—C15	178.8 (6)	C27—C30—C31—C32	-175.1 (6)
N1—C13—C14—C16	-177.9 (5)	N4—C30—C31—C33	-177.6 (5)
C10—C13—C14—C16	-0.1 (10)	C27—C30—C31—C33	7.6 (10)
N1—N2—C15—C14	-1.0 (6)	N4—N3—C32—C31	-1.3 (6)
N1—N2—C15—C17	-177.0 (5)	N4—N3—C32—C34	-176.9 (4)
C13—C14—C15—N2	0.0 (6)	C30—C31—C32—N3	1.0 (6)
C16—C14—C15—N2	178.9 (5)	C33—C31—C32—N3	178.2 (5)
C13—C14—C15—C17	175.3 (5)	C30—C31—C32—C34	175.7 (5)
C16—C14—C15—C17	-5.8 (10)	C33—C31—C32—C34	-7.1 (10)
N2—C15—C17—O1	178.8 (5)	N3—C32—C34—O3	179.8 (5)
C14—C15—C17—O1	3.8 (9)	C31—C32—C34—O3	5.2 (9)
N2—C15—C17—O2	-1.3 (8)	N3—C32—C34—O4	1.3 (7)
C14—C15—C17—O2	-176.3 (6)	C31—C32—C34—O4	-173.3 (6)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2 <i>A</i> \cdots O4	0.85	1.74	2.564 (7)	163
O3—H3 <i>B</i> \cdots O1	0.85	1.89	2.723 (6)	165
C16—H16 <i>C</i> \cdots C <i>g</i> 6 ⁱ	0.96	3.13	3.867 (4)	135
C33—H33 <i>B</i> \cdots C <i>g</i> 4 ⁱⁱ	0.96	3.29	3.857 (3)	120

Symmetry codes: (i) $-x+1, -y, -z$; (ii) $-x+1, -y+1, -z$.