

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Amino-1*H*-benzoimidazol-3-ium 4,4,4-trifluoro-1,3-dioxo-1-phenylbutan-2-ideGong-Chun Li,^{a*} Feng-Ling Yang^a and Chang-Sheng Yao^b

^aCollege of Chemistry and Chemical Engineering, Xuchang University, Xuchang, Henan Province 461000, People's Republic of China, and ^bSchool of Chemistry and Chemical Engineering, and Key Laboratory of Biotechnology for Medicinal Plants, Xuzhou Normal University, Xuzhou 221116, People's Republic of China
Correspondence e-mail: actaeli@mail.google.com


Received 12 October 2008; accepted 12 November 2008

Key indicators: single-crystal X-ray study; $T = 113$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.039; wR factor = 0.107; data-to-parameter ratio = 11.9.

In the title compound, $\text{C}_7\text{H}_8\text{N}_3^+ \cdot \text{C}_{10}\text{H}_6\text{F}_3\text{O}_2^-$, 1*H*-benzoimidazol-2-amine system adopts a planar conformation with an r.m.s. deviation of 0.0174 Å. The cation and anion in the asymmetric unit are linked by $\text{N}-\text{H} \cdots \text{O}$ hydrogen bonds. There are also additional intermolecular $\text{N}-\text{H} \cdots \text{O}$ hydrogen bonds and $\pi-\pi$ stacking interactions between the phenyl rings of neighbouring anions with centroid-centroid distances of 4.0976 (13) Å.

Related literature

For details of the bioactivity of organofluorine compounds, see: Hermann *et al.* (2003); Ulrich (2004).



Experimental

Crystal data

$\text{C}_7\text{H}_8\text{N}_3^+ \cdot \text{C}_{10}\text{H}_6\text{F}_3\text{O}_2^-$
 $M_r = 349.31$

Orthorhombic, $Pbca$
 $a = 16.755$ (3) Å

$b = 10.552$ (2) Å
 $c = 18.497$ (4) Å
 $V = 3270.0$ (11) Å³
 $Z = 8$

Mo $K\alpha$ radiation
 $\mu = 0.12$ mm⁻¹
 $T = 113$ (2) K
 $0.12 \times 0.10 \times 0.06$ mm

Data collection

Rigaku Saturn CCD diffractometer
Absorption correction: multi-scan
(*CrystalClear*; Rigaku/MS, 2002)
 $T_{\min} = 0.986$, $T_{\max} = 0.993$

31132 measured reflections
2881 independent reflections
2609 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.050$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.107$
 $S = 1.06$
2881 reflections
243 parameters
5 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\max} = 0.21$ e Å⁻³
 $\Delta\rho_{\min} = -0.20$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N1}-\text{H1A} \cdots \text{O2}^{\ddagger}$	0.877 (9)	2.004 (11)	2.7803 (18)	146.8 (16)
$\text{N3}-\text{H3A} \cdots \text{O1}^{\ddagger}$	0.895 (9)	1.949 (13)	2.7651 (17)	150.9 (19)
$\text{N3}-\text{H3A} \cdots \text{O2}^{\ddagger}$	0.895 (9)	2.363 (18)	2.9912 (18)	127.3 (17)
$\text{N1}-\text{H1B} \cdots \text{O1}$	0.879 (9)	2.030 (10)	2.8662 (18)	158.4 (16)
$\text{N2}-\text{H2A} \cdots \text{O2}$	0.905 (10)	1.862 (11)	2.7360 (18)	161.8 (19)

Symmetry code: (i) $-x + 2, y - \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku/MS, 2002); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the Natural Science Foundation of Henan Province, China (grant No. 082300420110) and the Natural Science Foundation of Henan Province Education Department, China (grant No. 2007150036) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2549).

References

- Hermann, B., Erwin, H. & Hansjorg, K. (2003). US Patent No. 2 003 176 284. Rigaku/MS (2002). *CrystalClear*. Rigaku/MS Inc., The Woodlands, Texas, USA.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Ulrich, H. (2004). US Patent No. 2 004 033 897.

supporting information

Acta Cryst. (2008). E64, o2460 [doi:10.1107/S1600536808037483]

2-Amino-1*H*-benzoimidazol-3-ium 4,4,4-trifluoro-1,3-dioxo-1-phenylbutan-2-ide

Gong-Chun Li, Feng-Ling Yang and Chang-Sheng Yao

S1. Comment

Compounds that contain fluorine exhibit particular bioactivity, for example, flumioxazin is a widely used herbicide (Hermann *et al.*, 2003; Ulrich, 2004). This led us to pay much attention to the synthesis and structure of similar fluoro-compounds. During the synthesis of trifluoromethylated heterocyclic compounds, an intermediate, the title compound, (I), Fig. 1, was isolated and we report its crystal structure here.

The 1*H*-benzo[*d*]imidazol-2-amine cation adopts a planar conformation with an rms deviation of 0.0174 for the fitted atoms. The phenyl ring is almost perpendicular to the fused heterocyclic rings, with a dihedral angle of 81.84 (4)° between them. The cation and anion in the asymmetric unit are linked by N1—H1B···O1 and N2—H2A···O2 hydrogen bonds.

The crystal packing is further stabilized by additional intermolecular N—H···O hydrogen bonds (Table 1, Fig. 2) and intermolecular $\pi\cdots\pi$ stacking interactions between the C5-C10 phenyl rings (symmetry code: 2-x, -y, -z) of neighbouring anions with centroid-to-centroid distances, plane-plane distances and displacement distances of 4.0976 (13), 3.786 and 1.565 Å respectively.

S2. Experimental

The title compound was synthesized by the reaction of 4,4,4-trifluoro-1-phenylbutane-1,3-dione (1 mmol) and 1*H*-benzo[*d*]imidazol-2-amine (1 mmol) in refluxing ethanol (20 mL) for a certain time (monitored by TLC). Cooling, the reaction mixture slowly to room temperature, gave single crystals suitable for X-ray diffraction.

S3. Refinement

Hydrogen atoms bound to nitrogen atoms were located in a difference Fourier map and were refined with d(N—H) restrained to 0.90 (1) Å and d(H1A···H1B) restrained to 1.50 (1) Å. Other H atoms were placed in calculated positions, d(C—H) = 0.93 Å, and included in the final cycles of refinement using a riding model, with $U_{iso}(H) = 1.2U_{eq}(C)$.




Figure 1

The asymmetric unit of the title compound, showing 30% probability displacement ellipsoids and the atom-numbering scheme. Hydrogen bonds are drawn as dashed lines.




Figure 2

The packing diagram of the title compound. Intermolecular hydrogen bonds are shown as dashed lines.

2-Amino-1*H*-benzimidazol-3-ium 4,4,4-trifluoro-1,3-dioxo-1-phenylbutan-2-ide

Crystal data

$C_7H_8N_3^+ \cdot C_{10}H_6F_3O_2^-$

$M_r = 349.31$

Orthorhombic, *Pbca*

Hall symbol: $-P\ 2ac\ 2ab$

$a = 16.755\ (3)\ \text{\AA}$

$b = 10.552\ (2)\ \text{\AA}$

$c = 18.497\ (4)\ \text{\AA}$

$V = 3270.0\ (11)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1440$

$D_x = 1.419 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 6974 reflections
 $\theta = 2.2\text{--}27.9^\circ$

$\mu = 0.12 \text{ mm}^{-1}$
 $T = 113 \text{ K}$
 Block, yellow
 $0.12 \times 0.10 \times 0.06 \text{ mm}$

Data collection

Rigaku Saturn CCD
 diffractometer
 Radiation source: rotating anode
 Confocal monochromator
 ω scans
 Absorption correction: multi-scan
 (*CrystalClear*; Rigaku/MSC, 2002)
 $T_{\min} = 0.986$, $T_{\max} = 0.993$

31132 measured reflections
 2881 independent reflections
 2609 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.050$
 $\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.2^\circ$
 $h = -19 \rightarrow 19$
 $k = -12 \rightarrow 12$
 $l = -22 \rightarrow 21$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.107$
 $S = 1.06$
 2881 reflections
 243 parameters
 5 restraints
 Primary atom site location: structure-invariant
 direct methods
 Secondary atom site location: difference Fourier
 map

Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0613P)^2 + 0.895P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.21 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.20 \text{ e \AA}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick,
 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
 Extinction coefficient: 0.0073 (10)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
F1	0.74257 (7)	0.54639 (12)	0.05657 (5)	0.0552 (3)
F2	0.77552 (6)	0.66729 (9)	0.14532 (6)	0.0473 (3)
F3	0.71669 (6)	0.49082 (11)	0.16580 (6)	0.0467 (3)
O1	1.01117 (6)	0.37243 (10)	0.11875 (6)	0.0301 (3)
O2	0.88568 (6)	0.49842 (10)	0.19269 (6)	0.0303 (3)
N1	1.03832 (8)	0.19158 (15)	0.23172 (8)	0.0379 (4)
N2	0.90662 (8)	0.26462 (13)	0.25377 (7)	0.0298 (3)
N3	0.94668 (8)	0.09546 (13)	0.31266 (7)	0.0295 (3)
C1	0.77211 (10)	0.54628 (16)	0.12397 (9)	0.0335 (4)
C2	0.85307 (9)	0.48052 (14)	0.13136 (8)	0.0277 (3)

C3	0.88043 (9)	0.40780 (15)	0.07420 (8)	0.0290 (4)
H3	0.8467	0.3955	0.0349	0.035*
C4	0.95774 (9)	0.35055 (14)	0.07232 (8)	0.0264 (3)
C5	0.97828 (9)	0.26266 (14)	0.01141 (8)	0.0263 (3)
C6	1.05759 (10)	0.25313 (16)	-0.01066 (9)	0.0324 (4)
H6	1.0962	0.3030	0.0116	0.039*
C7	1.07968 (10)	0.17084 (17)	-0.06508 (10)	0.0378 (4)
H7	1.1327	0.1665	-0.0797	0.045*
C8	1.02270 (11)	0.09443 (16)	-0.09797 (9)	0.0368 (4)
H8	1.0376	0.0379	-0.1341	0.044*
C9	0.94389 (10)	0.10278 (15)	-0.07679 (9)	0.0346 (4)
H9	0.9057	0.0518	-0.0989	0.042*
C10	0.92111 (10)	0.18696 (15)	-0.02263 (8)	0.0308 (4)
H10	0.8678	0.1928	-0.0091	0.037*
C11	0.96806 (9)	0.18451 (15)	0.26443 (8)	0.0296 (4)
C12	0.84329 (9)	0.22790 (15)	0.29854 (8)	0.0275 (4)
C13	0.76879 (9)	0.28105 (17)	0.31021 (8)	0.0321 (4)
H13	0.7518	0.3527	0.2853	0.038*
C14	0.72078 (10)	0.22158 (17)	0.36124 (9)	0.0349 (4)
H14	0.6705	0.2548	0.3710	0.042*
C15	0.74579 (10)	0.11347 (17)	0.39833 (9)	0.0339 (4)
H15	0.7118	0.0763	0.4319	0.041*
C16	0.82038 (10)	0.06027 (15)	0.38614 (8)	0.0309 (4)
H16	0.8371	-0.0120	0.4106	0.037*
C17	0.86862 (9)	0.11999 (14)	0.33575 (8)	0.0273 (3)
H1A	1.0783 (9)	0.1439 (16)	0.2457 (10)	0.046 (5)*
H1B	1.0438 (10)	0.2461 (14)	0.1961 (8)	0.043 (5)*
H2A	0.9093 (12)	0.3377 (13)	0.2283 (10)	0.051 (6)*
H3A	0.9769 (11)	0.0317 (15)	0.3290 (11)	0.051 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F1	0.0509 (7)	0.0792 (8)	0.0356 (6)	0.0313 (6)	-0.0129 (5)	-0.0053 (6)
F2	0.0462 (6)	0.0331 (6)	0.0628 (7)	0.0093 (4)	0.0013 (5)	-0.0034 (5)
F3	0.0293 (5)	0.0561 (7)	0.0546 (7)	-0.0022 (5)	0.0024 (5)	0.0043 (5)
O1	0.0292 (6)	0.0319 (6)	0.0291 (6)	-0.0017 (5)	-0.0052 (5)	-0.0008 (5)
O2	0.0307 (6)	0.0322 (6)	0.0279 (6)	-0.0019 (5)	-0.0042 (4)	-0.0015 (5)
N1	0.0316 (8)	0.0461 (9)	0.0358 (8)	0.0102 (7)	0.0039 (6)	0.0122 (7)
N2	0.0306 (7)	0.0313 (7)	0.0274 (7)	0.0048 (6)	-0.0006 (5)	0.0055 (6)
N3	0.0298 (7)	0.0304 (7)	0.0284 (7)	0.0056 (6)	-0.0008 (5)	0.0029 (6)
C1	0.0353 (9)	0.0355 (9)	0.0296 (8)	0.0026 (7)	-0.0016 (7)	0.0005 (7)
C2	0.0275 (8)	0.0253 (8)	0.0301 (8)	-0.0030 (6)	-0.0033 (6)	0.0048 (6)
C3	0.0295 (8)	0.0312 (8)	0.0263 (8)	0.0009 (6)	-0.0056 (6)	0.0006 (6)
C4	0.0296 (8)	0.0239 (8)	0.0256 (7)	-0.0038 (6)	-0.0027 (6)	0.0053 (6)
C5	0.0291 (8)	0.0251 (7)	0.0249 (7)	-0.0002 (6)	-0.0030 (6)	0.0044 (6)
C6	0.0285 (8)	0.0337 (9)	0.0350 (9)	-0.0024 (7)	-0.0036 (7)	-0.0011 (7)
C7	0.0309 (9)	0.0418 (10)	0.0407 (9)	0.0010 (7)	0.0045 (7)	-0.0035 (8)

C8	0.0452 (10)	0.0324 (9)	0.0327 (9)	0.0025 (7)	0.0014 (7)	-0.0036 (7)
C9	0.0391 (9)	0.0286 (8)	0.0360 (9)	-0.0057 (7)	-0.0058 (7)	-0.0025 (7)
C10	0.0287 (8)	0.0306 (8)	0.0330 (8)	-0.0030 (6)	-0.0021 (6)	0.0020 (7)
C11	0.0301 (9)	0.0336 (8)	0.0250 (8)	0.0036 (7)	-0.0028 (6)	0.0009 (6)
C12	0.0293 (8)	0.0305 (8)	0.0227 (7)	-0.0001 (6)	-0.0027 (6)	-0.0021 (6)
C13	0.0300 (9)	0.0369 (9)	0.0292 (8)	0.0055 (7)	-0.0047 (6)	-0.0011 (7)
C14	0.0260 (8)	0.0453 (10)	0.0332 (9)	0.0007 (7)	-0.0029 (6)	-0.0045 (8)
C15	0.0300 (8)	0.0421 (10)	0.0298 (8)	-0.0074 (7)	-0.0005 (7)	-0.0015 (7)
C16	0.0348 (9)	0.0300 (8)	0.0279 (8)	-0.0032 (7)	-0.0049 (6)	0.0018 (6)
C17	0.0279 (8)	0.0290 (8)	0.0249 (7)	0.0008 (6)	-0.0056 (6)	-0.0033 (6)

Geometric parameters (Å, °)

F1—C1	1.3415 (19)	C5—C10	1.397 (2)
F2—C1	1.338 (2)	C6—C7	1.380 (2)
F3—C1	1.3429 (19)	C6—H6	0.9300
O1—C4	1.2618 (18)	C7—C8	1.390 (3)
O2—C2	1.2732 (18)	C7—H7	0.9300
N1—C11	1.326 (2)	C8—C9	1.380 (2)
N1—H1A	0.877 (9)	C8—H8	0.9300
N1—H1B	0.879 (9)	C9—C10	1.392 (2)
N2—C11	1.346 (2)	C9—H9	0.9300
N2—C12	1.401 (2)	C10—H10	0.9300
N2—H2A	0.905 (10)	C12—C13	1.385 (2)
N3—C11	1.344 (2)	C12—C17	1.396 (2)
N3—C17	1.400 (2)	C13—C14	1.390 (2)
N3—H3A	0.895 (9)	C13—H13	0.9300
C1—C2	1.530 (2)	C14—C15	1.395 (2)
C2—C3	1.385 (2)	C14—H14	0.9300
C3—C4	1.430 (2)	C15—C16	1.388 (2)
C3—H3	0.9300	C15—H15	0.9300
C4—C5	1.499 (2)	C16—C17	1.385 (2)
C5—C6	1.394 (2)	C16—H16	0.9300
C11—N1—H1A	120.8 (12)	C8—C7—H7	120.0
C11—N1—H1B	118.1 (11)	C9—C8—C7	119.73 (16)
H1A—N1—H1B	121.1 (14)	C9—C8—H8	120.1
C11—N2—C12	108.60 (13)	C7—C8—H8	120.1
C11—N2—H2A	125.0 (13)	C8—C9—C10	120.49 (15)
C12—N2—H2A	125.5 (13)	C8—C9—H9	119.8
C11—N3—C17	108.79 (13)	C10—C9—H9	119.8
C11—N3—H3A	126.8 (14)	C9—C10—C5	120.07 (15)
C17—N3—H3A	124.4 (14)	C9—C10—H10	120.0
F2—C1—F1	106.81 (13)	C5—C10—H10	120.0
F2—C1—F3	105.99 (13)	N1—C11—N3	125.36 (14)
F1—C1—F3	106.30 (13)	N1—C11—N2	125.23 (15)
F2—C1—C2	111.64 (13)	N3—C11—N2	109.40 (14)
F1—C1—C2	114.24 (14)	C13—C12—C17	121.81 (14)

F3—C1—C2	111.35 (13)	C13—C12—N2	131.50 (15)
O2—C2—C3	128.36 (14)	C17—C12—N2	106.66 (13)
O2—C2—C1	113.13 (14)	C12—C13—C14	116.39 (15)
C3—C2—C1	118.47 (14)	C12—C13—H13	121.8
C2—C3—C4	123.55 (14)	C14—C13—H13	121.8
C2—C3—H3	118.2	C13—C14—C15	121.95 (15)
C4—C3—H3	118.2	C13—C14—H14	119.0
O1—C4—C3	123.29 (14)	C15—C14—H14	119.0
O1—C4—C5	117.51 (13)	C16—C15—C14	121.39 (15)
C3—C4—C5	119.18 (13)	C16—C15—H15	119.3
C6—C5—C10	118.71 (14)	C14—C15—H15	119.3
C6—C5—C4	118.92 (13)	C17—C16—C15	116.78 (15)
C10—C5—C4	122.34 (14)	C17—C16—H16	121.6
C7—C6—C5	120.98 (15)	C15—C16—H16	121.6
C7—C6—H6	119.5	C16—C17—C12	121.68 (14)
C5—C6—H6	119.5	C16—C17—N3	131.78 (14)
C6—C7—C8	120.01 (16)	C12—C17—N3	106.52 (13)
C6—C7—H7	120.0		
F2—C1—C2—O2	48.38 (18)	C4—C5—C10—C9	-177.14 (14)
F1—C1—C2—O2	169.71 (14)	C17—N3—C11—N1	179.40 (15)
F3—C1—C2—O2	-69.86 (17)	C17—N3—C11—N2	-1.38 (17)
F2—C1—C2—C3	-133.69 (15)	C12—N2—C11—N1	-179.13 (15)
F1—C1—C2—C3	-12.4 (2)	C12—N2—C11—N3	1.64 (18)
F3—C1—C2—C3	108.07 (16)	C11—N2—C12—C13	177.01 (16)
O2—C2—C3—C4	-7.8 (3)	C11—N2—C12—C17	-1.26 (17)
C1—C2—C3—C4	174.64 (14)	C17—C12—C13—C14	0.3 (2)
C2—C3—C4—O1	-8.1 (2)	N2—C12—C13—C14	-177.71 (16)
C2—C3—C4—C5	173.70 (14)	C12—C13—C14—C15	-0.6 (2)
O1—C4—C5—C6	-27.7 (2)	C13—C14—C15—C16	0.3 (2)
C3—C4—C5—C6	150.58 (15)	C14—C15—C16—C17	0.3 (2)
O1—C4—C5—C10	150.25 (14)	C15—C16—C17—C12	-0.5 (2)
C3—C4—C5—C10	-31.4 (2)	C15—C16—C17—N3	177.26 (15)
C10—C5—C6—C7	0.1 (2)	C13—C12—C17—C16	0.2 (2)
C4—C5—C6—C7	178.13 (15)	N2—C12—C17—C16	178.68 (14)
C5—C6—C7—C8	-1.0 (3)	C13—C12—C17—N3	-178.06 (14)
C6—C7—C8—C9	1.1 (3)	N2—C12—C17—N3	0.42 (16)
C7—C8—C9—C10	-0.1 (3)	C11—N3—C17—C16	-177.45 (16)
C8—C9—C10—C5	-0.8 (2)	C11—N3—C17—C12	0.57 (16)
C6—C5—C10—C9	0.9 (2)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1A \cdots O2 ⁱ	0.88 (1)	2.00 (1)	2.7803 (18)	147 (2)
N3—H3A \cdots O1 ⁱ	0.90 (1)	1.95 (1)	2.7651 (17)	151 (2)
N3—H3A \cdots O2 ⁱ	0.90 (1)	2.36 (2)	2.9912 (18)	127 (2)

supporting information

N1—H1B···O1	0.88 (1)	2.03 (1)	2.8662 (18)	158 (2)
N2—H2A···O2	0.91 (1)	1.86 (1)	2.7360 (18)	162 (2)

Symmetry code: (i) $-x+2, y-1/2, -z+1/2$.