

# A crystallographically isolated dimeric hydrolyzed chlorophosphazene dianion

Matthew J. Panzner, Wiley J. Youngs and Claire A. Tessier\*

 The University of Akron, Chemistry Department, Akron, OH 44325-3601, USA  
 Correspondence e-mail: tessier@uakron.edu


Received 17 November 2008; accepted 10 December 2008

 Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.009$  Å;  $R$  factor = 0.047;  $wR$  factor = 0.123; data-to-parameter ratio = 17.4.

Single crystals of the title compound bis[bis(1-ethyl-3-methylimidazol-2-ylidene)silver(I)] 1,5,5,7,11,11-hexachloro-2,8-dioxa-4,6,10,12,13,14-hexaaza-1 $\lambda^5$ ,3,5 $\lambda^5$ ,7 $\lambda^5$ ,9,11 $\lambda^5$ -hexaphosphatricyclo[7.3.1.1<sup>3,7</sup>]tetradeca-1(13),4,7(14),10-tetraene-6,12-diide 3,9-dioxide,  $[\text{Ag}(\text{C}_6\text{H}_{10}\text{N}_2)_2](\text{Cl}_6\text{N}_6\text{O}_4\text{P}_6)_{0.5}$ , were isolated from the reaction of the silver *N*-heteocyclic carbene complex  $[\text{Ag}(\text{C}_6\text{H}_{10}\text{N}_2)_2]\text{Cl}$  and hexachlorocyclotriphosphazene  $[\text{NPCl}_2]_3$  in the presence of water. The asymmetric unit contains one silver carbene cation with the carbene ligands bound to the Ag(I) in an almost linear arrangement and one half of a hydrolyzed phosphazene dianion. The second cation and additional half of the anion are generated by an inversion center.

## Related literature

For background on phosphazene hydrolysis products, see: Allcock (2003); Allcock *et al.* (1975); Gabler & Haw (1990); Murray *et al.* (1994); van de Grampel (1992). For related structures, see: Bartlett *et al.* (2006); Brandt *et al.* (1991); Bullen (1971); Meetsma *et al.* (1990).



## Experimental

### Crystal data

$[\text{Ag}(\text{C}_6\text{H}_{10}\text{N}_2)_2](\text{Cl}_6\text{N}_6\text{O}_4\text{P}_6)_{0.5}$	$\alpha = 78.916$ (3)°
$M_r = 601.48$	$\beta = 71.558$ (3)°
Triclinic, $P\bar{1}$	$\gamma = 76.107$ (3)°
$a = 9.3224$ (15) Å	$V = 1108.4$ (3) Å <sup>3</sup>
$b = 10.6190$ (18) Å	$Z = 2$
$c = 12.257$ (2) Å	Mo $K\alpha$ radiation

 $\mu = 1.51$  mm<sup>-1</sup>  
 $T = 100$  (2) K

 $0.30 \times 0.08 \times 0.04$  mm

### Data collection

Bruker SMART CCD area-detector diffractometer	8912 measured reflections
Absorption correction: multi-scan (SADABS; Bruker, 2001)	4459 independent reflections
$T_{\min} = 0.660$ , $T_{\max} = 0.942$	3484 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.031$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.047$	257 parameters
$wR(F^2) = 0.123$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\text{max}} = 2.25$ e Å <sup>-3</sup>
4459 reflections	$\Delta\rho_{\text{min}} = -1.02$ e Å <sup>-3</sup>

**Table 1**

Selected geometric parameters (Å, °).

Ag—C1	2.065 (5)	P1—N6	1.615 (4)
Ag—C7	2.070 (5)	P1—O2	1.657 (3)
Cl1—P2	2.0118 (16)	P2—N6	1.555 (4)
Cl2—P2	2.0171 (17)	P2—N7	1.578 (4)
Cl3—P3	2.0217 (16)	P3—N5 <sup>i</sup>	1.554 (4)
P1—O1	1.468 (3)	P3—O2	1.578 (3)
P1—N5	1.610 (4)	P3—N7 <sup>i</sup>	1.594 (4)
Cl1—Ag—C7	178.72 (19)		

 Symmetry code: (i)  $-x + 1, -y + 1, -z + 1$ .

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

We thank the National Science Foundation (USA) for support of this work under grants CHE-0316944 and CHE-0616601 and grant CHE-0116041 for the funds used to support the X-ray facilities at the University of Akron.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BR2087).

## References

- Allcock, H. R. (2003). *Chemistry and Applications of Polyphosphazenes*, pp. 137–187. New Jersey: Wiley-Interscience.
- Allcock, H. R., Gardener, J. E. & Smeltz, K. M. (1975). *Macromolecules*, **8**, 36–42.
- Bartlett, S. W., Coles, S. J., Davies, D. B., Hursthouse, M. B., İbişoğlu, H., Kiliç, A., Shaw, R. A. & Ün, İ. (2006). *Acta Cryst.* **B62**, 321–329.
- Brandt, K., van de Grampel, J. C., Meetsma, A. & Jekel, A. P. (1991). *Recl Trav. Chim. Pays Bas*, **110**, 27–28.
- Bruker (2001). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bullen, G. J. (1971). *J. Chem. Soc. A*, 1450–1453.
- Gabler, D. G. & Haw, J. F. (1990). *Inorg. Chem.* **29**, 4018–4021.
- Meetsma, A., van der Lee, A., Jekel, A. P., van de Grampel, J. C. & Brandt, K. (1990). *Acta Cryst.* **C46**, 909–911.
- Murray, M., Penkett, C. J., Rillie, I. M. & Singh, G. (1994). *Phosphorus, Sulfur, Silicon*. **93–94**, 313–316.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Grampel, J. C. van de (1992). *Coord. Chem. Rev.* **112**, 247–271.

## supporting information

*Acta Cryst.* (2009). E65, m105 [doi:10.1107/S1600536808042116]

## A crystallographically isolated dimeric hydrolyzed chlorophosphazene dianion

Matthew J. Panzner, Wiley J. Youngs and Claire A. Tessier

### S1. Comment

Hexachlorocyclotriphosphazene  $[\text{NPCl}_2]_3$  is used as a starting material for the synthesis of poly(dichlorophosphazene). This conversion can be achieved by a ring opening melt polymerization process at high temperature, generally greater than 473 °K. It has long been contested that trace amounts of water not only accelerate the rate of polymerization but are necessary to generate active species needed to simply promote polymer formation (Allcock, 2003; Allcock *et al.*, 1975). Isolation of hydrolyzed species gives insight into the still unclear role that water plays in this polymerization reaction (Gabler *et al.*, 1990). As part of our broader studies on the irreproducibility in the synthesis of poly(dichlorophosphazene) we report the first crystal structure of a dimeric hydrolyzed chlorophosphazene dianion.


The asymmetric unit consists of one silver N-heterocyclic carbene cation  $[\text{AgC}_{12}\text{H}_{20}\text{N}_4]^+$  and half of a hydrolyzed phosphazene dianion  $[\text{Cl}_6\text{N}_6\text{O}_4\text{P}_6]^{2-}$ . The second cation and other half of the dianion are generated by a crystallographic inversion center at  $x,y,z$  (1/2,1/2,1/2) located between one of the bridging oxygen atoms of the dimeric dianion and its symmetry generated equivalent. Each of the Ag(I) atoms is bound to two identical N-heterocyclic carbene ligands in a linear fashion. Two partially hydrolyzed phosphazene rings are joined by bridging oxygen atoms from a phosphorus atom of one ring to the other to form the dimeric dianion. The P—O—P bonds of the bridging oxygen atoms are inequivalent. The P—N bond distances of the individual rings deviate from the reported values of hexachlorocyclotriphosphazene, all six being virtually equivalent (Bartlett *et al.*, 2006; Bullen, 1971). Two of the P—N bonds on each of the rings show significant double bond character while the remaining P—N bonds are lengthened to give more single bond character.

### S2. Experimental

The title compound, bis[bis(1-ethyl-3-methylimidazol-2-ylidene)silver(I)] 1,5,5,7,11,11-hexachloro-2,8-dioxa-4,6,10,12,13,14-hexaaza-1 $\lambda^5$ ,3,5 $\lambda^5$ ,7 $\lambda^5$ ,9,11 $\lambda^5$ -hexaphosphatricyclo[7.3.1.1<sup>3,7</sup>]tetradeca-1(13),4,7(14),10-tetraene-6,12-diide 3,9-dioxide,  $[\text{Ag}(\text{C}_6\text{H}_{10}\text{N}_2)_2](\text{Cl}_6\text{N}_6\text{O}_4\text{P}_6)_{0.5}$ , was isolated from the reaction of silver N-heterocyclic carbene complex  $[\text{AgC}_{12}\text{H}_{20}\text{N}_4]^+\text{Cl}^-$  (generated *in situ*) and hexachlorocyclotriphosphazene,  $[\text{NPCl}_2]_3$ . A small amount of crystals were obtained after removing volatiles from the reaction.

### S3. Refinement

Hydrogen atoms were calculated and placed in geometrically idealized positions with C—H distances of 0.95 Å (aromatic), 0.99 Å (methylene), and 0.98 Å (methyl). H atoms were constrained to ride on the parent carbon atom with  $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$  for aromatic and methylene and  $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{C})$  for methyl protons.

**Figure 1**

Structure of the title compound, first crystallographically characterized dimeric oxygen-bridged chlorophosphazene dianion. A numbering scheme of the non-H atoms is shown and thermal ellipsoids shown at 50% probability.

**bis[bis(1-ethyl-3-methylimidazol-2-ylidene)silver(I)] 1,5,5,7,11,11-hexachloro-2,8-dioxa-4,6,10,12,13,14-hexaaza-1 $\lambda$ <sup>5</sup>,3,5 $\lambda$ <sup>5</sup>,7 $\lambda$ <sup>5</sup>,9,11 $\lambda$ <sup>5</sup>-hexaphosphatrimethylene-1(13),4,7(14),10-tetraene-6,12-dioxide 3,9-dioxide**

*Crystal data*

$[\text{Ag}(\text{C}_6\text{H}_{10}\text{N}_2)_2](\text{Cl}_6\text{N}_6\text{O}_4\text{P}_6)_{0.5}$

$M_r = 601.48$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 9.3224(15)\ \text{\AA}$

$b = 10.6190(18)\ \text{\AA}$

$c = 12.257(2)\ \text{\AA}$

$\alpha = 78.916(3)^\circ$

$\beta = 71.558(3)^\circ$

$\gamma = 76.107(3)^\circ$

$V = 1108.4(3)\ \text{\AA}^3$

$Z = 2$

$F(000) = 600$

$D_x = 1.802\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2618 reflections

$\theta = 2.5\text{--}26.8^\circ$

$\mu = 1.51\ \text{mm}^{-1}$

$T = 100\ \text{K}$

Plate, colourless

$0.30 \times 0.08 \times 0.04\ \text{mm}$

*Data collection*

Bruker SMART CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2001)

$T_{\text{min}} = 0.660$ ,  $T_{\text{max}} = 0.942$

8912 measured reflections

4459 independent reflections

3484 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\text{max}} = 26.3^\circ$ ,  $\theta_{\text{min}} = 1.8^\circ$

$h = -11 \rightarrow 11$

$k = -13 \rightarrow 13$

$l = -15 \rightarrow 15$

Refinement

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.047$   
 $wR(F^2) = 0.123$   
 $S = 1.04$   
 4459 reflections  
 257 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0737P)^2]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 2.25 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -1.02 \text{ e } \text{\AA}^{-3}$

Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Ag	0.92396 (4)	0.82923 (3)	0.02673 (3)	0.02921 (14)
Cl1	0.20816 (15)	0.96514 (11)	0.50044 (11)	0.0364 (3)
Cl2	0.06691 (14)	0.72027 (14)	0.58126 (11)	0.0402 (3)
Cl3	0.32100 (13)	0.28263 (12)	0.39490 (11)	0.0312 (3)
P1	0.50886 (13)	0.63373 (11)	0.34302 (10)	0.0220 (3)
P2	0.27569 (13)	0.76965 (11)	0.50900 (10)	0.0218 (3)
P3	0.46494 (13)	0.36960 (11)	0.43519 (10)	0.0196 (3)
O1	0.5964 (4)	0.6691 (3)	0.2226 (3)	0.0298 (8)
O2	0.4577 (3)	0.4948 (3)	0.3425 (3)	0.0243 (7)
N1	0.6862 (5)	1.0763 (4)	0.0964 (4)	0.0346 (10)
N2	0.6984 (5)	1.0504 (4)	-0.0734 (4)	0.0330 (10)
N3	1.1582 (4)	0.6223 (4)	0.1266 (3)	0.0298 (9)
N4	1.1637 (4)	0.5782 (4)	-0.0374 (3)	0.0277 (9)
N5	0.6039 (4)	0.5991 (3)	0.4376 (3)	0.0219 (8)
N6	0.3458 (4)	0.7321 (4)	0.3838 (3)	0.0259 (9)
N7	0.3694 (4)	0.7221 (3)	0.6012 (3)	0.0207 (8)
C1	0.7588 (5)	0.9961 (4)	0.0151 (4)	0.0287 (11)
C2	0.5818 (6)	1.1759 (5)	0.0603 (6)	0.0454 (15)
H2	0.5168	1.2441	0.1033	0.055*
C3	0.5876 (6)	1.1600 (5)	-0.0462 (5)	0.0414 (14)
H3	0.5274	1.2135	-0.0937	0.050*
C4	0.7138 (7)	1.0544 (6)	0.2095 (5)	0.0455 (14)
H4A	0.8242	1.0172	0.2013	0.055*
H4B	0.6891	1.1391	0.2404	0.055*
C5	0.6220 (9)	0.9672 (7)	0.2893 (6)	0.0638 (19)

H5A	0.5127	1.0074	0.3028	0.096*
H5B	0.6483	0.9494	0.3629	0.096*
H5C	0.6423	0.8850	0.2569	0.096*
C6	0.7431 (8)	0.9969 (6)	-0.1845 (5)	0.0513 (16)
H6A	0.8545	0.9887	-0.2195	0.077*
H6B	0.6889	1.0558	-0.2370	0.077*
H6C	0.7155	0.9106	-0.1707	0.077*
C7	1.0921 (5)	0.6642 (5)	0.0399 (4)	0.0263 (10)
C8	1.2702 (6)	0.5110 (5)	0.1032 (5)	0.0398 (13)
H8	1.3328	0.4632	0.1511	0.048*
C9	1.2735 (6)	0.4839 (5)	0.0001 (4)	0.0375 (13)
H9	1.3389	0.4132	-0.0394	0.045*
C10	1.1113 (6)	0.6846 (5)	0.2325 (4)	0.0369 (12)
H10A	1.2010	0.6708	0.2631	0.044*
H10B	1.0781	0.7800	0.2135	0.044*
C11	0.9870 (8)	0.6337 (8)	0.3212 (6)	0.074 (2)
H11A	0.9042	0.6342	0.2878	0.111*
H11B	0.9471	0.6884	0.3843	0.111*
H11C	1.0255	0.5439	0.3516	0.111*
C12	1.1316 (6)	0.5854 (5)	-0.1481 (4)	0.0313 (11)
H12A	1.0230	0.6247	-0.1405	0.047*
H12B	1.1532	0.4971	-0.1697	0.047*
H12C	1.1970	0.6390	-0.2081	0.047*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Ag	0.0342 (2)	0.0232 (2)	0.0293 (2)	0.00745 (15)	-0.01754 (17)	-0.00302 (15)
Cl1	0.0483 (7)	0.0219 (6)	0.0319 (7)	0.0102 (5)	-0.0150 (6)	-0.0023 (5)
Cl2	0.0273 (6)	0.0564 (9)	0.0366 (7)	-0.0102 (6)	-0.0147 (5)	0.0073 (6)
Cl3	0.0295 (6)	0.0343 (7)	0.0365 (7)	-0.0071 (5)	-0.0155 (5)	-0.0084 (5)
P1	0.0282 (6)	0.0178 (6)	0.0176 (6)	0.0025 (5)	-0.0096 (5)	0.0003 (4)
P2	0.0234 (6)	0.0208 (6)	0.0185 (6)	0.0026 (5)	-0.0095 (5)	0.0015 (5)
P3	0.0223 (6)	0.0175 (6)	0.0200 (6)	-0.0007 (4)	-0.0114 (5)	0.0005 (4)
O1	0.0401 (19)	0.0250 (17)	0.0163 (17)	0.0015 (14)	-0.0058 (14)	0.0028 (13)
O2	0.0313 (17)	0.0224 (16)	0.0219 (17)	0.0008 (13)	-0.0180 (14)	0.0010 (13)
N1	0.037 (2)	0.025 (2)	0.039 (3)	0.0035 (18)	-0.015 (2)	-0.0027 (19)
N2	0.036 (2)	0.025 (2)	0.035 (2)	-0.0018 (18)	-0.0168 (19)	0.0090 (18)
N3	0.030 (2)	0.033 (2)	0.027 (2)	0.0016 (17)	-0.0141 (18)	-0.0029 (18)
N4	0.029 (2)	0.027 (2)	0.022 (2)	0.0010 (17)	-0.0080 (17)	0.0022 (16)
N5	0.0208 (18)	0.022 (2)	0.022 (2)	0.0016 (15)	-0.0099 (15)	-0.0025 (15)
N6	0.032 (2)	0.024 (2)	0.0180 (19)	0.0097 (16)	-0.0143 (16)	0.0007 (15)
N7	0.0247 (19)	0.0208 (19)	0.0169 (19)	-0.0001 (15)	-0.0103 (15)	-0.0007 (15)
C1	0.034 (3)	0.023 (2)	0.033 (3)	-0.003 (2)	-0.019 (2)	0.000 (2)
C2	0.039 (3)	0.027 (3)	0.058 (4)	0.005 (2)	-0.010 (3)	0.002 (3)
C3	0.033 (3)	0.032 (3)	0.048 (4)	0.000 (2)	-0.015 (3)	0.018 (3)
C4	0.053 (4)	0.041 (3)	0.045 (3)	-0.003 (3)	-0.015 (3)	-0.018 (3)
C5	0.088 (5)	0.062 (4)	0.055 (4)	-0.026 (4)	-0.032 (4)	-0.004 (3)

C6	0.076 (4)	0.048 (4)	0.033 (3)	-0.007 (3)	-0.030 (3)	0.006 (3)
C7	0.025 (2)	0.027 (3)	0.025 (2)	-0.0039 (19)	-0.010 (2)	0.0021 (19)
C8	0.034 (3)	0.040 (3)	0.044 (3)	0.010 (2)	-0.024 (3)	-0.001 (3)
C9	0.035 (3)	0.036 (3)	0.031 (3)	0.013 (2)	-0.011 (2)	-0.003 (2)
C10	0.037 (3)	0.045 (3)	0.031 (3)	-0.002 (2)	-0.017 (2)	-0.008 (2)
C11	0.086 (5)	0.110 (6)	0.034 (4)	-0.055 (5)	0.013 (3)	-0.029 (4)
C12	0.032 (3)	0.040 (3)	0.016 (2)	-0.004 (2)	-0.003 (2)	0.000 (2)

*Geometric parameters (Å, °)*

Ag—C1	2.065 (5)	N5—P3 <sup>i</sup>	1.554 (4)
Ag—C7	2.070 (5)	N7—P3 <sup>i</sup>	1.594 (4)
C11—P2	2.0118 (16)	C2—C3	1.331 (8)
C12—P2	2.0171 (17)	C2—H2	0.9500
C13—P3	2.0217 (16)	C3—H3	0.9500
P1—O1	1.468 (3)	C4—C5	1.432 (9)
P1—N5	1.610 (4)	C4—H4A	0.9900
P1—N6	1.615 (4)	C4—H4B	0.9900
P1—O2	1.657 (3)	C5—H5A	0.9800
P2—N6	1.555 (4)	C5—H5B	0.9800
P2—N7	1.578 (4)	C5—H5C	0.9800
P3—N5 <sup>i</sup>	1.554 (4)	C6—H6A	0.9800
P3—O2	1.578 (3)	C6—H6B	0.9800
P3—N7 <sup>i</sup>	1.594 (4)	C6—H6C	0.9800
N1—C1	1.341 (6)	C8—C9	1.338 (8)
N1—C2	1.365 (6)	C8—H8	0.9500
N1—C4	1.454 (7)	C9—H9	0.9500
N2—C1	1.348 (6)	C10—C11	1.448 (8)
N2—C3	1.374 (6)	C10—H10A	0.9900
N2—C6	1.475 (7)	C10—H10B	0.9900
N3—C7	1.345 (6)	C11—H11A	0.9800
N3—C8	1.385 (6)	C11—H11B	0.9800
N3—C10	1.465 (6)	C11—H11C	0.9800
N4—C7	1.347 (6)	C12—H12A	0.9800
N4—C9	1.374 (6)	C12—H12B	0.9800
N4—C12	1.464 (6)	C12—H12C	0.9800
C1—Ag—C7	178.72 (19)	C5—C4—H4A	109.4
O1—P1—N5	116.2 (2)	N1—C4—H4A	109.4
O1—P1—N6	113.33 (19)	C5—C4—H4B	109.4
N5—P1—N6	112.96 (19)	N1—C4—H4B	109.4
O1—P1—O2	104.90 (18)	H4A—C4—H4B	108.0
N5—P1—O2	104.43 (17)	C4—C5—H5A	109.5
N6—P1—O2	103.29 (19)	C4—C5—H5B	109.5
N6—P2—N7	120.75 (19)	H5A—C5—H5B	109.5
N6—P2—C11	108.88 (15)	C4—C5—H5C	109.5
N7—P2—C11	108.31 (15)	H5A—C5—H5C	109.5
N6—P2—C12	110.61 (17)	H5B—C5—H5C	109.5

N7—P2—C12	107.54 (15)	N2—C6—H6A	109.5
C11—P2—C12	98.43 (7)	N2—C6—H6B	109.5
N5 <sup>i</sup> —P3—O2	113.38 (18)	H6A—C6—H6B	109.5
N5 <sup>i</sup> —P3—N7 <sup>i</sup>	119.10 (19)	N2—C6—H6C	109.5
O2—P3—N7 <sup>i</sup>	109.28 (18)	H6A—C6—H6C	109.5
N5 <sup>i</sup> —P3—C13	109.53 (15)	H6B—C6—H6C	109.5
O2—P3—C13	97.59 (12)	N3—C7—N4	104.7 (4)
N7 <sup>i</sup> —P3—C13	105.49 (14)	N3—C7—Ag	127.0 (3)
P3—O2—P1	126.57 (19)	N4—C7—Ag	128.2 (3)
C1—N1—C2	111.3 (5)	C9—C8—N3	106.8 (4)
C1—N1—C4	123.2 (4)	C9—C8—H8	126.6
C2—N1—C4	125.5 (5)	N3—C8—H8	126.6
C1—N2—C3	111.5 (4)	C8—C9—N4	106.6 (4)
C1—N2—C6	124.2 (4)	C8—C9—H9	126.7
C3—N2—C6	124.3 (4)	N4—C9—H9	126.7
C7—N3—C8	110.6 (4)	C11—C10—N3	112.4 (5)
C7—N3—C10	123.6 (4)	C11—C10—H10A	109.1
C8—N3—C10	125.7 (4)	N3—C10—H10A	109.1
C7—N4—C9	111.2 (4)	C11—C10—H10B	109.1
C7—N4—C12	124.6 (4)	N3—C10—H10B	109.1
C9—N4—C12	124.2 (4)	H10A—C10—H10B	107.9
P3 <sup>i</sup> —N5—P1	124.3 (2)	C10—C11—H11A	109.5
P2—N6—P1	123.1 (2)	C10—C11—H11B	109.5
P2—N7—P3 <sup>i</sup>	118.6 (2)	H11A—C11—H11B	109.5
N1—C1—N2	103.8 (4)	C10—C11—H11C	109.5
N1—C1—Ag	127.0 (4)	H11A—C11—H11C	109.5
N2—C1—Ag	129.2 (4)	H11B—C11—H11C	109.5
C3—C2—N1	107.4 (5)	N4—C12—H12A	109.5
C3—C2—H2	126.3	N4—C12—H12B	109.5
N1—C2—H2	126.3	H12A—C12—H12B	109.5
C2—C3—N2	105.9 (4)	N4—C12—H12C	109.5
C2—C3—H3	127.0	H12A—C12—H12C	109.5
N2—C3—H3	127.0	H12B—C12—H12C	109.5
C5—C4—N1	111.3 (5)		
N5 <sup>i</sup> —P3—O2—P1	47.2 (3)	C7—Ag—C1—N1	-47 (9)
N7 <sup>i</sup> —P3—O2—P1	-88.3 (3)	C7—Ag—C1—N2	133 (8)
C13—P3—O2—P1	162.3 (2)	C1—N1—C2—C3	-0.3 (7)
O1—P1—O2—P3	136.3 (2)	C4—N1—C2—C3	-178.1 (5)
N5—P1—O2—P3	13.5 (3)	N1—C2—C3—N2	-0.7 (6)
N6—P1—O2—P3	-104.8 (3)	C1—N2—C3—C2	1.5 (6)
O1—P1—N5—P3 <sup>i</sup>	146.1 (3)	C6—N2—C3—C2	-179.9 (5)
N6—P1—N5—P3 <sup>i</sup>	12.6 (4)	C1—N1—C4—C5	-84.8 (7)
O2—P1—N5—P3 <sup>i</sup>	-98.9 (3)	C2—N1—C4—C5	92.9 (7)
N7—P2—N6—P1	5.5 (4)	C8—N3—C7—N4	0.0 (6)
C11—P2—N6—P1	131.7 (2)	C10—N3—C7—N4	-177.7 (4)
C12—P2—N6—P1	-121.2 (3)	C8—N3—C7—Ag	-177.6 (4)
O1—P1—N6—P2	-144.9 (3)	C10—N3—C7—Ag	4.7 (7)

---

N5—P1—N6—P2	-10.1 (4)	C9—N4—C7—N3	-0.2 (6)
O2—P1—N6—P2	102.2 (3)	C12—N4—C7—N3	-179.0 (4)
N6—P2—N7—P3 <sup>i</sup>	-2.2 (4)	C9—N4—C7—Ag	177.5 (4)
C11—P2—N7—P3 <sup>i</sup>	-128.6 (2)	C12—N4—C7—Ag	-1.4 (7)
C12—P2—N7—P3 <sup>i</sup>	125.9 (2)	C1—Ag—C7—N3	40 (9)
C2—N1—C1—N2	1.2 (6)	C1—Ag—C7—N4	-137 (8)
C4—N1—C1—N2	179.1 (5)	C7—N3—C8—C9	0.1 (6)
C2—N1—C1—Ag	-178.8 (4)	C10—N3—C8—C9	177.7 (5)
C4—N1—C1—Ag	-0.9 (7)	N3—C8—C9—N4	-0.2 (6)
C3—N2—C1—N1	-1.6 (6)	C7—N4—C9—C8	0.2 (6)
C6—N2—C1—N1	179.8 (5)	C12—N4—C9—C8	179.0 (5)
C3—N2—C1—Ag	178.3 (4)	C7—N3—C10—C11	87.5 (7)
C6—N2—C1—Ag	-0.2 (8)	C8—N3—C10—C11	-89.9 (7)

---

Symmetry code: (i)  $-x+1, -y+1, -z+1$ .