

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

***trans*-1,2,3-Tris(4-methoxybenzoyl)-cyclopropane**

Jingjing Sun* and Nengfang She

Key Laboratory of Pesticides and Chemical Biology of the Ministry of Education, College of Chemistry, Central China Normal University, Wuhan 430079, People's Republic of China

Correspondence e-mail: ajsun@mails.ccnu.edu.cn


Received 10 December 2008; accepted 12 February 2009

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.057; wR factor = 0.155; data-to-parameter ratio = 14.4.

In the title compound, $\text{C}_{27}\text{H}_{24}\text{O}_6$, the packing of the molecules is mainly governed by $\text{C}-\text{H}\cdots\text{O}$ interactions.

Related literature

For related structures, see: Saba (1990). For background to the chemistry of cyclopropanes as a versatile tool in organic synthesis, see: Wong (1989).



Experimental

Crystal data

$\text{C}_{27}\text{H}_{24}\text{O}_6$
 $M_r = 444.46$
 Triclinic, $P\bar{1}$

$a = 10.1897$ (6) Å
 $b = 10.626$ (6) Å
 $c = 10.6931$ (6) Å

$\alpha = 90.736$ (1)°
 $\beta = 103.194$ (1)°
 $\gamma = 92.432$ (1)°
 $V = 1125.9$ (6) Å³
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 0.09$ mm⁻¹
 $T = 298$ K
 $0.30 \times 0.20 \times 0.10$ mm

Data collection

Bruker SMART CCD area-detector diffractometer
 Absorption correction: none
 9852 measured reflections

4349 independent reflections
 2862 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.155$
 $S = 1.06$
 4349 reflections

301 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.19$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.17$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C26}-\text{H26}\cdots\text{O2}^{\text{i}}$	0.93	2.54	3.467 (3)	175
$\text{C11}-\text{H11}\cdots\text{O6}^{\text{ii}}$	0.98	2.51	3.270 (3)	134
$\text{C7}-\text{H7}\cdots\text{O5}^{\text{iii}}$	0.93	2.52	3.171 (3)	128
$\text{C9}-\text{H9}\cdots\text{O3}^{\text{iii}}$	0.98	2.55	3.497 (3)	164
$\text{C27}-\text{H27B}\cdots\text{O4}^{\text{iv}}$	0.96	2.58	3.136 (3)	117

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $x+1, y, z$; (iii) $-x+1, -y+2, -z$; (iv) $-x, -y+1, -z+1$.

Data collection: *SMART* (Bruker, 2001); cell refinement: *SAINT* (Bruker, 2001); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Professor An-Xin Wu for technical assistance and Dr Xiang-Gao Meng for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2123).

References

- Bruker (2001). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Saba, A. (1990). *J. Chem. Res. Synop.* **9**, 288–289.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Wong, H. N. C. (1989). *Chem. Rev.* **89**, 165–198.

supporting information

Acta Cryst. (2009). E65, o588 [doi:10.1107/S1600536809005145]

trans*-1,2,3-Tris(4-methoxybenzoyl)cyclopropane*Jingjing Sun and Nengfang She****S1. Comment**

The unusual bonding of cyclopropanes and the strain release associated with cleavage of cyclopropanes offer the possibility of recognizing that structural unit when it is a part of a larger molecule. We report here the molecular structure of the title cyclopropane derivative (Fig. 1), which is an important intermediate in organic synthesis (Saba, 1990). Since numerous methodologies have been developed for the construction of three-membered carbocycles, the chemistry of cyclopropanes has emerged as a versatile tool in organic synthesis (Wong, 1989).


The crystal packing is stabilized by intermolecular C—H···O interaction.

S2. Experimental

The title compound was synthesized according to the procedure reported (Saba, 1990). Crystals appropriate for X-ray data collection were obtained by slow evaporation of the dichloromethane solution at 293 K.

S3. Refinement

All H atoms were positioned in geometrically idealized positions and constrained to ride on their parent atoms, with C—H distances in the range 0.93–0.98 Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$.

**Figure 1**

The molecular structure of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are presented as a small spheres of arbitrary radius.

trans-1,2,3-Tris(4-methoxybenzoyl)cyclopropane

Crystal data

$C_{27}H_{24}O_6$

$M_r = 444.46$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 10.1897(6)\ \text{\AA}$

$b = 10.626(6)\ \text{\AA}$

$c = 10.6931(6)\ \text{\AA}$

$\alpha = 90.736(1)^\circ$

$\beta = 103.194(1)^\circ$

$\gamma = 92.432(1)^\circ$

$V = 1125.9(6)\ \text{\AA}^3$

$Z = 2$

$F(000) = 468$

$D_x = 1.311\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 1774 reflections

$\theta = 2.7\text{--}21.3^\circ$

$\mu = 0.09\ \text{mm}^{-1}$

$T = 298$ K
Block, colourless

$0.30 \times 0.20 \times 0.10$ mm

Data collection

Bruker SMART CCD area-detector
diffractometer
Radiation source: Fine-focus sealed tube
Graphite monochromator
 φ and ω scans
9852 measured reflections
4349 independent reflections

2862 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.031$
 $\theta_{\text{max}} = 26.0^\circ$, $\theta_{\text{min}} = 1.9^\circ$
 $h = -12 \rightarrow 12$
 $k = -10 \rightarrow 13$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
Least-squares matrix: Full
 $R[F^2 > 2\sigma(F^2)] = 0.057$
 $wR(F^2) = 0.155$
 $S = 1.06$
4349 reflections
301 parameters
0 restraints
Primary atom site location: Direct

Secondary atom site location: Difmap
Hydrogen site location: Geom
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0697P)^2 + 0.0779P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.19 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.17 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.0074 (3)	1.5583 (3)	0.2073 (3)	0.0938 (10)
H1A	0.9340	1.6141	0.1903	0.141*
H1B	1.0773	1.5892	0.1672	0.141*
H1C	1.0425	1.5543	0.2983	0.141*
C2	0.8628 (2)	1.3733 (2)	0.2028 (2)	0.0559 (6)
C3	0.8010 (3)	1.4209 (2)	0.2944 (2)	0.0632 (7)
H3	0.8256	1.5011	0.3302	0.076*
C4	0.7018 (3)	1.3478 (2)	0.3320 (2)	0.0598 (7)
H4	0.6627	1.3784	0.3961	0.072*
C5	0.6594 (2)	1.2303 (2)	0.27708 (19)	0.0452 (6)
C6	0.7213 (2)	1.1851 (2)	0.1834 (2)	0.0480 (6)
H6	0.6935	1.1070	0.1440	0.058*
C7	0.8233 (2)	1.2554 (2)	0.1489 (2)	0.0578 (7)
H7	0.8662	1.2229	0.0884	0.069*
C8	0.5550 (2)	1.1548 (2)	0.32226 (19)	0.0489 (6)
C9	0.4782 (2)	1.0509 (2)	0.23775 (18)	0.0447 (6)

H9	0.4560	1.0668	0.1454	0.054*
C10	0.3734 (2)	0.9763 (2)	0.28795 (19)	0.0453 (6)
H10	0.3694	0.9979	0.3763	0.054*
C11	0.5019 (2)	0.9174 (2)	0.27438 (19)	0.0452 (6)
H11	0.5681	0.9061	0.3552	0.054*
C12	0.5012 (2)	0.8198 (2)	0.1703 (2)	0.0480 (6)
C13	0.4473 (2)	0.6911 (2)	0.1861 (2)	0.0467 (6)
C14	0.4281 (3)	0.6476 (3)	0.3027 (2)	0.0592 (7)
H14	0.4515	0.6999	0.3756	0.071*
C15	0.3747 (3)	0.5276 (3)	0.3114 (2)	0.0684 (8)
H15	0.3643	0.4989	0.3905	0.082*
C16	0.3365 (3)	0.4493 (2)	0.2036 (2)	0.0571 (6)
C17	0.3553 (3)	0.4921 (3)	0.0870 (2)	0.0578 (7)
H17	0.3294	0.4409	0.0135	0.069*
C18	0.4123 (2)	0.6105 (2)	0.0802 (2)	0.0529 (6)
H18	0.4278	0.6372	0.0021	0.064*
C19	0.2147 (3)	0.2623 (3)	0.1091 (3)	0.0829 (9)
H19A	0.1527	0.3131	0.0523	0.124*
H19B	0.1663	0.1912	0.1343	0.124*
H19C	0.2810	0.2334	0.0659	0.124*
C20	0.2404 (2)	0.9402 (2)	0.20001 (19)	0.0437 (6)
C21	0.1269 (2)	0.9056 (2)	0.25933 (19)	0.0425 (5)
C22	-0.0052 (2)	0.9130 (2)	0.1882 (2)	0.0503 (6)
H22	-0.0211	0.9394	0.1037	0.060*
C23	-0.1126 (2)	0.8816 (2)	0.2419 (2)	0.0582 (7)
H23	-0.2003	0.8875	0.1936	0.070*
C24	-0.0906 (2)	0.8411 (2)	0.3674 (2)	0.0538 (6)
C25	0.0391 (2)	0.8319 (3)	0.4392 (2)	0.0593 (7)
H25	0.0545	0.8039	0.5231	0.071*
C26	0.1462 (2)	0.8650 (2)	0.3847 (2)	0.0534 (6)
H26	0.2337	0.8598	0.4337	0.064*
C27	-0.1880 (3)	0.7676 (3)	0.5382 (3)	0.0860 (9)
H27A	-0.1411	0.6906	0.5457	0.129*
H27B	-0.2753	0.7524	0.5563	0.129*
H27C	-0.1372	0.8295	0.5982	0.129*
O2	0.53287 (17)	1.17456 (18)	0.42848 (14)	0.0672 (5)
O1	0.96041 (19)	1.43598 (18)	0.15687 (19)	0.0816 (6)
O3	0.54286 (18)	0.84862 (17)	0.07605 (15)	0.0635 (5)
O4	0.2801 (2)	0.33557 (18)	0.22053 (17)	0.0786 (6)
O5	0.22625 (15)	0.94253 (16)	0.08352 (13)	0.0579 (5)
O6	-0.20364 (17)	0.8128 (2)	0.41048 (18)	0.0797 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.086 (2)	0.058 (2)	0.140 (3)	-0.0223 (17)	0.036 (2)	-0.0041 (19)
C2	0.0474 (14)	0.0519 (16)	0.0689 (15)	-0.0061 (12)	0.0160 (12)	0.0002 (12)
C3	0.0651 (16)	0.0477 (17)	0.0761 (17)	-0.0103 (13)	0.0178 (13)	-0.0154 (13)

C4	0.0622 (15)	0.0631 (18)	0.0557 (14)	-0.0040 (13)	0.0191 (12)	-0.0128 (12)
C5	0.0447 (12)	0.0474 (15)	0.0409 (11)	-0.0032 (11)	0.0057 (9)	0.0003 (10)
C6	0.0504 (13)	0.0439 (14)	0.0487 (12)	-0.0027 (11)	0.0103 (10)	-0.0032 (10)
C7	0.0532 (14)	0.0607 (18)	0.0636 (15)	-0.0030 (13)	0.0233 (12)	-0.0097 (12)
C8	0.0461 (13)	0.0625 (17)	0.0343 (11)	0.0013 (11)	0.0016 (9)	0.0022 (10)
C9	0.0431 (12)	0.0574 (16)	0.0320 (10)	-0.0077 (11)	0.0073 (9)	0.0032 (10)
C10	0.0432 (12)	0.0569 (15)	0.0354 (10)	-0.0034 (11)	0.0094 (9)	-0.0037 (10)
C11	0.0382 (12)	0.0564 (16)	0.0388 (11)	-0.0011 (10)	0.0047 (9)	0.0037 (10)
C12	0.0381 (12)	0.0635 (17)	0.0435 (12)	0.0040 (11)	0.0115 (10)	0.0020 (11)
C13	0.0441 (12)	0.0542 (16)	0.0445 (12)	0.0069 (11)	0.0149 (10)	0.0039 (10)
C14	0.0696 (17)	0.0643 (19)	0.0442 (13)	-0.0035 (14)	0.0154 (11)	0.0006 (11)
C15	0.091 (2)	0.071 (2)	0.0454 (13)	-0.0064 (16)	0.0213 (13)	0.0065 (13)
C16	0.0659 (16)	0.0509 (17)	0.0565 (14)	0.0060 (13)	0.0171 (12)	0.0070 (12)
C17	0.0695 (17)	0.0575 (18)	0.0483 (13)	0.0088 (13)	0.0168 (12)	-0.0048 (12)
C18	0.0595 (15)	0.0570 (17)	0.0471 (13)	0.0121 (13)	0.0205 (11)	0.0042 (11)
C19	0.099 (2)	0.061 (2)	0.086 (2)	-0.0007 (17)	0.0178 (17)	-0.0101 (15)
C20	0.0434 (12)	0.0473 (15)	0.0403 (12)	0.0001 (10)	0.0101 (9)	-0.0047 (10)
C21	0.0397 (12)	0.0444 (14)	0.0428 (11)	-0.0024 (10)	0.0089 (9)	-0.0008 (10)
C22	0.0469 (13)	0.0543 (16)	0.0471 (12)	-0.0035 (11)	0.0063 (10)	0.0066 (11)
C23	0.0384 (13)	0.0705 (18)	0.0630 (15)	0.0004 (12)	0.0056 (11)	0.0106 (13)
C24	0.0415 (13)	0.0589 (17)	0.0637 (15)	0.0017 (11)	0.0169 (11)	0.0124 (12)
C25	0.0526 (15)	0.0770 (19)	0.0490 (13)	0.0006 (13)	0.0128 (11)	0.0148 (12)
C26	0.0381 (12)	0.0714 (18)	0.0478 (13)	-0.0002 (12)	0.0042 (10)	0.0080 (12)
C27	0.0723 (19)	0.111 (3)	0.086 (2)	0.0035 (18)	0.0399 (16)	0.0242 (18)
O2	0.0693 (11)	0.0922 (14)	0.0383 (9)	-0.0202 (10)	0.0137 (8)	-0.0112 (8)
O1	0.0740 (13)	0.0648 (14)	0.1145 (15)	-0.0186 (10)	0.0440 (11)	-0.0110 (11)
O3	0.0715 (11)	0.0723 (13)	0.0538 (9)	-0.0054 (9)	0.0307 (8)	0.0021 (8)
O4	0.1083 (16)	0.0589 (13)	0.0671 (11)	-0.0104 (11)	0.0191 (10)	0.0065 (9)
O5	0.0540 (10)	0.0799 (13)	0.0384 (8)	-0.0060 (9)	0.0097 (7)	-0.0086 (8)
O6	0.0481 (10)	0.1137 (17)	0.0840 (13)	0.0054 (10)	0.0268 (9)	0.0337 (11)

Geometric parameters (Å, °)

C1—O1	1.422 (3)	C14—C15	1.378 (3)
C1—H1A	0.9600	C14—H14	0.9300
C1—H1B	0.9600	C15—C16	1.383 (3)
C1—H1C	0.9600	C15—H15	0.9300
C2—O1	1.360 (3)	C16—O4	1.348 (3)
C2—C7	1.377 (3)	C16—C17	1.384 (3)
C2—C3	1.381 (3)	C17—C18	1.373 (3)
C3—C4	1.383 (3)	C17—H17	0.9300
C3—H3	0.9300	C18—H18	0.9300
C4—C5	1.381 (3)	C19—O4	1.428 (3)
C4—H4	0.9300	C19—H19A	0.9600
C5—C6	1.390 (3)	C19—H19B	0.9600
C5—C8	1.477 (3)	C19—H19C	0.9600
C6—C7	1.376 (3)	C20—O5	1.222 (2)
C6—H6	0.9300	C20—C21	1.477 (3)

C7—H7	0.9300	C21—C26	1.387 (3)
C8—O2	1.225 (2)	C21—C22	1.393 (3)
C8—C9	1.492 (3)	C22—C23	1.378 (3)
C9—C11	1.491 (3)	C22—H22	0.9300
C9—C10	1.503 (3)	C23—C24	1.387 (3)
C9—H9	0.9800	C23—H23	0.9300
C10—C20	1.494 (3)	C24—O6	1.358 (3)
C10—C11	1.510 (3)	C24—C25	1.376 (3)
C10—H10	0.9800	C25—C26	1.384 (3)
C11—C12	1.510 (3)	C25—H25	0.9300
C11—H11	0.9800	C26—H26	0.9300
C12—O3	1.218 (2)	C27—O6	1.430 (3)
C12—C13	1.478 (3)	C27—H27A	0.9600
C13—C18	1.382 (3)	C27—H27B	0.9600
C13—C14	1.387 (3)	C27—H27C	0.9600
O1—C1—H1A	109.5	C15—C14—C13	120.5 (2)
O1—C1—H1B	109.5	C15—C14—H14	119.8
H1A—C1—H1B	109.5	C13—C14—H14	119.8
O1—C1—H1C	109.5	C14—C15—C16	120.7 (2)
H1A—C1—H1C	109.5	C14—C15—H15	119.6
H1B—C1—H1C	109.5	C16—C15—H15	119.6
O1—C2—C7	115.4 (2)	O4—C16—C15	116.4 (2)
O1—C2—C3	124.8 (2)	O4—C16—C17	124.5 (2)
C7—C2—C3	119.8 (2)	C15—C16—C17	119.1 (2)
C2—C3—C4	119.1 (2)	C18—C17—C16	119.7 (2)
C2—C3—H3	120.5	C18—C17—H17	120.2
C4—C3—H3	120.5	C16—C17—H17	120.2
C5—C4—C3	121.8 (2)	C17—C18—C13	121.8 (2)
C5—C4—H4	119.1	C17—C18—H18	119.1
C3—C4—H4	119.1	C13—C18—H18	119.1
C4—C5—C6	118.1 (2)	O4—C19—H19A	109.5
C4—C5—C8	119.8 (2)	O4—C19—H19B	109.5
C6—C5—C8	122.0 (2)	H19A—C19—H19B	109.5
C7—C6—C5	120.3 (2)	O4—C19—H19C	109.5
C7—C6—H6	119.8	H19A—C19—H19C	109.5
C5—C6—H6	119.8	H19B—C19—H19C	109.5
C6—C7—C2	120.8 (2)	O5—C20—C21	121.68 (18)
C6—C7—H7	119.6	O5—C20—C10	120.80 (19)
C2—C7—H7	119.6	C21—C20—C10	117.50 (17)
O2—C8—C5	121.07 (19)	C26—C21—C22	117.9 (2)
O2—C8—C9	119.9 (2)	C26—C21—C20	122.47 (19)
C5—C8—C9	119.03 (18)	C22—C21—C20	119.67 (19)
C11—C9—C8	119.55 (18)	C23—C22—C21	120.6 (2)
C11—C9—C10	60.57 (15)	C23—C22—H22	119.7
C8—C9—C10	117.15 (17)	C21—C22—H22	119.7
C11—C9—H9	116.0	C22—C23—C24	120.4 (2)
C8—C9—H9	116.0	C22—C23—H23	119.8

C10—C9—H9	116.0	C24—C23—H23	119.8
C20—C10—C9	119.72 (17)	O6—C24—C25	124.6 (2)
C20—C10—C11	121.69 (18)	O6—C24—C23	115.4 (2)
C9—C10—C11	59.33 (14)	C25—C24—C23	120.0 (2)
C20—C10—H10	114.9	C24—C25—C26	119.1 (2)
C9—C10—H10	114.9	C24—C25—H25	120.5
C11—C10—H10	114.9	C26—C25—H25	120.5
C9—C11—C10	60.10 (15)	C25—C26—C21	122.0 (2)
C9—C11—C12	118.96 (18)	C25—C26—H26	119.0
C10—C11—C12	121.17 (17)	C21—C26—H26	119.0
C9—C11—H11	115.2	O6—C27—H27A	109.5
C10—C11—H11	115.2	O6—C27—H27B	109.5
C12—C11—H11	115.2	H27A—C27—H27B	109.5
O3—C12—C13	121.9 (2)	O6—C27—H27C	109.5
O3—C12—C11	119.9 (2)	H27A—C27—H27C	109.5
C13—C12—C11	118.13 (19)	H27B—C27—H27C	109.5
C18—C13—C14	118.1 (2)	C2—O1—C1	119.2 (2)
C18—C13—C12	119.0 (2)	C16—O4—C19	118.2 (2)
C14—C13—C12	122.9 (2)	C24—O6—C27	118.2 (2)
O1—C2—C3—C4	179.4 (2)	C18—C13—C14—C15	-0.5 (4)
C7—C2—C3—C4	1.2 (4)	C12—C13—C14—C15	178.6 (2)
C2—C3—C4—C5	-2.5 (4)	C13—C14—C15—C16	-1.4 (4)
C3—C4—C5—C6	1.4 (4)	C14—C15—C16—O4	-177.3 (2)
C3—C4—C5—C8	178.9 (2)	C14—C15—C16—C17	1.4 (4)
C4—C5—C6—C7	1.1 (3)	O4—C16—C17—C18	179.2 (2)
C8—C5—C6—C7	-176.4 (2)	C15—C16—C17—C18	0.6 (4)
C5—C6—C7—C2	-2.3 (4)	C16—C17—C18—C13	-2.5 (4)
O1—C2—C7—C6	-177.2 (2)	C14—C13—C18—C17	2.5 (4)
C3—C2—C7—C6	1.1 (4)	C12—C13—C18—C17	-176.6 (2)
C4—C5—C8—O2	-21.4 (3)	C9—C10—C20—O5	18.0 (3)
C6—C5—C8—O2	156.0 (2)	C11—C10—C20—O5	-52.3 (3)
C4—C5—C8—C9	160.2 (2)	C9—C10—C20—C21	-160.3 (2)
C6—C5—C8—C9	-22.4 (3)	C11—C10—C20—C21	129.4 (2)
O2—C8—C9—C11	-68.5 (3)	O5—C20—C21—C26	159.8 (2)
C5—C8—C9—C11	109.9 (2)	C10—C20—C21—C26	-21.9 (3)
O2—C8—C9—C10	1.3 (3)	O5—C20—C21—C22	-20.1 (3)
C5—C8—C9—C10	179.8 (2)	C10—C20—C21—C22	158.2 (2)
C11—C9—C10—C20	-111.4 (2)	C26—C21—C22—C23	0.4 (3)
C8—C9—C10—C20	138.3 (2)	C20—C21—C22—C23	-179.7 (2)
C8—C9—C10—C11	-110.3 (2)	C21—C22—C23—C24	-0.5 (4)
C8—C9—C11—C10	106.4 (2)	C22—C23—C24—O6	179.9 (2)
C8—C9—C11—C12	-142.2 (2)	C22—C23—C24—C25	-0.1 (4)
C10—C9—C11—C12	111.4 (2)	O6—C24—C25—C26	-179.3 (2)
C20—C10—C11—C9	108.1 (2)	C23—C24—C25—C26	0.7 (4)
C20—C10—C11—C12	0.4 (3)	C24—C25—C26—C21	-0.8 (4)
C9—C10—C11—C12	-107.8 (2)	C22—C21—C26—C25	0.3 (4)
C9—C11—C12—O3	33.7 (3)	C20—C21—C26—C25	-179.6 (2)

C10—C11—C12—O3	104.4 (3)	C7—C2—O1—C1	-179.9 (3)
C9—C11—C12—C13	-145.5 (2)	C3—C2—O1—C1	1.9 (4)
C10—C11—C12—C13	-74.8 (3)	C15—C16—O4—C19	167.5 (2)
O3—C12—C13—C18	-16.1 (3)	C17—C16—O4—C19	-11.1 (4)
C11—C12—C13—C18	163.0 (2)	C25—C24—O6—C27	-1.4 (4)
O3—C12—C13—C14	164.8 (2)	C23—C24—O6—C27	178.6 (3)
C11—C12—C13—C14	-16.0 (3)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C26—H26...O2 ⁱ	0.93	2.54	3.467 (3)	175
C11—H11...O6 ⁱⁱ	0.98	2.51	3.270 (3)	134
C7—H7...O5 ⁱⁱⁱ	0.93	2.52	3.171 (3)	128
C9—H9...O3 ⁱⁱⁱ	0.98	2.55	3.497 (3)	164
C27—H27B...O4 ^{iv}	0.96	2.58	3.136 (3)	117

Symmetry codes: (i) $-x+1, -y+2, -z+1$; (ii) $x+1, y, z$; (iii) $-x+1, -y+2, -z$; (iv) $-x, -y+1, -z+1$.