

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Bis(2,5-dimethylanilinium) tetrachloridozincate(II)

Sofiane Souissi, Wajda Smirani* and Mohamed Rzaigui

Laboratoire de Chimie des Matériaux, Faculté des Sciences de Bizerte, 7021

Zarzouna Bizerte, Tunisia

Correspondence e-mail: wajda_sta@yahoo.fr


Received 17 March 2009; accepted 18 March 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; R factor = 0.079; wR factor = 0.229; data-to-parameter ratio = 18.4.

In the title compound, $(\text{C}_8\text{H}_{12}\text{N})_2[\text{ZnCl}_4]$, the Zn^{2+} ion adopts a distorted tetrahedral coordination geometry. In the crystal, the cations and anions are linked by $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds, leading to ribbons propagating parallel to the a axis.

Related literature

For related structures, see: Guo *et al.* (2007); Smirani & Rzaigui (2009). For background on hybrid materials, see: Tao *et al.* (2003); Bringley & Rajeswaran (2006).



Experimental

Crystal data

 $(\text{C}_8\text{H}_{12}\text{N})_2[\text{ZnCl}_4]$ $M_r = 451.58$ Monoclinic, $P2_1/n$ $a = 7.425$ (2) Å $b = 12.884$ (2) Å $c = 22.809$ (2) Å $\beta = 96.16$ (2)° $V = 2169.5$ (7) Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 1.62$ mm⁻¹ $T = 293$ K $0.20 \times 0.13 \times 0.10$ mm

Data collection

Enraf–Nonius Turbo CAD-4 diffractometer

Absorption correction: none
6539 measured reflections3947 independent reflections
2621 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.033$ 2 standard reflections
frequency: 120 min
intensity decay: 5%

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.079$ $wR(F^2) = 0.229$ $S = 1.05$

3947 reflections

214 parameters

H-atom parameters not refined

 $\Delta\rho_{\text{max}} = 0.50$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.61$ e Å⁻³

Table 1

Selected bond lengths (Å).

Zn1—Cl1	2.248 (2)	Zn1—Cl3	2.274 (2)
Zn1—Cl2	2.2502 (16)	Zn1—Cl4	2.2721 (18)

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1A \cdots Cl4 ⁱ	0.89	2.25	3.125 (6)	169
N1—H1B \cdots Cl3 ⁱⁱ	0.89	2.54	3.304 (6)	145
N1—H1C \cdots Cl2 ⁱⁱⁱ	0.89	2.31	3.172 (7)	162
N2—H2A \cdots Cl1 ⁱ	0.89	2.34	3.219 (6)	171
N2—H2B \cdots Cl4 ^{iv}	0.89	2.70	3.505 (7)	151
N2—H2C \cdots Cl3 ⁱⁱ	0.89	2.39	3.262 (6)	168

Symmetry codes: (i) $x, y + 1, z$; (ii) $-x, -y + 1, -z$; (iii) $-x + 1, -y + 1, -z$; (iv) $x - 1, y + 1, z$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB2930).

References

- Bringley, J. F. & Rajeswaran, M. (2006). *Acta Cryst.* **E62**, m1304–m1305.
Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
Guo, N., Yi, J., Chen, Y., Liao, S. & Fu, Z. (2007). *Acta Cryst.* **E63**, m2571.
Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Smirani, W. & Rzaigui, M. (2009). *Acta Cryst.* **E65**, o83.
Tao, J., Yin, X., Jiang, Y. B., Yang, L. F., Huang, R. B. & Zheng, L. S. (2003). *Eur. J. Inorg. Chem.* pp. 2678–2682.

supporting information

Acta Cryst. (2009). E65, m442 [doi:10.1107/S1600536809009982]

Bis(2,5-dimethylanilinium) tetrachloridozincate(II)

Sofiane Souissi, Wajda Smirani and Mohamed Rzaigui

S1. Comment

Inorganic-organic hybrid materials are of great interest in solid state chemistry due to their enormous variety of intriguing structural topologies and their fascinating properties as well as great potential applications in many fields (Tao *et al.*, 2003; Bringley & Rajeswaran, 2006). Here we report the crystal structure of bis(2,5-xylidinium) tetrachlorozincate (I).

As shown in Fig. 1, the asymmetric unit of (I) is built up from two 2,5-xylidinium cations and one tetrachlorozincate (II) anion. The Zn (II) ion is in a tetrahedral coordination environment composed of four Cl anions (Table 1). The Cl—Zn—Cl bond angles range from 106.13 (8) to 112.46 (8)°. These values indicate that the anionic $[\text{ZnCl}_4]^{2-}$ tetrahedron is slightly distorted (Guo *et al.*, 2007). The examination of the organic cation shows that the values of the N—C, C—C distances and N—C—C, C—C—C angles range from 1.343 (12) to 1.512 (11) Å and 115.90 (7) to 123.0 (6)°, respectively. These values show no significant difference from those obtained in other crystals involving the same organic groups (Smirani and Rzaigui, 2009).


A projection of the structure along the direction *a* shows that the $[\text{ZnCl}_4]^{2-}$ anions are connected *via* N—H...Cl hydrogen bonds originating from NH_3^+ groups, so as to built inorganic ribbons at $x = 0$ and $x = 1/2$ (Fig. 2, Table 2). The 2,5-xylidinium cations are anchored onto the successive ribbons *via* hydrogen bonds and electrostatic and van der Waals interactions, to compensate their negative charges.

S2. Experimental


An aqueous solution of 2,5-xylidine, HCl and ZnCl_2 in a 2:2:1 molar ratio was prepared and colourless blocks of (I) grew as the water evaporated over the course of a few days.

S3. Refinement

All H atoms were positioned geometrically and treated as riding on their parent atoms: N—H = 0.89, C—H = 0.93–0.96 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl-C,N})$.

**Figure 1**

View of the molecular structure of (I): displacement ellipsoids for non-H atoms are drawn at the 30% probability level.

**Figure 2**

A view of the packing in (I) viewed along the *a* axis.

Bis(2,5-dimethylanilinium) tetrachloridozincate(II)

Crystal data

$(C_8H_{12}N)_2[ZnCl_4]$

$M_r = 451.58$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 7.425$ (2) Å
 $b = 12.884$ (2) Å
 $c = 22.809$ (2) Å
 $\beta = 96.16$ (2)°
 $V = 2169.5$ (7) Å³
 $Z = 4$
 $F(000) = 928$
 $D_x = 1.383$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 25 reflections
 $\theta = 9.9$ – 11.0 °
 $\mu = 1.62$ mm⁻¹
 $T = 293$ K
 Block, colourless
 $0.20 \times 0.13 \times 0.10$ mm

Data collection

Enraf–Nonius Turbo CAD-4
 diffractometer
 Radiation source: Enraf Nonius FR590
 Nonprofiled ω scans
 6539 measured reflections
 3947 independent reflections
 2621 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.033$
 $\theta_{\text{max}} = 28.0$ °, $\theta_{\text{min}} = 2.4$ °
 $h = -9 \rightarrow 9$
 $k = 0 \rightarrow 17$
 $l = -10 \rightarrow 17$
 2 standard reflections every 120 min
 intensity decay: 5%

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.079$
 $wR(F^2) = 0.229$
 $S = 1.05$
 3947 reflections
 214 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters not refined
 $w = 1/[\sigma^2(F_o^2) + (0.1531P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.012$
 $\Delta\rho_{\text{max}} = 0.50$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.61$ e Å⁻³

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.0360 (2)	0.16857 (16)	0.12060 (10)	0.0688 (6)
C12	0.52806 (19)	0.20922 (13)	0.10654 (9)	0.0537 (5)
C13	0.1904 (2)	0.22519 (13)	-0.02438 (9)	0.0583 (6)
C14	0.3103 (3)	-0.02537 (13)	0.04807 (11)	0.0808 (7)
N1	0.2378 (7)	0.7499 (4)	-0.0004 (3)	0.0562 (16)
H1A	0.2629	0.8096	0.0183	0.084*
H1B	0.1189	0.7451	-0.0105	0.084*
H1C	0.2950	0.7474	-0.0328	0.084*
C1	0.2981 (7)	0.6631 (4)	0.0386 (3)	0.0414 (17)
C3	0.3769 (8)	0.6844 (4)	0.0940 (3)	0.0490 (19)

H3	0.3922	0.7531	0.1061	0.059*
C4	0.4345 (8)	0.6053 (5)	0.1327 (3)	0.0527 (18)
C2	0.2704 (7)	0.5621 (4)	0.0171 (3)	0.0407 (16)
C5	0.3295 (8)	0.4834 (4)	0.0569 (3)	0.0513 (19)
H5	0.3153	0.4146	0.0451	0.062*
C9	0.1839 (10)	0.5398 (6)	-0.0438 (3)	0.062 (2)
H9A	0.0653	0.5703	-0.0490	0.093*
H9B	0.1742	0.4661	-0.0495	0.093*
H9C	0.2567	0.5686	-0.0721	0.093*
C8	0.4080 (9)	0.5049 (5)	0.1130 (4)	0.0530 (19)
H8	0.4441	0.4502	0.1382	0.064*
C7	0.5214 (14)	0.6282 (7)	0.1943 (4)	0.086 (3)
H7A	0.6431	0.6522	0.1925	0.129*
H7B	0.5235	0.5661	0.2177	0.129*
H7C	0.4529	0.6808	0.2119	0.129*
N2	-0.2368 (8)	0.9737 (4)	0.1100 (3)	0.0565 (17)
H2A	-0.1525	1.0231	0.1114	0.085*
H2B	-0.3429	0.9999	0.0952	0.085*
H2C	-0.2061	0.9220	0.0872	0.085*
C10	-0.2512 (9)	0.9346 (4)	0.1696 (3)	0.0456 (17)
C13	-0.1163 (9)	0.8706 (5)	0.1964 (3)	0.0483 (19)
C11	-0.3993 (9)	0.9637 (6)	0.1974 (4)	0.062 (2)
H11	-0.4863	1.0074	0.1782	0.075*
C14	0.0465 (11)	0.8399 (7)	0.1675 (4)	0.082 (3)
H14A	0.0186	0.7799	0.1433	0.123*
H14B	0.1443	0.8240	0.1971	0.123*
H14C	0.0813	0.8961	0.1434	0.123*
C12	-0.4195 (10)	0.9281 (7)	0.2541 (4)	0.065 (2)
C16	-0.2860 (11)	0.8619 (6)	0.2797 (4)	0.062 (2)
H16	-0.2972	0.8346	0.3169	0.075*
C15	-0.1412 (11)	0.8360 (6)	0.2523 (4)	0.062 (2)
H15	-0.0538	0.7929	0.2718	0.075*
C17	-0.5803 (14)	0.9593 (10)	0.2849 (5)	0.120 (4)
H17A	-0.6893	0.9349	0.2627	0.180*
H17B	-0.5844	1.0335	0.2879	0.180*
H17C	-0.5701	0.9293	0.3236	0.180*
Zn1	0.26302 (8)	0.14604 (5)	0.06412 (4)	0.0418 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C11	0.0506 (9)	0.0874 (13)	0.0709 (18)	0.0007 (8)	0.0174 (9)	-0.0083 (11)
C12	0.0423 (7)	0.0579 (9)	0.0584 (15)	-0.0086 (6)	-0.0064 (7)	-0.0055 (8)
C13	0.0709 (10)	0.0511 (9)	0.0506 (16)	-0.0023 (7)	-0.0040 (9)	0.0047 (8)
C14	0.1136 (16)	0.0354 (8)	0.0909 (19)	-0.0001 (9)	-0.0004 (13)	-0.0092 (9)
N1	0.067 (3)	0.039 (3)	0.063 (5)	0.003 (2)	0.004 (3)	0.008 (3)
C1	0.036 (3)	0.033 (3)	0.056 (6)	0.003 (2)	0.012 (3)	0.003 (3)
C3	0.047 (3)	0.036 (3)	0.063 (6)	0.000 (2)	0.001 (3)	-0.004 (3)

C4	0.048 (3)	0.052 (3)	0.057 (6)	-0.001 (3)	-0.001 (3)	0.004 (3)
C2	0.037 (3)	0.041 (3)	0.046 (5)	-0.004 (2)	0.012 (3)	-0.001 (3)
C5	0.047 (3)	0.033 (3)	0.076 (6)	-0.001 (2)	0.011 (3)	0.005 (3)
C9	0.066 (4)	0.056 (4)	0.064 (7)	-0.008 (3)	0.006 (4)	-0.012 (4)
C8	0.053 (3)	0.042 (3)	0.063 (6)	0.005 (3)	0.000 (4)	0.008 (3)
C7	0.101 (7)	0.075 (5)	0.077 (8)	-0.001 (5)	-0.014 (6)	0.001 (5)
N2	0.075 (4)	0.054 (3)	0.041 (5)	0.004 (3)	0.005 (3)	0.008 (3)
C10	0.060 (4)	0.047 (3)	0.029 (5)	-0.006 (3)	0.000 (3)	0.001 (3)
C13	0.057 (4)	0.053 (3)	0.033 (6)	-0.003 (3)	-0.004 (3)	0.001 (3)
C11	0.058 (4)	0.069 (4)	0.058 (7)	0.013 (3)	-0.003 (4)	0.000 (4)
C14	0.068 (5)	0.109 (7)	0.067 (8)	0.026 (5)	0.002 (5)	-0.002 (5)
C12	0.067 (4)	0.087 (5)	0.042 (7)	-0.006 (4)	0.012 (4)	-0.002 (4)
C16	0.075 (5)	0.076 (5)	0.034 (6)	-0.009 (4)	-0.001 (4)	0.009 (4)
C15	0.068 (5)	0.065 (4)	0.051 (7)	0.006 (3)	-0.008 (4)	0.007 (4)
C17	0.085 (6)	0.195 (13)	0.082 (9)	0.028 (7)	0.020 (6)	-0.011 (8)
Zn1	0.0383 (4)	0.0382 (4)	0.0481 (8)	-0.0011 (3)	0.0007 (3)	-0.0031 (3)

Geometric parameters (Å, °)

Zn1—C11	2.248 (2)	C7—H7B	0.9600
Zn1—C12	2.2502 (16)	C7—H7C	0.9600
Zn1—C13	2.274 (2)	N2—C10	1.463 (9)
Zn1—C14	2.2721 (18)	N2—H2A	0.8900
N1—C1	1.468 (8)	N2—H2B	0.8900
N1—H1A	0.8900	N2—H2C	0.8900
N1—H1B	0.8900	C10—C11	1.379 (10)
N1—H1C	0.8900	C10—C13	1.388 (9)
C1—C3	1.364 (10)	C13—C15	1.382 (11)
C1—C2	1.399 (8)	C13—C14	1.491 (11)
C3—C4	1.386 (9)	C11—C12	1.396 (11)
C3—H3	0.9300	C11—H11	0.9300
C4—C8	1.377 (9)	C14—H14A	0.9600
C4—C7	1.512 (11)	C14—H14B	0.9600
C2—C5	1.400 (9)	C14—H14C	0.9600
C2—C9	1.495 (10)	C12—C16	1.388 (11)
C5—C8	1.376 (10)	C12—C17	1.503 (13)
C5—H5	0.9300	C16—C15	1.343 (12)
C9—H9A	0.9600	C16—H16	0.9300
C9—H9B	0.9600	C15—H15	0.9300
C9—H9C	0.9600	C17—H17A	0.9600
C8—H8	0.9300	C17—H17B	0.9600
C7—H7A	0.9600	C17—H17C	0.9600
C1—N1—H1A	109.5	C10—N2—H2C	109.5
C1—N1—H1B	109.5	H2A—N2—H2C	109.5
H1A—N1—H1B	109.5	H2B—N2—H2C	109.5
C1—N1—H1C	109.5	C11—C10—C13	122.2 (7)
H1A—N1—H1C	109.5	C11—C10—N2	118.4 (6)

H1B—N1—H1C	109.5	C13—C10—N2	119.5 (6)
C3—C1—C2	123.0 (6)	C15—C13—C10	115.9 (7)
C3—C1—N1	118.8 (5)	C15—C13—C14	121.2 (7)
C2—C1—N1	118.2 (6)	C10—C13—C14	122.9 (7)
C1—C3—C4	121.0 (6)	C10—C11—C12	120.5 (7)
C1—C3—H3	119.5	C10—C11—H11	119.8
C4—C3—H3	119.5	C12—C11—H11	119.8
C8—C4—C3	117.4 (7)	C13—C14—H14A	109.5
C8—C4—C7	121.2 (7)	C13—C14—H14B	109.5
C3—C4—C7	121.3 (7)	H14A—C14—H14B	109.5
C1—C2—C5	115.0 (6)	C13—C14—H14C	109.5
C1—C2—C9	122.5 (6)	H14A—C14—H14C	109.5
C5—C2—C9	122.5 (6)	H14B—C14—H14C	109.5
C8—C5—C2	122.0 (6)	C16—C12—C11	116.7 (7)
C8—C5—H5	119.0	C16—C12—C17	122.3 (9)
C2—C5—H5	119.0	C11—C12—C17	121.0 (8)
C2—C9—H9A	109.5	C15—C16—C12	121.8 (8)
C2—C9—H9B	109.5	C15—C16—H16	119.1
H9A—C9—H9B	109.5	C12—C16—H16	119.1
C2—C9—H9C	109.5	C16—C15—C13	122.9 (7)
H9A—C9—H9C	109.5	C16—C15—H15	118.5
H9B—C9—H9C	109.5	C13—C15—H15	118.5
C5—C8—C4	121.6 (6)	C12—C17—H17A	109.5
C5—C8—H8	119.2	C12—C17—H17B	109.5
C4—C8—H8	119.2	H17A—C17—H17B	109.5
C4—C7—H7A	109.5	C12—C17—H17C	109.5
C4—C7—H7B	109.5	H17A—C17—H17C	109.5
H7A—C7—H7B	109.5	H17B—C17—H17C	109.5
C4—C7—H7C	109.5	Cl1—Zn1—Cl2	112.46 (8)
H7A—C7—H7C	109.5	Cl1—Zn1—Cl4	110.87 (9)
H7B—C7—H7C	109.5	Cl2—Zn1—Cl4	106.13 (8)
C10—N2—H2A	109.5	Cl1—Zn1—Cl3	109.26 (8)
C10—N2—H2B	109.5	Cl2—Zn1—Cl3	109.42 (7)
H2A—N2—H2B	109.5	Cl4—Zn1—Cl3	108.59 (9)
C2—C1—C3—C4	-0.1 (9)	C11—C10—C13—C15	1.3 (10)
N1—C1—C3—C4	179.4 (6)	N2—C10—C13—C15	-179.2 (6)
C1—C3—C4—C8	-0.4 (10)	C11—C10—C13—C14	-178.8 (7)
C1—C3—C4—C7	-179.8 (7)	N2—C10—C13—C14	0.7 (10)
C3—C1—C2—C5	0.1 (8)	C13—C10—C11—C12	-0.7 (11)
N1—C1—C2—C5	-179.3 (5)	N2—C10—C11—C12	179.7 (6)
C3—C1—C2—C9	-180.0 (6)	C10—C11—C12—C16	-1.0 (11)
N1—C1—C2—C9	0.6 (9)	C10—C11—C12—C17	180.0 (9)
C1—C2—C5—C8	0.3 (9)	C11—C12—C16—C15	2.2 (12)
C9—C2—C5—C8	-179.6 (6)	C17—C12—C16—C15	-178.7 (9)
C2—C5—C8—C4	-0.8 (10)	C12—C16—C15—C13	-1.8 (12)
C3—C4—C8—C5	0.8 (10)	C10—C13—C15—C16	0.0 (11)
C7—C4—C8—C5	-179.8 (7)	C14—C13—C15—C16	-179.9 (8)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1A \cdots C14 ⁱ	0.89	2.25	3.125 (6)	169
N1—H1B \cdots C13 ⁱⁱ	0.89	2.54	3.304 (6)	145
N1—H1C \cdots C12 ⁱⁱⁱ	0.89	2.31	3.172 (7)	162
N2—H2A \cdots C11 ⁱ	0.89	2.34	3.219 (6)	171
N2—H2B \cdots C14 ^{iv}	0.89	2.70	3.505 (7)	151
N2—H2C \cdots C13 ⁱⁱ	0.89	2.39	3.262 (6)	168

Symmetry codes: (i) $x, y+1, z$; (ii) $-x, -y+1, -z$; (iii) $-x+1, -y+1, -z$; (iv) $x-1, y+1, z$.