

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

7,7',8,8'-Tetramethoxy-4,4'-dimethyl-3,3'-bicomarin

Hoong-Kun Fun,^{a*} Samuel Robinson Jebas,^{a§} Mehtab Parveen,^b Zakia Khanam^b and Raza Murad Ghalib^b^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 Universiti Sains Malaysia, Penang, Malaysia, and ^bDepartment of Chemistry, Aligarh Muslim University, Aligarh 202002 (UP), India

Correspondence e-mail: hkfun@usm.my


Received 5 May 2009; accepted 8 May 2009

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.059; wR factor = 0.155; data-to-parameter ratio = 18.7.

In the crystal structure, the title compound, $\text{C}_{24}\text{H}_{22}\text{O}_8$, lies on a twofold rotation axis and the asymmetric unit comprises one half-molecule. The dihedral angle formed by the coumarin unit with the symmetry-related part is 74.78 (14)°. One of the methoxy groups attached to the coumarin unit is considerably twisted, making an angle of 87.17 (17)° with respect to the coumarin unit; the other is twisted by 0.66 (19)°. No classical hydrogen bonds are found in the structure; only a weak $\text{C}-\text{H}\cdots\pi$ interaction and short intramolecular $\text{O}\cdots\text{O}$ contacts [2.683 (2)–2.701 (2) Å] are observed.

Related literature

For the biological activity of coumarins, see: El-Agrody *et al.* (2001); El-Faragy (1991); Emmanuel-Giota *et al.* (2001); Ghate *et al.* (2005); Laakso *et al.* (1994); Nofal *et al.* (2000); Pratibha & Shreeya (1999); Shaker (1996); Yang *et al.* (2005). For the pharmaceutical properties of coumarin derivatives, see: Kennedy & Thornes (1997). For natural and synthetic coumarins, see: Carlton *et al.* (1996); Zhou *et al.* (2000). For related bond-length data, see: Allen *et al.* (1987). For stability of the temperature controller used in the data collection, see: Cosier & Glazer (1986).



* Thomson Reuters Researcher ID: A-3561-2009.

§ Thomson Reuters Researcher ID: A-5473-2009. Permanent address: Department of Physics, Karunya University, Karunya Nagar, Coimbatore 641 114, India.

Experimental

Crystal data

$\text{C}_{24}\text{H}_{22}\text{O}_8$
 $M_r = 438.42$
 Monoclinic, $C2/c$
 $a = 21.715$ (9) Å
 $b = 7.138$ (3) Å
 $c = 15.511$ (6) Å
 $\beta = 121.801$ (5)°

$V = 2043.3$ (14) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.11$ mm⁻¹
 $T = 100$ K
 $0.28 \times 0.19 \times 0.06$ mm

Data collection

Bruker SMART APEXII CCD
 area-detector diffractometer
 Absorption correction: multi-scan
 (SADABS; Bruker, 2005)
 $T_{\min} = 0.971$, $T_{\max} = 0.994$

27961 measured reflections
 3527 independent reflections
 2710 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.065$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.059$
 $wR(F^2) = 0.155$
 $S = 1.08$
 3527 reflections

189 parameters
 All H-atom parameters refined
 $\Delta\rho_{\text{max}} = 0.50$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.21$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C6}-\text{H6}\cdots\text{Cg1}^i$	0.96 (2)	2.86 (2)	3.676 (2)	143.5 (18)

Symmetry code: (i) $x, -y, z - \frac{1}{2}$. Cg1 is the centroid of the C4–C9 ring.

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL and PLATON (Spek, 2009).

HKF and SRJ thank the Malaysian Government and Universiti Sains Malaysia for the Science Fund grant No. 305/PFIZIK/613312. SRJ thanks Universiti Sains Malaysia for a post-doctoral research fellowship. HKF also thanks Universiti Sains Malaysia for the Research University Golden Goose grant No. 1001/PFIZIK/811012.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS2417).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–S19.
- Bruker (2005). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Carlton, B. D., Aubrun, J. C. & Simon, G. S. (1996). *Fundam. Appl. Toxicol.* **30**, 145–151.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst.* **19**, 105–107.
- El-Agrody, A. M., Abd El-Latif, M. S., El-Hady, N. A., Fakery, A. H. & Bedair, A. H. (2001). *Molecules*, **6**, 519–527.
- El-Faragy, A. F. (1991). *Egypt. J. Pharm. Sci.* **32**, 625–625.
- Emmanuel-Giota, A. A., Fylaktakidou, K. C., Hadjipavlou-Litina, D. J., Litinas, K. E. & Nicolaidis, D. N. (2001). *J. Heterocycl. Chem.* **38**, 3, 717–722.
- Ghate, M., Kusanur, R. A. & Kulkarni, M. V. (2005). *Eur. J. Med. Chem.* **40**, 882–887.

- Kennedy, R. O. & Thornes, R. D. (1997). *Coumarins: Biology, Applications and Mode of Action*. New York: John Wiley & Sons.
- Laakso, J. A., Narske, E. D., Gloer, J. B., Wicklow, D. T. & Dowd, P. F. (1994). *J. Nat. Prod.*, **57**, 128–133.
- Nofal, Z. M., El-Zahar, M. & Abd El-Karim, S. (2000). *Molecules*, **5**, 99–113.
- Pratibha, S. & Shreeya, P. (1999). *Indian J. Chem.* **38B**, 1139–1142.
- Shaker, R. M. (1996). *Pharmazie*, **51**, 148–148.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Yang, H., Protiva, P., Gil, R. R., Jiang, B., Baggett, S., Basile, M. J., Reynertson, K. A., Weinstein, I. B. & Kennelly, E. J. (2005). *Planta Med.* **71**, 852–60.
- Zhou, P., Takaishi, Y. & Duan, H. (2000). *Phytochemistry*, **53**, 689–697.

supporting information

Acta Cryst. (2009). E65, o1294–o1295 [doi:10.1107/S1600536809017334]

7,7',8,8'-Tetramethoxy-4,4'-dimethyl-3,3'-bicoumarin

Hoong-Kun Fun, Samuel Robinson Jebas, Mehtab Parveen, Zakia Khanam and Raza Murad Ghalib

S1. Comment

Coumarins are a large group of naturally occurring oxygen heterocycles representing 2*H*-1-benzopyran-2-one derivative. Many natural coumarins are reputed for their wide range of biological activities such as antibacterial (El-Agrody *et al.*, 2001; Pratibha & Shreeya, 1999), antifungal (Shaker, 1996; El-Faragy, 1991), antioxidant (Yang *et al.*, 2005), analgesic (Ghate *et al.*, 2005), anti-inflammatory (Emmanuel-Giota *et al.*, 2001) and antitumor (Nofal *et al.*, 2000) properties. Bi and tri-coumarins are comparatively new groups which are widely spread in nature and their biological properties are also well known (Laakso *et al.*, 1994). One of the characteristic pharmacological properties of coumarin derivatives is the anticoagulant action (Kennedy & Thornes, 1997). A large number of natural and semisynthetic coumarin and bicoumarin derivatives have been reported to demonstrate chemopreventive (Carlton *et al.*, 1996) and anti-HIV (Zhou *et al.*, 2000) activities. Keeping in view of these biological importance of coumarins and their dimers, we have synthesized the title compound (I) and report here its structure.

The asymmetric unit of (I) (Fig. 1), contains half of the 7,7',8,8'-4,4'-dimethyl-3,3'-bicoumarin molecule. The other half is symmetry generated [symmetry code: $-x, y, -z + 1/2$]. The coumarin unit is planar with the maximum deviation from the mean plane of 0.0295 (15) Å for atom C2. One of the methyl group attached to the coumarin unit is twisted as evidenced by the torsion angle of C10—O3—C8—C9 = 87.17 (17)°. The dihedral angle formed by the coumarin unit (O1/C1—C9) with the symmetry related coumarin unit (O1A/C1A—C9A) is 74.78 (14)°, indicating that they are almost perpendicular to each other. The bond lengths (Allen *et al.*, 1987) and bond angles are normal.


The crystal packing (Fig. 2) (Table 1) is stabilized by weak C—H \cdots π interactions and intramolecular O \cdots O = 2.683 (2) to 2.701 (2) Å short contacts.

S2. Experimental


A mixture of 7,8-dimethoxy-4-methyl coumarin (2.20 g, 10 mmol) and manganese(III) acetate (0.774 g, 1 mmol) was stirred at room temperature, then 70% perchloric acid (0.8 g, 6 mmol) was added. The reaction mixture was heated under reflux at 114°C with stirring in the atmosphere of nitrogen for 3 h. The reaction mixture was cooled and diluted with 50 ml of benzene. The benzene solution was washed with water and aq. NaHCO₃, dried over anhydrous Na₂SO₄ and left to evaporate. The residue showed two major compounds which were separated by column chromatography followed by preparative thin layer chromatography (Benzene: EtOAc, 9:1) into the title compound (I) (260 mg, 12%).

S3. Refinement

All the hydrogen atoms were located from the Fourier map and allowed to refine freely.

**Figure 1**

The molecular structure of (I), showing 50% probability displacement ellipsoids and the atom numbering scheme. [Symmetry code: $-x, y, -z + 1/2$ to generate equivalent atoms].

**Figure 2**

The crystal packing of (I). Molecules are stacked along the *b* axis.

(I)*Crystal data* $C_{24}H_{22}O_8$ $M_r = 438.42$ Monoclinic, $C2/c$ Hall symbol: $-C 2yc$ $a = 21.715 (9) \text{ \AA}$ $b = 7.138 (3) \text{ \AA}$ $c = 15.511 (6) \text{ \AA}$ $\beta = 121.801 (5)^\circ$ $V = 2043.3 (14) \text{ \AA}^3$ $Z = 4$ $F(000) = 920$ $D_x = 1.425 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 6562 reflections

 $\theta = 2.7\text{--}31.8^\circ$ $\mu = 0.11 \text{ mm}^{-1}$ $T = 100 \text{ K}$

Plate, colourless

 $0.28 \times 0.19 \times 0.06 \text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2005)

$T_{\min} = 0.971$, $T_{\max} = 0.994$

27961 measured reflections

3527 independent reflections

2710 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.065$

$\theta_{\text{max}} = 32.0^\circ$, $\theta_{\text{min}} = 2.2^\circ$

$h = -32 \rightarrow 32$

$k = -10 \rightarrow 10$

$l = -23 \rightarrow 23$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.059$

$wR(F^2) = 0.155$

$S = 1.08$

3527 reflections

189 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

All H-atom parameters refined

$w = 1/[\sigma^2(F_o^2) + (0.0752P)^2 + 1.2567P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} < 0.001$

$\Delta\rho_{\text{max}} = 0.50 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.21 \text{ e } \text{\AA}^{-3}$

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cyrosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.15443 (5)	0.04445 (13)	0.38705 (7)	0.0178 (2)
O2	0.05473 (6)	-0.06775 (15)	0.37300 (8)	0.0242 (2)
O3	0.30066 (5)	0.03278 (14)	0.48961 (7)	0.0212 (2)
O4	0.37511 (5)	0.26053 (15)	0.43926 (8)	0.0221 (2)
C1	0.08045 (7)	0.05014 (19)	0.34517 (10)	0.0175 (3)
C2	0.03949 (7)	0.19639 (19)	0.27048 (10)	0.0163 (3)
C3	0.07228 (7)	0.32194 (19)	0.24158 (10)	0.0168 (3)
C4	0.15015 (7)	0.31354 (18)	0.28870 (10)	0.0161 (3)
C5	0.19034 (8)	0.44106 (19)	0.26927 (10)	0.0185 (3)
C6	0.26498 (8)	0.42852 (19)	0.31768 (11)	0.0194 (3)
C7	0.30199 (7)	0.28719 (19)	0.38881 (10)	0.0174 (3)
C8	0.26401 (7)	0.15966 (18)	0.41265 (9)	0.0164 (3)
C9	0.18892 (7)	0.17416 (18)	0.36139 (10)	0.0154 (2)
C10	0.30773 (13)	-0.1493 (2)	0.45770 (15)	0.0362 (4)

C11	0.41539 (8)	0.3869 (2)	0.41549 (13)	0.0267 (3)
C12	0.02974 (8)	0.4697 (2)	0.16315 (12)	0.0247 (3)
H5	0.1654 (10)	0.540 (3)	0.2226 (15)	0.024 (5)*
H6	0.2908 (11)	0.516 (3)	0.3016 (16)	0.033 (5)*
H10A	0.3396 (14)	-0.139 (4)	0.429 (2)	0.063 (8)*
H10B	0.3324 (13)	-0.224 (3)	0.514 (2)	0.046 (6)*
H10C	0.2573 (16)	-0.202 (4)	0.410 (2)	0.067 (8)*
H11A	0.3985 (12)	0.382 (3)	0.3405 (19)	0.044 (6)*
H11B	0.4125 (11)	0.519 (3)	0.4338 (15)	0.028 (5)*
H11C	0.4642 (10)	0.340 (3)	0.4535 (14)	0.022 (4)*
H12A	0.0472 (11)	0.483 (3)	0.1149 (17)	0.035 (5)*
H12B	0.0359 (12)	0.591 (3)	0.1955 (18)	0.041 (6)*
H12C	-0.0217 (12)	0.440 (3)	0.1264 (17)	0.036 (5)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0162 (5)	0.0193 (5)	0.0163 (4)	0.0013 (3)	0.0074 (4)	0.0050 (4)
O2	0.0213 (5)	0.0253 (5)	0.0244 (5)	-0.0003 (4)	0.0111 (4)	0.0083 (4)
O3	0.0217 (5)	0.0219 (5)	0.0134 (4)	0.0047 (4)	0.0048 (4)	0.0029 (4)
O4	0.0144 (5)	0.0250 (5)	0.0230 (5)	-0.0015 (4)	0.0071 (4)	-0.0020 (4)
C1	0.0167 (6)	0.0199 (6)	0.0149 (6)	0.0003 (5)	0.0076 (5)	0.0008 (5)
C2	0.0155 (6)	0.0179 (6)	0.0140 (6)	0.0004 (4)	0.0067 (5)	-0.0003 (4)
C3	0.0166 (6)	0.0176 (6)	0.0145 (6)	0.0013 (4)	0.0071 (5)	0.0018 (4)
C4	0.0166 (6)	0.0174 (6)	0.0133 (5)	0.0013 (4)	0.0072 (5)	0.0009 (4)
C5	0.0199 (6)	0.0185 (6)	0.0165 (6)	0.0007 (5)	0.0091 (5)	0.0027 (5)
C6	0.0202 (6)	0.0200 (6)	0.0186 (6)	-0.0016 (5)	0.0106 (5)	-0.0002 (5)
C7	0.0142 (6)	0.0211 (6)	0.0143 (6)	-0.0007 (4)	0.0058 (5)	-0.0037 (5)
C8	0.0168 (6)	0.0176 (6)	0.0108 (5)	0.0018 (4)	0.0045 (5)	-0.0003 (4)
C9	0.0175 (6)	0.0155 (6)	0.0125 (5)	-0.0012 (4)	0.0075 (5)	-0.0006 (4)
C10	0.0564 (12)	0.0239 (8)	0.0302 (9)	0.0173 (8)	0.0241 (9)	0.0093 (7)
C11	0.0188 (7)	0.0267 (8)	0.0339 (8)	-0.0059 (6)	0.0134 (6)	-0.0047 (6)
C12	0.0183 (7)	0.0265 (7)	0.0254 (7)	0.0030 (5)	0.0089 (6)	0.0112 (6)

Geometric parameters (Å, °)

O1—C9	1.3750 (16)	C5—H5	0.952 (19)
O1—C1	1.3801 (17)	C6—C7	1.395 (2)
O2—C1	1.2085 (17)	C6—H6	0.96 (2)
O3—C8	1.3701 (16)	C7—C8	1.4025 (19)
O3—C10	1.428 (2)	C8—C9	1.3912 (19)
O4—C7	1.3640 (17)	C10—H10A	1.01 (3)
O4—C11	1.4334 (19)	C10—H10B	0.92 (3)
C1—C2	1.4618 (19)	C10—H10C	1.02 (3)
C2—C3	1.3592 (19)	C11—H11A	1.02 (2)
C2—C2 ⁱ	1.482 (3)	C11—H11B	1.00 (2)
C3—C4	1.4472 (19)	C11—H11C	0.963 (19)
C3—C12	1.5036 (19)	C12—H12A	1.01 (2)

C4—C5	1.3993 (19)	C12—H12B	0.97 (2)
C4—C9	1.4034 (18)	C12—H12C	0.97 (2)
C5—C6	1.384 (2)		
C9—O1—C1	121.36 (10)	O3—C8—C9	121.00 (12)
C8—O3—C10	114.77 (12)	O3—C8—C7	120.42 (12)
C7—O4—C11	116.63 (12)	C9—C8—C7	118.42 (12)
O2—C1—O1	116.99 (12)	O1—C9—C8	115.95 (11)
O2—C1—C2	125.28 (13)	O1—C9—C4	121.49 (12)
O1—C1—C2	117.72 (11)	C8—C9—C4	122.56 (12)
C3—C2—C1	121.77 (12)	O3—C10—H10A	108.3 (16)
C3—C2—C2 ⁱ	123.20 (11)	O3—C10—H10B	108.3 (15)
C1—C2—C2 ⁱ	115.03 (10)	H10A—C10—H10B	106 (2)
C2—C3—C4	118.80 (12)	O3—C10—H10C	108.6 (16)
C2—C3—C12	121.60 (13)	H10A—C10—H10C	115 (2)
C4—C3—C12	119.59 (12)	H10B—C10—H10C	110 (2)
C5—C4—C9	117.13 (12)	O4—C11—H11A	111.7 (13)
C5—C4—C3	124.03 (12)	O4—C11—H11B	112.6 (11)
C9—C4—C3	118.80 (12)	H11A—C11—H11B	108.8 (17)
C6—C5—C4	121.74 (13)	O4—C11—H11C	104.2 (11)
C6—C5—H5	119.7 (11)	H11A—C11—H11C	107.7 (16)
C4—C5—H5	118.6 (11)	H11B—C11—H11C	111.8 (16)
C5—C6—C7	119.83 (13)	C3—C12—H12A	111.0 (12)
C5—C6—H6	119.7 (13)	C3—C12—H12B	110.3 (14)
C7—C6—H6	120.5 (13)	H12A—C12—H12B	107.2 (17)
O4—C7—C6	124.46 (12)	C3—C12—H12C	110.3 (12)
O4—C7—C8	115.26 (12)	H12A—C12—H12C	110.5 (18)
C6—C7—C8	120.28 (13)	H12B—C12—H12C	107.4 (17)
C9—O1—C1—O2	-179.03 (12)	C5—C6—C7—O4	178.75 (12)
C9—O1—C1—C2	1.45 (18)	C5—C6—C7—C8	-1.0 (2)
O2—C1—C2—C3	-178.65 (14)	C10—O3—C8—C9	87.17 (17)
O1—C1—C2—C3	0.83 (19)	C10—O3—C8—C7	-97.47 (17)
O2—C1—C2—C2 ⁱ	1.3 (2)	O4—C7—C8—O3	6.82 (18)
O1—C1—C2—C2 ⁱ	-179.20 (11)	C6—C7—C8—O3	-173.39 (12)
C1—C2—C3—C4	-2.0 (2)	O4—C7—C8—C9	-177.70 (11)
C2 ⁱ —C2—C3—C4	178.01 (13)	C6—C7—C8—C9	2.09 (19)
C1—C2—C3—C12	178.72 (13)	C1—O1—C9—C8	176.73 (11)
C2 ⁱ —C2—C3—C12	-1.3 (2)	C1—O1—C9—C4	-2.48 (18)
C2—C3—C4—C5	-176.43 (13)	O3—C8—C9—O1	-5.18 (18)
C12—C3—C4—C5	2.8 (2)	C7—C8—C9—O1	179.37 (11)
C2—C3—C4—C9	1.01 (19)	O3—C8—C9—C4	174.02 (12)
C12—C3—C4—C9	-179.71 (13)	C7—C8—C9—C4	-1.42 (19)
C9—C4—C5—C6	1.4 (2)	C5—C4—C9—O1	178.84 (11)
C3—C4—C5—C6	178.94 (13)	C3—C4—C9—O1	1.22 (19)
C4—C5—C6—C7	-0.8 (2)	C5—C4—C9—C8	-0.32 (19)

C11—O4—C7—C6	-0.66 (19)	C3—C4—C9—C8	-177.94 (12)
C11—O4—C7—C8	179.12 (12)		

Symmetry code: (i) $-x, y, -z+1/2$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C6—H6...Cg1 ⁱⁱ	0.96 (2)	2.86 (2)	3.676 (2)	143.5 (18)

Symmetry code: (ii) $x, -y, z-1/2$.