

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2-(Dimethylaminomethyl)phenyl phenyl telluride

Tapash Chakraborty,<sup>a</sup> Harkesh B. Singh<sup>a\*</sup> and Ray J. Butcher<sup>b</sup><sup>a</sup>Department of Chemistry, Indian Institute of Technology Bombay, Powai 400 076, India, and <sup>b</sup>Department of Chemistry, Howard University, 525 College Street NW, Washington, DC 20059, USA

Correspondence e-mail: chhsbia@chem.iitb.ac.in


Received 19 September 2009; accepted 21 September 2009

Key indicators: single-crystal X-ray study;  $T = 110$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.023;  $wR$  factor = 0.058; data-to-parameter ratio = 31.0.

The title compound,  $\text{C}_{15}\text{H}_{17}\text{N}\text{Te}$ , is a heteroleptic Te,  $N$ -bidentate ligand having a short  $\text{Te}\cdots\text{N}$  contact [2.8079 (16) Å] involving a secondary bonding interaction between the amino N and  $\text{Te}^{\text{II}}$  atoms. The  $\text{Te}-\text{C}$  bond [2.136 (2) Å] *trans* to the amino group is slightly elongated compared to the other  $\text{Te}-\text{C}$  bond [2.1242 (18) Å] due to the hypervalent interaction. The bond angle for the *trans*  $\text{N}-\text{Te}-\text{C}$  atoms [164.92 (6)°] deviates significantly from linearity.

## Related literature

For Heck and cross-coupling reactions, see: Cella *et al.* (2006); Nishibayashi *et al.* (1996*a,b*); Zeni & Comasseto (1999); Zeni *et al.* (2001). For intramolecularly coordinated tellurides, see: Detty *et al.* (1995); Drake *et al.* (2001); Engman *et al.* (2004); Kaur *et al.* (1995, 2009); Menon *et al.* (1996); Panda *et al.* (1999); Singh *et al.* (1990). For van der Waals and covalent radii, see: Bondi (1964); Cordero *et al.* (2008).



## Experimental

## Crystal data

$\text{C}_{15}\text{H}_{17}\text{N}\text{Te}$   
 $M_r = 338.90$   
 Monoclinic,  $P2_1/c$   
 $a = 8.5736$  (3) Å  
 $b = 13.2472$  (5) Å  
 $c = 12.6719$  (4) Å  
 $\beta = 95.933$  (3)°

$V = 1431.52$  (9) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 2.06$  mm<sup>-1</sup>  
 $T = 110$  K  
 $0.49 \times 0.41 \times 0.27$  mm

## Data collection

Oxford Diffraction Gemini R CCD diffractometer  
 Absorption correction: multi-scan (CrysAlis Pro; Oxford Diffraction, 2009)  
 $T_{\min} = 0.728$ ,  $T_{\max} = 1.000$   
 20642 measured reflections  
 4836 independent reflections  
 2926 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$   
 $wR(F^2) = 0.058$   
 $S = 0.97$   
 4836 reflections  
 156 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.58$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.47$  e Å<sup>-3</sup>

Table 1

Selected geometric parameters (Å, °).

Te—C1	2.1242 (18)	Te—N	2.8079 (16)
Te—C10	2.136 (2)		
C1—Te—C10	94.19 (7)	C10—Te—N	164.92 (6)
C1—Te—N	70.77 (6)		

Data collection: *CrysAlis Pro* (Oxford Diffraction, 2009); cell refinement: *CrysAlis Pro*; data reduction: *CrysAlis Pro*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

HSB is grateful to the Department of Science and Technology (DST) for the award of a Ramanna Fellowship. TC is grateful to the CSIR for a JRF/SRF fellowship. RJB wishes to acknowledge the NSF-MRI program (grant CHE-0619278) for funds to purchase the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5068).

## References

- Bondi, A. (1964). *J. Phys. Chem.* **68**, 441–451.  
 Cella, R., Cunha, R. L. O., Reis, A. E. S., Pimenta, D. C., Klitzke, C. F. & Stefani, H. A. (2006). *J. Org. Chem.* **71**, 244–250.  
 Cordero, B., Gómez, V., Platero-Prats, A. E., Revés, M., Echeverría, J., Cremades, E., Barragán, F. & Alvarez, S. (2008). *Dalton Trans.* pp. 2832–2838.  
 Detty, R. M., Friedman, A. E. & McMillan, M. (1995). *Organometallics*, **14**, 1442–1449.  
 Drake, J. E., Hursthouse, M. B., Kulcsar, M., Light, M. E. & Silvestru, A. (2001). *J. Organomet. Chem.* **623**, 153–160.  
 Engman, L., Wojtoń, A., Oleksyn, B. J. & Śliwiński, J. (2004). *Phosphorus Sulfur Silicon Relat. Elem.* **179**, 285–292.  
 Kaur, R., Menon, S. C., Panda, A., Singh, H. B., Patel, R. P. & Butcher, R. J. (2009). *Organometallics*, **14**, 2363–2371.  
 Kaur, R., Singh, H. B. & Butcher, R. J. (1995). *Organometallics*, **14**, 4745–4763.  
 Menon, S. C., Singh, H. B., Jasinski, J. M., Jasinski, J. P. & Butcher, R. J. (1996). *Organometallics*, **15**, 1707–1712.  
 Nishibayashi, Y., Cho, C. S., Ohe, K. & Uemura, S. (1996*a*). *J. Organomet. Chem.* **526**, 335–339.  
 Nishibayashi, Y., Cho, C. S. & Uemura, S. (1996*b*). *J. Organomet. Chem.* **507**, 197–200.  
 Oxford Diffraction (2009). *CrysAlis Pro*. Oxford Diffraction Ltd, Yarnton, England.  
 Panda, A., Mughes, G., Singh, H. B. & Butcher, R. J. (1999). *Organometallics*, **18**, 1986–1993.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Singh, H. B., Sudha, N., West, A. A. & Hamor, T. A. (1990). *J. Chem. Soc. Dalton Trans.* pp. 907–913.  
Zeni, C. & Comasseto, J. V. (1999). *Tetrahedron Lett.* **40**, 4619–4622.

Zeni, G., Menezes, P. H., Moro, A. V., Braga, A. L., Silveira, C. C. & Stefani, H. A. (2001). *Synlett*, **9**, 1473–1475.

## supporting information

*Acta Cryst.* (2009). E65, o2562–o2563 [doi:10.1107/S1600536809038161]

## 2-(Dimethylaminomethyl)phenyl phenyl telluride

Tapash Chakraborty, Harkesh B. Singh and Ray J. Butcher

### S1. Comment

The structure of the title compound, (I), is shown below. Dimensions are available in the archived CIF.

Hydrid Te, N ligands with soft Te and hard N are of current interest due to their intramolecular coordination as the *trans* bond is activated and easily cleaved by metals (Kaur *et al.* 2009). Thus incorporation of soft tellurium and hard nitrogen into mixed donor Te, N bidentate ligands makes them interesting and promising as candidates for catalysts in combination with soft transition metals like Pd and the Rh group metals. Their coordination properties can be varied by changing the nitrogen function. Thus by introduction of a secondary bonding interaction which weakens the bond *trans* to Te in *ortho*-coordinated or suitably arranged substrates, the catalytic transformation of tellurides as substrates in Heck-type (Nishibayashi *et al.* 1996a; Nishibayashi *et al.* 1996b) and cross coupling (Zeni & Comasseto, 1999; Zeni *et al.* 2001; Cella *et al.* 2006) reactions and their coordination properties (Kaur *et al.* 2009) can be influenced. Our group (Singh *et al.* 1990; Kaur *et al.* 1995; Menon *et al.* 1996; Panda *et al.* 1999) as well as others (Detty *et al.* 1995; Drake *et al.* 2001, Engman *et al.* 2004) have been involved in the synthesis and studies of such intramolecularly coordinated organotellurides. The title compound, but not the structure, has been reported previously by Detty and co-workers as well as by our group (Kaur *et al.* 1995).


In the structure of the title compound, considering the bonding geometry around the Te as V-shaped with a longer intramolecular Te $\cdots$ N secondary interaction, a pseudo five-membered puckered ring can be envisioned. The Te $\cdots$ N distance of 2.8079 (16) Å is much greater than the sum of their covalent radii (2.11 Å; Cordero *et al.* 2008) but less than the sum of their van der Waal radii (3.61 Å; Bondi, 1964) and greater than the corresponding distance in similar compounds *viz.*, 2-NMe<sub>2</sub>CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>TeCl (2.362 (3) Å; Engman *et al.* 2004), 2-NMe<sub>2</sub>CH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>TeI (2.366 (4) Å; Kaur *et al.* 1995) or 8-(dimethylamino)-1-naphthyl phenyl telluride (2.713 (1) Å; Menon *et al.* 1996). Due to this hypervalent intramolecular Te $\cdots$ N contact the Te—C bond (2.137 (2) Å) *trans* to the amino group gets slightly elongated compared to the other Te—C bond (2.1249 (18) Å). The bond angle for the *trans* N—Te—C atoms (164.92 (6)°) deviates significantly from linearity.

### S2. Experimental

The title compound was prepared by the reported procedure (Kaur *et al.* 1995). A saturated solution of the compound was made in warm n-pentane and allowed to evaporate slowly at room temperature to grow crystals suitable for diffraction.


### S3. Refinement

H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms with C—H distances ranging from 0.95 to 0.99 Å and  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  [ $1.5U_{\text{eq}}(\text{C})$  for CH<sub>3</sub> H atoms].



**Figure 1**

The molecular structure of  $C_{15}H_{17}NTe$  showing the atom numbering scheme and 50% probability displacement ellipsoids. The secondary interaction between the N and Te is shown as a dashed line.



**Figure 2**

The molecular packing for  $C_{15}H_{17}NTe$  viewed down the  $a$  axis. The secondary interaction between the N and Te is shown by dashed lines.

### 2-(Dimethylaminomethyl)phenyl phenyl telluride

#### Crystal data

$C_{15}H_{17}NTe$

$M_r = 338.90$

Monoclinic,  $P2_1/c$

Hall symbol:  $-P\ 2ybc$

$a = 8.5736$  (3) Å

$b = 13.2472$  (5) Å

$c = 12.6719$  (4) Å

$\beta = 95.933$  (3)°

$V = 1431.52$  (9) Å<sup>3</sup>

$Z = 4$

$F(000) = 664$

$D_x = 1.572$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 8619 reflections

$\theta = 4.9$ – $32.4$ °

$\mu = 2.06$  mm<sup>-1</sup>

$T = 110$  K  $0.49 \times 0.41 \times 0.27$  mm  
 Prism, colorless

*Data collection*

Oxford Diffraction Gemini R CCD diffractometer	20642 measured reflections
Radiation source: Enhance (Mo) X-ray Source	4836 independent reflections
Graphite monochromator	2926 reflections with $I > 2\sigma(I)$
Detector resolution: $10.5081$ pixels $\text{mm}^{-1}$	$R_{\text{int}} = 0.028$
$\omega$ scans	$\theta_{\text{max}} = 32.4^\circ$ , $\theta_{\text{min}} = 4.9^\circ$
Absorption correction: multi-scan ( <i>CrysAlis PRO</i> ; Oxford Diffraction, 2009)	$h = -12 \rightarrow 12$
$T_{\text{min}} = 0.728$ , $T_{\text{max}} = 1.000$	$k = -18 \rightarrow 19$
	$l = -18 \rightarrow 18$

*Refinement*

Refinement on $F^2$	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.023$	H-atom parameters constrained
$wR(F^2) = 0.058$	$w = 1/[\sigma^2(F_o^2) + (0.0289P)^2]$
$S = 0.97$	where $P = (F_o^2 + 2F_c^2)/3$
4836 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
156 parameters	$\Delta\rho_{\text{max}} = 0.58 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.47 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

*Special details*

**Experimental.** *CrysAlis RED*, (Oxford Diffraction, 2009) Empirical absorption correction using spherical harmonics, implemented in *SCALE3 ABSPACK* scaling algorithm.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Te	0.886656 (14)	0.206826 (9)	0.327530 (10)	0.04605 (6)
C7	1.0873 (2)	-0.00329 (16)	0.33777 (15)	0.0462 (5)
H7A	1.1484	0.0230	0.4025	0.055*
H7B	1.1349	-0.0681	0.3190	0.055*
C1	0.8142 (2)	0.05794 (13)	0.36112 (13)	0.0373 (4)
C2	0.6624 (2)	0.03853 (16)	0.38466 (14)	0.0460 (4)
H2A	0.5888	0.0922	0.3840	0.055*
C3	0.6175 (2)	-0.05790 (16)	0.40897 (16)	0.0533 (5)
H3A	0.5134	-0.0704	0.4249	0.064*
C4	0.7225 (3)	-0.13543 (17)	0.41016 (16)	0.0566 (5)
H4A	0.6923	-0.2016	0.4287	0.068*
C5	0.8725 (3)	-0.11769 (15)	0.38443 (15)	0.0497 (5)

H5A	0.9435	-0.1726	0.3831	0.060*
C6	0.9219 (2)	-0.02150 (14)	0.36041 (14)	0.0395 (4)
N	1.09600 (18)	0.06793 (12)	0.25179 (13)	0.0470 (4)
C8	1.0372 (3)	0.0248 (2)	0.14988 (17)	0.0711 (7)
H8A	0.9302	-0.0002	0.1533	0.107*
H8B	1.1050	-0.0311	0.1329	0.107*
H8C	1.0366	0.0767	0.0947	0.107*
C9	1.2542 (3)	0.1090 (2)	0.2514 (2)	0.0739 (7)
H9A	1.2852	0.1426	0.3192	0.111*
H9B	1.2559	0.1579	0.1934	0.111*
H9C	1.3277	0.0540	0.2412	0.111*
C10	0.6957 (2)	0.27717 (13)	0.39468 (15)	0.0433 (4)
C11	0.5679 (3)	0.31290 (16)	0.32958 (17)	0.0554 (5)
H11A	0.5633	0.3032	0.2550	0.066*
C12	0.4467 (3)	0.36259 (18)	0.37197 (19)	0.0631 (6)
H12A	0.3587	0.3856	0.3266	0.076*
C13	0.4534 (3)	0.37856 (17)	0.47833 (19)	0.0609 (6)
H13A	0.3709	0.4136	0.5071	0.073*
C14	0.5789 (3)	0.34409 (18)	0.54402 (18)	0.0627 (6)
H14A	0.5827	0.3548	0.6184	0.075*
C15	0.6989 (3)	0.29428 (15)	0.50326 (16)	0.0536 (5)
H15B	0.7856	0.2710	0.5497	0.064*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Te	0.04845 (9)	0.04144 (8)	0.04904 (9)	0.00098 (6)	0.00871 (6)	0.00703 (6)
C7	0.0416 (10)	0.0500 (12)	0.0469 (11)	0.0059 (9)	0.0049 (8)	0.0024 (8)
C1	0.0395 (9)	0.0410 (10)	0.0314 (8)	-0.0007 (8)	0.0031 (7)	-0.0031 (7)
C2	0.0416 (10)	0.0501 (12)	0.0468 (10)	0.0001 (9)	0.0069 (8)	-0.0066 (9)
C3	0.0495 (11)	0.0563 (14)	0.0564 (12)	-0.0160 (10)	0.0169 (9)	-0.0145 (10)
C4	0.0731 (15)	0.0465 (13)	0.0521 (12)	-0.0174 (11)	0.0154 (10)	-0.0081 (9)
C5	0.0640 (13)	0.0393 (11)	0.0463 (11)	0.0035 (10)	0.0080 (9)	-0.0023 (9)
C6	0.0424 (10)	0.0416 (10)	0.0341 (9)	0.0011 (8)	0.0025 (7)	0.0013 (8)
N	0.0431 (9)	0.0491 (10)	0.0503 (9)	0.0027 (7)	0.0128 (7)	0.0049 (8)
C8	0.0833 (16)	0.0851 (19)	0.0466 (13)	0.0158 (15)	0.0149 (11)	0.0025 (12)
C9	0.0555 (13)	0.0756 (18)	0.0948 (18)	-0.0037 (12)	0.0279 (12)	0.0159 (14)
C10	0.0504 (11)	0.0327 (10)	0.0469 (11)	-0.0005 (8)	0.0049 (9)	0.0028 (8)
C11	0.0605 (13)	0.0542 (14)	0.0499 (12)	0.0097 (10)	-0.0022 (10)	-0.0012 (9)
C12	0.0563 (13)	0.0605 (15)	0.0712 (15)	0.0165 (11)	0.0005 (11)	0.0013 (12)
C13	0.0652 (14)	0.0455 (12)	0.0755 (15)	0.0070 (11)	0.0238 (12)	0.0006 (11)
C14	0.0827 (16)	0.0591 (15)	0.0486 (12)	0.0050 (12)	0.0171 (12)	-0.0042 (10)
C15	0.0603 (13)	0.0534 (13)	0.0462 (11)	0.0050 (10)	0.0006 (9)	0.0030 (9)

*Geometric parameters (Å, °)*

Te—C1	2.1242 (18)	N—C9	1.462 (3)
Te—C10	2.136 (2)	C8—H8A	0.9800

Te—N	2.8079 (16)	C8—H8B	0.9800
C7—N	1.449 (2)	C8—H8C	0.9800
C7—C6	1.495 (2)	C9—H9A	0.9800
C7—H7A	0.9900	C9—H9B	0.9800
C7—H7B	0.9900	C9—H9C	0.9800
C1—C2	1.389 (3)	C10—C11	1.385 (3)
C1—C6	1.401 (3)	C10—C15	1.392 (3)
C2—C3	1.378 (3)	C11—C12	1.384 (3)
C2—H2A	0.9500	C11—H11A	0.9500
C3—C4	1.365 (3)	C12—C13	1.360 (3)
C3—H3A	0.9500	C12—H12A	0.9500
C4—C5	1.379 (3)	C13—C14	1.369 (3)
C4—H4A	0.9500	C13—H13A	0.9500
C5—C6	1.387 (3)	C14—C15	1.368 (3)
C5—H5A	0.9500	C14—H14A	0.9500
N—C8	1.454 (3)	C15—H15B	0.9500
C1—Te—C10	94.19 (7)	C9—N—Te	112.55 (14)
C1—Te—N	70.77 (6)	N—C8—H8A	109.5
C10—Te—N	164.92 (6)	N—C8—H8B	109.5
N—C7—C6	111.92 (14)	H8A—C8—H8B	109.5
N—C7—H7A	109.2	N—C8—H8C	109.5
C6—C7—H7A	109.2	H8A—C8—H8C	109.5
N—C7—H7B	109.2	H8B—C8—H8C	109.5
C6—C7—H7B	109.2	N—C9—H9A	109.5
H7A—C7—H7B	107.9	N—C9—H9B	109.5
C2—C1—C6	119.70 (17)	H9A—C9—H9B	109.5
C2—C1—Te	120.99 (14)	N—C9—H9C	109.5
C6—C1—Te	119.30 (13)	H9A—C9—H9C	109.5
C3—C2—C1	120.59 (18)	H9B—C9—H9C	109.5
C3—C2—H2A	119.7	C11—C10—C15	117.82 (19)
C1—C2—H2A	119.7	C11—C10—Te	120.25 (15)
C4—C3—C2	120.08 (19)	C15—C10—Te	121.81 (14)
C4—C3—H3A	120.0	C12—C11—C10	120.7 (2)
C2—C3—H3A	120.0	C12—C11—H11A	119.6
C3—C4—C5	119.9 (2)	C10—C11—H11A	119.6
C3—C4—H4A	120.0	C13—C12—C11	120.2 (2)
C5—C4—H4A	120.0	C13—C12—H12A	119.9
C4—C5—C6	121.48 (19)	C11—C12—H12A	119.9
C4—C5—H5A	119.3	C12—C13—C14	120.0 (2)
C6—C5—H5A	119.3	C12—C13—H13A	120.0
C5—C6—C1	118.17 (17)	C14—C13—H13A	120.0
C5—C6—C7	120.53 (17)	C15—C14—C13	120.4 (2)
C1—C6—C7	121.27 (17)	C15—C14—H14A	119.8
C7—N—C8	111.84 (17)	C13—C14—H14A	119.8
C7—N—C9	111.41 (16)	C14—C15—C10	120.87 (19)
C8—N—C9	112.33 (17)	C14—C15—H15B	119.6
C7—N—Te	94.88 (10)	C10—C15—H15B	119.6



---

C8—N—Te	112.69 (13)		
C10—Te—C1—C2	17.26 (15)	C6—C7—N—Te	45.84 (15)
N—Te—C1—C2	-161.59 (15)	C1—Te—N—C7	-35.48 (10)
C10—Te—C1—C6	-162.12 (13)	C10—Te—N—C7	-39.9 (3)
N—Te—C1—C6	19.02 (12)	C1—Te—N—C8	80.65 (15)
C6—C1—C2—C3	1.0 (3)	C10—Te—N—C8	76.3 (3)
Te—C1—C2—C3	-178.36 (13)	C1—Te—N—C9	-151.08 (15)
C1—C2—C3—C4	0.0 (3)	C10—Te—N—C9	-155.5 (2)
C2—C3—C4—C5	-1.6 (3)	C1—Te—C10—C11	-101.37 (16)
C3—C4—C5—C6	2.2 (3)	N—Te—C10—C11	-97.2 (3)
C4—C5—C6—C1	-1.1 (3)	C1—Te—C10—C15	82.60 (16)
C4—C5—C6—C7	177.09 (17)	N—Te—C10—C15	86.8 (3)
C2—C1—C6—C5	-0.5 (3)	C15—C10—C11—C12	-0.9 (3)
Te—C1—C6—C5	178.88 (13)	Te—C10—C11—C12	-177.11 (17)
C2—C1—C6—C7	-178.66 (16)	C10—C11—C12—C13	1.2 (4)
Te—C1—C6—C7	0.7 (2)	C11—C12—C13—C14	-1.0 (4)
N—C7—C6—C5	140.26 (18)	C12—C13—C14—C15	0.6 (4)
N—C7—C6—C1	-41.6 (2)	C13—C14—C15—C10	-0.3 (4)
C6—C7—N—C8	-71.0 (2)	C11—C10—C15—C14	0.5 (3)
C6—C7—N—C9	162.38 (17)	Te—C10—C15—C14	176.60 (17)

---