

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Methyl-2H-1,3-oxazine-2,6(3H)-dione

Damon Parrish, Fredrick Leuschner, Gretchen M. Rehberg and Margaret E. Kastner*

Department of Chemistry, Bucknell University, Lewisburg, PA 17837, USA

Correspondence e-mail: kastner@bucknell.edu


Received 13 August 2009; accepted 28 August 2009

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.049; wR factor = 0.191; data-to-parameter ratio = 15.6.

In the title compound, $\text{C}_5\text{H}_5\text{NO}_3$, the planar (maximum deviation = 0.075 Å for the ring O atom) molecules form $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds in a zigzag chain ($\text{C}-\text{O}\cdots\text{N}$ bond angle $\simeq 140^\circ$) between glide-related molecules.

Related literature

For synthetic background, see: Warren *et al.* (1975); Rehberg & Glass (1995). For related structures, see: Copley *et al.* (2005); Parrish, Leuschner *et al.* (2009); Parrish, Tivitmahaisoon *et al.* (2009).



Experimental

Crystal data

$\text{C}_5\text{H}_5\text{NO}_3$
 $M_r = 127.1$
 Monoclinic, $P2_1/n$

$a = 7.254$ (3) Å
 $b = 6.683$ (2) Å
 $c = 11.689$ (5) Å

$\beta = 98.11$ (4) $^\circ$
 $V = 561.0$ (4) Å³
 $Z = 4$
 Mo $K\alpha$ radiation

$\mu = 0.13$ mm⁻¹
 $T = 293$ K
 $0.46 \times 0.30 \times 0.10$ mm

Data collection

Bruker R3/V diffractometer
 Absorption correction: none
 1410 measured reflections
 1294 independent reflections
 910 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.012$
 3 standard reflections
 every 97 reflections
 intensity decay: none

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.191$
 $S = 0.93$
 1294 reflections

83 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.23$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N3}-\text{H3}\cdots\text{O6}^i$	0.86	2.02	2.877 (3)	173

Symmetry code: (i) $x - \frac{1}{2}, -y + \frac{1}{2}, z - \frac{1}{2}$.

Data collection: *XSCANS* (Bruker, 1996); cell refinement: *XSCANS* (Bruker, 1996); data reduction: *XSCANS* (Bruker, 1996); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).

The authors thank the National Science Foundation for grant No. ILI8951058.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2197).

References

- Bruker (1996). *XSCANS*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Copley, R. C. B., Deprez, L. S., Lewis, T. C. & Price, S. L. (2005). *CrystEngComm*, **7**, 421–428.
 Parrish, D., Leuschner, F., Rehberg, G. M. & Kastner, M. E. (2009). *Acta Cryst.* **E65**, o2356.
 Parrish, D., Tivitmahaisoon, P., Rehberg, G. M. & Kastner, M. E. (2009). *Acta Cryst.* **E65**, o2355.
 Rehberg, G. M. & Glass, B. M. (1995). *Org. Prep. Proced. Int.* **27**, 651–652.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Warren, J. D., MacMillan, J. H. & Washburne, S. S. (1975). *J. Org. Chem.* **40**, 743–746.

supporting information

Acta Cryst. (2009). E65, o2354 [doi:10.1107/S1600536809034618]

4-Methyl-2H-1,3-oxazine-2,6(3H)-dione

Damon Parrish, Fredrick Leuschner, Gretchen M. Rehberg and Margaret E. Kastner

S1. Comment

The synthesis of derivatives of 3-oxauracil has previously been reported (Warren *et al.*, 1975) and an improved synthesis of the unsubstituted 3-oxauracil was reported by Rehberg & Glass (1995). The structure of the unsubstituted 3-oxauracil and its monohydrate have been reported (Copley *et al.*, 2005). Three derivatives of 3-oxauracil (4-methyl, 4-bromo, and 4,5-dichloro) have been prepared in our laboratory in route to the synthesis of 1-aza-1,3-butadienes. In this paper, we report the crystal structure of the title compound, (I).

In the title compound (Fig. 1) only one intermolecular H-bond is formed between N3 and O6 of glide-related molecules (details are given in Table 1). Although the molecules of (I) are planar, the H-bonding chains are staggered as shown in Figure 2. The hydrogen bonding networks in (I) differs significantly from the hydrogen bonding in 4,5-dichloro (Parrish, Leuschner *et al.*, 2009) and 4-bromo (Parrish, Tivitmahaisoon *et al.*, 2009) derivatives.

S2. Experimental

Citraconic anhydride (3-methylfuran-2,5-dione, 2.0 ml, 22 mmol) and trimethylsilyl azide (3.0 ml, 23 mmol) were added to 10 ml dichloromethane at 273 K and stirred under nitrogen for 4 h. Upon warming to room temperature over night, a white precipitate formed. Ethanol (2.5 ml) was added, the mixture stirred 2 additional hours, and then the solvent was removed under reduced pressure to obtain the title compound; yield: 1.7 g (13 mmol, 59%). Crystals of the title compound were grown from a solution of acetone at room temperature by slow evaporation.

S3. Refinement

Hydrogen positions were calculated and refined using a riding model using the following C—H distances: methyl 0.96 Å, methylene 0.93 Å, and N—H 0.88 Å with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C5/N3})$ and $1.5U_{\text{eq}}(\text{C7})$.





Figure 1

The molecular structure of the title compound, with atom labels and 50% probability displacement ellipsoids for non-H atoms.

**Figure 2**

The packing of the title compound viewed down the a axis; intermolecular hydrogen bonds have been represented by dashed lines.

4-Methyl-2H-1,3-oxazine-2,6(3H)-dione

Crystal data

$C_5H_5NO_3$

$M_r = 127.1$

Monoclinic, $P2_1/n$

Hall symbol: $-P\ 2_1n$

$a = 7.254\ (3)\ \text{\AA}$

$b = 6.683\ (2)\ \text{\AA}$

$c = 11.689\ (5)\ \text{\AA}$

$\beta = 98.11\ (4)^\circ$

$V = 561.0\ (4)\ \text{\AA}^3$

$Z = 4$

$F(000) = 264$

Data collection

Bruker R3/V

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\theta - 2\theta$ scans

1410 measured reflections

1294 independent reflections

910 reflections with $I > 2\sigma(I)$

$D_x = 1.505\ \text{Mg m}^{-3}$

$D_m = 1.46\ \text{Mg m}^{-3}$

D_m measured by floatation in
bromoform/hexane solution

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 20 reflections

$\theta = 10\text{--}12.5^\circ$

$\mu = 0.13\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Plates, colorless

$0.46 \times 0.30 \times 0.10\ \text{mm}$

$R_{\text{int}} = 0.012$

$\theta_{\text{max}} = 27.6^\circ$, $\theta_{\text{min}} = 3.1^\circ$

$h = 0 \rightarrow 9$

$k = 0 \rightarrow 8$

$l = -15 \rightarrow 15$

3 standard reflections every 97 reflections

intensity decay: none

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.049$
 $wR(F^2) = 0.191$
 $S = 0.93$
 1294 reflections
 83 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1301P)^2 + 0.1905P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.005$
 $\Delta\rho_{\max} = 0.23 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.24 \text{ e } \text{Å}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.6042 (2)	0.1383 (2)	0.68042 (13)	0.0466 (5)
C2	0.4998 (3)	0.2078 (3)	0.58306 (19)	0.0422 (5)
O2	0.3337 (2)	0.2161 (3)	0.57888 (19)	0.0694 (6)
N3	0.5939 (2)	0.2571 (3)	0.49490 (14)	0.0406 (5)
H3	0.5305	0.2967	0.4312	0.049*
C4	0.7833 (3)	0.2477 (3)	0.50089 (18)	0.0392 (5)
C5	0.8847 (3)	0.1896 (3)	0.59971 (19)	0.0430 (5)
H5	1.0137	0.1846	0.6048	0.052*
C6	0.7988 (3)	0.1357 (3)	0.69668 (18)	0.0435 (5)
O6	0.8682 (3)	0.0868 (3)	0.79242 (15)	0.0695 (7)
C7	0.8619 (4)	0.3002 (5)	0.3938 (2)	0.0630 (8)
H7A	0.9955	0.2979	0.4093	0.095*
H7B	0.8210	0.4317	0.3687	0.095*
H7C	0.8201	0.2050	0.3343	0.095*

Atomic displacement parameters (Å^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0448 (9)	0.0555 (10)	0.0417 (8)	-0.0028 (7)	0.0137 (6)	0.0020 (7)
C2	0.0329 (10)	0.0455 (12)	0.0490 (11)	0.0003 (8)	0.0086 (8)	-0.0068 (9)
O2	0.0324 (9)	0.0847 (14)	0.0936 (15)	-0.0020 (9)	0.0174 (9)	-0.0026 (11)
N3	0.0342 (9)	0.0511 (10)	0.0350 (9)	0.0041 (7)	0.0003 (7)	0.0010 (7)
C4	0.0374 (10)	0.0384 (10)	0.0445 (11)	0.0029 (8)	0.0152 (8)	-0.0025 (9)
C5	0.0294 (9)	0.0458 (12)	0.0534 (12)	0.0012 (9)	0.0049 (8)	-0.0021 (10)
C6	0.0447 (11)	0.0395 (11)	0.0435 (11)	-0.0024 (9)	-0.0038 (9)	-0.0025 (9)

O6	0.0831 (14)	0.0687 (13)	0.0494 (10)	-0.0096 (10)	-0.0167 (9)	0.0110 (9)
C7	0.0651 (16)	0.0733 (18)	0.0573 (14)	0.0053 (13)	0.0313 (12)	0.0133 (13)

Geometric parameters (Å, °)

O1—C2	1.357 (3)	C4—C7	1.489 (3)
O1—C6	1.398 (3)	C5—C6	1.415 (3)
C2—O2	1.200 (3)	C5—H5	0.9300
C2—N3	1.354 (3)	C6—O6	1.206 (3)
N3—C4	1.368 (3)	C7—H7A	0.9600
N3—H3	0.8600	C7—H7B	0.9600
C4—C5	1.337 (3)	C7—H7C	0.9600
C2—O1—C6	123.50 (17)	C4—C5—H5	119.5
O2—C2—N3	124.6 (2)	C6—C5—H5	119.5
O2—C2—O1	119.2 (2)	O6—C6—O1	114.3 (2)
N3—C2—O1	116.10 (18)	O6—C6—C5	129.7 (2)
C2—N3—C4	124.08 (18)	O1—C6—C5	115.99 (18)
C2—N3—H3	118.0	C4—C7—H7A	109.5
C4—N3—H3	118.0	C4—C7—H7B	109.5
C5—C4—N3	118.95 (18)	H7A—C7—H7B	109.5
C5—C4—C7	124.5 (2)	C4—C7—H7C	109.5
N3—C4—C7	116.6 (2)	H7A—C7—H7C	109.5
C4—C5—C6	121.03 (19)	H7B—C7—H7C	109.5
C6—O1—C2—O2	-175.5 (2)	N3—C4—C5—C6	0.9 (3)
C6—O1—C2—N3	6.7 (3)	C7—C4—C5—C6	-177.8 (2)
O2—C2—N3—C4	179.7 (2)	C2—O1—C6—O6	173.2 (2)
O1—C2—N3—C4	-2.6 (3)	C2—O1—C6—C5	-6.9 (3)
C2—N3—C4—C5	-1.1 (3)	C4—C5—C6—O6	-177.3 (2)
C2—N3—C4—C7	177.7 (2)	C4—C5—C6—O1	2.8 (3)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N3—H3 \cdots O6 ⁱ	0.86	2.02	2.877 (3)	173

Symmetry code: (i) $x-1/2, -y+1/2, z-1/2$.