

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

# 1,3-Bis(hydroxymethyl)benzimidazolin-2-one

H. C. Devarajgowda,<sup>a\*</sup> V. Madhura,<sup>b</sup>  
 B. S. Palakshamurthy,<sup>a</sup> S. Jeyaseelan<sup>a</sup> and Manohar V.  
 Kulkarni<sup>b</sup>

<sup>a</sup>Department of Physics, Yuvaraja's College (Constituent College), University of Mysore, Mysore 570 005, Karnataka, India, and <sup>b</sup>Department of Chemistry, Karnataka University, Dharwad 580 003, Karnataka, India  
 Correspondence e-mail: devarajgowda@yahoo.com


Received 24 September 2009; accepted 7 October 2009

Key indicators: single-crystal X-ray study;  $T = 273$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  
 $R$  factor = 0.040;  $wR$  factor = 0.110; data-to-parameter ratio = 14.1.

The title compound,  $\text{C}_9\text{H}_{10}\text{N}_2\text{O}_3$ , crystallizes with one and a half molecules in the asymmetric unit, one lying on a general position and the other on a twofold rotation axis. The dihedral angle between the two independent benzimidazole ring systems is  $18.96(5)^\circ$ . In the crystal, molecules are linked into a three-dimensional network by  $\text{O}-\text{H}\cdots\text{O}$  hydrogen bonding involving  $N$ -hydroxymethyl and carbonyl groups, and  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds.

## Related literature

For general background to 2-benzimidazolones, see: Raghu *et al.* (2005); Porret & Hebermeier (1974); Habermeier (1976); Trask-Morrel *et al.* (1988); Hammach *et al.* (2006); Bansal *et al.* (1981). For related structures, see: Anklekar & Kulkarni (1995); Schwiebert *et al.* (1996). For the synthesis, see: Zinner & Spangenberg (1958).



## Experimental

### Crystal data

$\text{C}_9\text{H}_{10}\text{N}_2\text{O}_3$   
 $M_r = 194.19$   
 Monoclinic,  $C2/c$   
 $a = 13.5515(14)$  Å  
 $b = 11.0848(12)$  Å  
 $c = 17.6253(19)$  Å  
 $\beta = 94.216(2)^\circ$   
 $V = 2640.4(5)$  Å<sup>3</sup>  
 $Z = 12$   
 Mo  $K\alpha$  radiation

$\mu = 0.11$  mm<sup>-1</sup>  
 $T = 273$  K  
 $0.22 \times 0.20 \times 0.10$  mm

### Data collection

Bruker SMART CCD area-detector  
 diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Sheldrick, 2004)  
 $T_{\min} = 0.976$ ,  $T_{\max} = 0.986$   
 13521 measured reflections  
 2684 independent reflections  
 2395 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.025$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$   
 $wR(F^2) = 0.110$   
 $S = 1.10$   
 2684 reflections  
 191 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.16$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.24$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1A}\cdots\text{O4}$	0.82	1.99	2.8003 (16)	169
$\text{O2}-\text{H2A}\cdots\text{O5}^{\text{ii}}$	0.82	1.93	2.7503 (18)	176
$\text{O5}-\text{H5A}\cdots\text{O3}^{\text{ii}}$	0.82	1.84	2.6551 (17)	175
$\text{C3}-\text{H3}\cdots\text{O2}^{\text{iii}}$	0.93	2.58	3.489 (2)	164
$\text{C14}-\text{H14B}\cdots\text{O1}^{\text{ii}}$	0.97	2.54	3.364 (2)	143

Symmetry codes: (i)  $x, -y + 1, z - \frac{1}{2}$ ; (ii)  $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$ ; (iii)  $-x + \frac{3}{2}, -y + \frac{3}{2}, -z$ .

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997); software used to prepare material for publication: SHELXL97.

The authors thank Professor T. N. Guru Row and Miss Brinda Selvaraj, Solid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore, for their help with the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2925).

## References

- Anklekar, K. Y. & Kulkarni, M. V. (1995). *Indian J. Chem. Sect. B*, **34**, 677–678.  
 Bansal, P. C., Ian, H., Pitman, I. H. & Higuchi, T. (1981). *J. Pharm. Sci.* **70**, 855–857.  
 Bruker (2001). SMART and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
 Habermeier, J. (1976). *Ger. Offen.* 2 453 450.  
 Hammach, A., Barbosa, A., Gaenzler, F. C., Fadra, T., Goldberg, D., Hao, M.-H., Kroe, R. R., Liu, P., Qian, K. C., Ralph, M., Sarko, C., Soleymanzadeh, F. & Moss, N. (2006). *Bioorg. Med. Chem. Lett.* **16**, 6316–6320.  
 Porret, D. & Hebermeier, J. (1974). *Ger. Offen.* 2 342 432.  
 Raghu, A. V., Gadaginamath, G. S. & Amminabhavi, T. M. (2005). *J. Appl. Polym. Sci.* **98**, 2236–2244.  
 Schwiebert, K. E., Chin, D. N., MacDonald, J. C. & Whitesides, G. M. (1996). *J. Am. Chem. Soc.* **118**, 4018–4029.  
 Sheldrick, G. M. (2004). SADABS. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Trask-Morrel, B. J., Franklin, W. E. & Liu, R. H. (1988). *Text. Chem. Color*, pp. 20–21.  
 Zinner, H. & Spangenberg, B. (1958). *Chem. Ber.* **91**, 1432–1437.

## supporting information

*Acta Cryst.* (2009). E65, o2735 [https://doi.org/10.1107/S1600536809040963]

**1,3-Bis(hydroxymethyl)benzimidazolin-2-one**

**H. C. Devarajegowda, V. Madhura, B. S. Palakshamurthy, S. Jeyaseelan and Manohar V. Kulkarni**

**S1. Comment**

1,3-Bishydroxyalkylated benzimidazolones are an important class of functionalized benzimidazoles which have been found to be useful as polymer intermediates (Raghu *et al.*, 2005; Porret *et al.*, 1974), fire retardants (Habermeier, 1976), and in curing process in textile industry (Trask-Morrel *et al.*, 1988). Solid state chemistry of hydroxy methylated benzimidazole derivatives leading to thermal extrusion of formaldehyde has been reported (Anklekar *et al.*, 1995) by our group. Design and synthesis of benzimidazolone p38 MAP kinase inhibitors (Hammach *et al.*, 2006) is based on the analysis of their crystal structure data. Benzimidazolones have been reported to crystallize as hydrogen bonded molecular tapes (Schwiebert *et al.*, 1996) which has been used to engineer structures of organic solids. In view of the therapeutic importance of aromatic and hetero aromatic compounds (Bansal *et al.*, 1981) containing *N*-hydroxymethyl group, we report here the crystal structure of title compound.

The asymmetric unit of the title compound contains one and a half molecules, one lying on a general position and the other on a twofold rotation axis (Fig.1). Atoms O4 and C13 lie on the twofold rotation axis. The two independent benzimidazole ring systems form a dihedral angle of 18.96 (5)°. There are no intramolecular hydrogen bonding between *N*-hydroxymethyl and carbonyl groups.


In the crystal, O—H...O hydrogen bonding involving *N*-hydroxymethyl and carbonyl groups results in the formation of three-dimensional network (Fig.2). In addition, C—H...O hydrogen bonds (Table 1) are observed. This type of intermolecular association is similar to that observed in the structure of benzimidazolone (Schwiebert *et al.*, 1996).

**S2. Experimental**


The title compound was prepared by following a literature method (Zinner *et al.*, 1958). A mixture of 2-hydroxy benzimidazole (13.4 g, 0.01 *M*) and 37% formalin (30 ml, 1*M*) was refluxed for 30 minutes in presence of 100 ml water. The solid product formed was filtered and single crystals were grown by slow evaporation in water (yield 92%, m.p. 433 K). Spectral data IR<sub>ν<sub>CO</sub></sub> = 1700 cm<sup>-1</sup>, ν<sub>OH</sub> = 3300 cm<sup>-1</sup>. <sup>1</sup>H NMR -(CDCl<sub>3</sub>+DMSO-*d*<sub>6</sub>)δ p.p.m. - 5.3(4*H*, d, CH<sub>2</sub>) appeared as singlet on D<sub>2</sub>O exchange, 6.2(2*H*, t, OH) vanished on D<sub>2</sub>O exchange, 7.4–7.8 (4*H*, m, Ar-H). Mass *m/z* = 134 (100%).

**S3. Refinement**

H atoms were positioned at calculated positions [O-H = 0.82 Å and C-H = 0.93–0.97 Å] and refined using a riding model with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  and  $1.5U_{\text{eq}}(\text{O})$ .

**Figure 1**

The asymmetric unit of the title compound. Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

Packing of the molecules viewed down the *b* axis. Hydrogen bonds are shown as dashed lines.

## 1,3-Bis(hydroxymethyl)benzimidazolin-2-one

## Crystal data

C<sub>9</sub>H<sub>10</sub>N<sub>2</sub>O<sub>3</sub> $M_r = 194.19$ Monoclinic,  $C2/c$ Hall symbol:  $-C\ 2yc$  $a = 13.5515\ (14)\ \text{\AA}$  $b = 11.0848\ (12)\ \text{\AA}$  $c = 17.6253\ (19)\ \text{\AA}$  $\beta = 94.216\ (2)^\circ$  $V = 2640.4\ (5)\ \text{\AA}^3$  $Z = 12$  $F(000) = 1224$  $D_x = 1.465\ \text{Mg m}^{-3}$ 

Melting point: 433 K

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$ 

Cell parameters from 2684 reflections

 $\theta = 2.3\text{--}26.4^\circ$  $\mu = 0.11\ \text{mm}^{-1}$  $T = 273\ \text{K}$ 

Plate, white

 $0.22 \times 0.20 \times 0.10\ \text{mm}$ 

## Data collection

Bruker SMART CCD area-detector  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\omega$  and  $\varphi$  scansAbsorption correction: multi-scan  
(*SADABS*; Sheldrick, 2004) $T_{\min} = 0.976$ ,  $T_{\max} = 0.986$ 

13521 measured reflections

2684 independent reflections

2395 reflections with  $I > 2\sigma(I)$  $R_{\text{int}} = 0.025$  $\theta_{\max} = 26.4^\circ$ ,  $\theta_{\min} = 2.3^\circ$  $h = -16 \rightarrow 16$  $k = -13 \rightarrow 13$  $l = -22 \rightarrow 22$ 

## Refinement

Refinement on  $F^2$ 

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.040$  $wR(F^2) = 0.110$  $S = 1.10$ 

2684 reflections

191 parameters

0 restraints

Primary atom site location: structure-invariant  
direct methodsSecondary atom site location: difference Fourier  
mapHydrogen site location: inferred from  
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0486P)^2 + 1.5821P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{\max} = 0.001$  $\Delta\rho_{\max} = 0.16\ \text{e \AA}^{-3}$  $\Delta\rho_{\min} = -0.24\ \text{e \AA}^{-3}$ 

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.36019 (9)	0.44213 (11)	0.19000 (7)	0.0555 (3)
H1A	0.4009	0.3984	0.2131	0.083*
O2	0.51268 (9)	0.85515 (11)	-0.04107 (7)	0.0560 (3)
H2A	0.4597	0.8565	-0.0667	0.084*

O3	0.32846 (8)	0.69395 (11)	0.07135 (7)	0.0525 (3)
N1	0.44553 (9)	0.54147 (11)	0.09507 (7)	0.0390 (3)
N2	0.49663 (9)	0.72131 (11)	0.06132 (7)	0.0420 (3)
C1	0.61458 (12)	0.44272 (15)	0.11034 (9)	0.0469 (4)
H1	0.5932	0.3665	0.1238	0.056*
C2	0.71420 (13)	0.46756 (18)	0.10526 (10)	0.0568 (5)
H2	0.7604	0.4063	0.1146	0.068*
C3	0.74639 (13)	0.5819 (2)	0.08647 (10)	0.0582 (5)
H3	0.8138	0.5963	0.0847	0.070*
C4	0.67978 (12)	0.67501 (17)	0.07028 (9)	0.0499 (4)
H4	0.7012	0.7516	0.0576	0.060*
C5	0.58057 (11)	0.64950 (13)	0.07380 (8)	0.0388 (3)
C6	0.54853 (11)	0.53545 (13)	0.09463 (8)	0.0372 (3)
C7	0.41401 (11)	0.65604 (14)	0.07545 (8)	0.0398 (3)
C8	0.49544 (14)	0.84582 (14)	0.03582 (10)	0.0511 (4)
H8A	0.4317	0.8813	0.0440	0.061*
H8B	0.5458	0.8909	0.0658	0.061*
C9	0.37880 (12)	0.44495 (14)	0.11304 (9)	0.0456 (4)
H9A	0.3166	0.4549	0.0827	0.055*
H9B	0.4072	0.3684	0.0993	0.055*
O4	0.5000	0.27510 (13)	0.2500	0.0467 (4)
O5	0.33811 (9)	0.12882 (13)	0.36950 (7)	0.0628 (4)
H5A	0.2848	0.1462	0.3858	0.094*
N3	0.42148 (9)	0.09011 (11)	0.26254 (7)	0.0396 (3)
C10	0.45046 (15)	-0.24348 (15)	0.25741 (9)	0.0553 (5)
H10	0.4179	-0.3167	0.2617	0.066*
C11	0.39872 (13)	-0.13699 (15)	0.26601 (9)	0.0487 (4)
H11	0.3325	-0.1370	0.2765	0.058*
C12	0.45049 (11)	-0.03066 (13)	0.25817 (8)	0.0385 (3)
C13	0.5000	0.16353 (19)	0.2500	0.0379 (4)
C14	0.32941 (11)	0.13248 (16)	0.28957 (9)	0.0468 (4)
H14A	0.3162	0.2143	0.2722	0.056*
H14B	0.2752	0.0814	0.2701	0.056*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O1	0.0523 (7)	0.0567 (7)	0.0602 (7)	0.0107 (6)	0.0216 (5)	0.0184 (6)
O2	0.0488 (7)	0.0629 (8)	0.0571 (7)	0.0001 (6)	0.0090 (5)	0.0226 (6)
O3	0.0415 (6)	0.0568 (7)	0.0603 (7)	0.0122 (5)	0.0109 (5)	0.0124 (5)
N1	0.0370 (6)	0.0371 (6)	0.0436 (7)	0.0016 (5)	0.0077 (5)	0.0042 (5)
N2	0.0454 (7)	0.0366 (7)	0.0451 (7)	0.0036 (5)	0.0122 (5)	0.0077 (5)
C1	0.0514 (9)	0.0423 (8)	0.0477 (9)	0.0097 (7)	0.0092 (7)	0.0043 (7)
C2	0.0463 (10)	0.0675 (12)	0.0574 (10)	0.0190 (8)	0.0094 (8)	0.0066 (9)
C3	0.0382 (9)	0.0814 (13)	0.0561 (10)	0.0022 (8)	0.0110 (7)	0.0020 (9)
C4	0.0479 (9)	0.0547 (10)	0.0486 (9)	-0.0070 (7)	0.0143 (7)	0.0017 (7)
C5	0.0424 (8)	0.0402 (8)	0.0349 (7)	0.0031 (6)	0.0095 (6)	0.0012 (6)
C6	0.0392 (8)	0.0397 (8)	0.0335 (7)	0.0020 (6)	0.0079 (5)	-0.0001 (6)

C7	0.0419 (8)	0.0417 (8)	0.0366 (7)	0.0048 (6)	0.0084 (6)	0.0045 (6)
C8	0.0629 (11)	0.0366 (8)	0.0547 (10)	0.0025 (7)	0.0103 (8)	0.0081 (7)
C9	0.0437 (8)	0.0404 (8)	0.0528 (9)	-0.0040 (7)	0.0052 (7)	0.0027 (7)
O4	0.0426 (8)	0.0344 (8)	0.0635 (10)	0.000	0.0068 (7)	0.000
O5	0.0400 (6)	0.0918 (10)	0.0575 (7)	0.0004 (6)	0.0105 (5)	-0.0143 (7)
N3	0.0328 (6)	0.0377 (7)	0.0490 (7)	-0.0017 (5)	0.0069 (5)	-0.0008 (5)
C10	0.0860 (13)	0.0370 (8)	0.0426 (9)	-0.0116 (8)	0.0021 (8)	0.0014 (7)
C11	0.0562 (10)	0.0459 (9)	0.0439 (8)	-0.0124 (7)	0.0032 (7)	0.0019 (7)
C12	0.0416 (8)	0.0376 (8)	0.0363 (7)	-0.0008 (6)	0.0020 (6)	0.0000 (6)
C13	0.0346 (10)	0.0381 (11)	0.0410 (11)	0.000	0.0025 (8)	0.000
C14	0.0329 (8)	0.0504 (9)	0.0575 (10)	0.0001 (7)	0.0052 (6)	-0.0001 (7)

*Geometric parameters (Å, °)*

O1—C9	1.3981 (19)	C5—C6	1.395 (2)
O1—H1A	0.82	C8—H8A	0.97
O2—C8	1.396 (2)	C8—H8B	0.97
O2—H2A	0.82	C9—H9A	0.97
O3—C7	1.2304 (18)	C9—H9B	0.97
N1—C7	1.3759 (19)	O4—C13	1.237 (3)
N1—C6	1.3980 (19)	O5—C14	1.406 (2)
N1—C9	1.4509 (19)	O5—H5A	0.82
N2—C7	1.371 (2)	N3—C13	1.3704 (17)
N2—C5	1.3925 (19)	N3—C12	1.3990 (19)
N2—C8	1.451 (2)	N3—C14	1.4462 (19)
C1—C6	1.378 (2)	C10—C10 <sup>i</sup>	1.387 (4)
C1—C2	1.387 (2)	C10—C11	1.387 (3)
C1—H1	0.93	C11—C12	1.384 (2)
C2—C3	1.388 (3)	C11—H11	0.93
C2—H2	0.93	C12—C12 <sup>i</sup>	1.393 (3)
C3—C4	1.387 (3)	C13—N3 <sup>i</sup>	1.3704 (17)
C3—H3	0.93	C14—H14A	0.97
C4—C5	1.380 (2)	C14—H14B	0.97
C4—H4	0.93		
C9—O1—H1A	109.4	O2—C8—H8B	109.2
C8—O2—H2A	109.5	N2—C8—H8B	109.2
C7—N1—C6	109.53 (12)	H8A—C8—H8B	107.9
C7—N1—C9	123.27 (13)	O1—C9—N1	112.84 (13)
C6—N1—C9	127.20 (12)	O1—C9—H9A	109.0
C7—N2—C5	109.76 (12)	N1—C9—H9A	109.0
C7—N2—C8	124.58 (14)	O1—C9—H9B	109.0
C5—N2—C8	125.66 (13)	N1—C9—H9B	109.0
C6—C1—C2	117.39 (16)	H9A—C9—H9B	107.8
C6—C1—H1	121.3	C14—O5—H5A	109.5
C2—C1—H1	121.3	C13—N3—C12	109.55 (12)
C1—C2—C3	121.40 (16)	C13—N3—C14	124.00 (13)
C1—C2—H2	119.3	C12—N3—C14	125.57 (13)

C3—C2—H2	119.3	C10 <sup>i</sup> —C10—C11	121.66 (10)
C4—C3—C2	121.16 (16)	C10 <sup>i</sup> —C10—H10	119.2
C4—C3—H3	119.4	C11—C10—H10	119.2
C2—C3—H3	119.4	C12—C11—C10	116.74 (16)
C5—C4—C3	117.36 (16)	C12—C11—H11	121.6
C5—C4—H4	121.3	C10—C11—H11	121.6
C3—C4—H4	121.3	C11—C12—C12 <sup>i</sup>	121.59 (10)
C4—C5—N2	131.54 (15)	C11—C12—N3	131.53 (14)
C4—C5—C6	121.39 (14)	C12 <sup>i</sup> —C12—N3	106.88 (8)
N2—C5—C6	107.05 (13)	O4—C13—N3	126.43 (9)
C1—C6—C5	121.26 (14)	O4—C13—N3 <sup>i</sup>	126.43 (9)
C1—C6—N1	131.97 (14)	N3—C13—N3 <sup>i</sup>	107.14 (18)
C5—C6—N1	106.76 (12)	O5—C14—N3	108.05 (13)
O3—C7—N2	125.99 (14)	O5—C14—H14A	110.1
O3—C7—N1	127.14 (14)	N3—C14—H14A	110.1
N2—C7—N1	106.87 (12)	O5—C14—H14B	110.1
O2—C8—N2	111.86 (14)	N3—C14—H14B	110.1
O2—C8—H8A	109.2	H14A—C14—H14B	108.4
N2—C8—H8A	109.2		
C6—C1—C2—C3	1.2 (3)	C8—N2—C7—N1	177.40 (14)
C1—C2—C3—C4	-1.5 (3)	C6—N1—C7—O3	-178.71 (15)
C2—C3—C4—C5	0.1 (3)	C9—N1—C7—O3	1.2 (2)
C3—C4—C5—N2	179.76 (16)	C6—N1—C7—N2	1.10 (16)
C3—C4—C5—C6	1.6 (2)	C9—N1—C7—N2	-178.97 (13)
C7—N2—C5—C4	-177.04 (16)	C7—N2—C8—O2	-105.28 (17)
C8—N2—C5—C4	4.1 (3)	C5—N2—C8—O2	73.5 (2)
C7—N2—C5—C6	1.36 (16)	C7—N1—C9—O1	-88.27 (17)
C8—N2—C5—C6	-177.55 (14)	C6—N1—C9—O1	91.64 (17)
C2—C1—C6—C5	0.5 (2)	C10 <sup>i</sup> —C10—C11—C12	0.8 (3)
C2—C1—C6—N1	-179.32 (15)	C10—C11—C12—C12 <sup>i</sup>	0.5 (3)
C4—C5—C6—C1	-1.9 (2)	C10—C11—C12—N3	179.38 (15)
N2—C5—C6—C1	179.46 (14)	C13—N3—C12—C11	-179.51 (14)
C4—C5—C6—N1	177.94 (13)	C14—N3—C12—C11	10.9 (3)
N2—C5—C6—N1	-0.65 (16)	C13—N3—C12—C12 <sup>i</sup>	-0.50 (18)
C7—N1—C6—C1	179.60 (16)	C14—N3—C12—C12 <sup>i</sup>	-170.07 (15)
C9—N1—C6—C1	-0.3 (3)	C12—N3—C13—O4	-179.81 (7)
C7—N1—C6—C5	-0.27 (16)	C14—N3—C13—O4	-10.04 (16)
C9—N1—C6—C5	179.81 (14)	C12—N3—C13—N3 <sup>i</sup>	0.19 (7)
C5—N2—C7—O3	178.30 (15)	C14—N3—C13—N3 <sup>i</sup>	169.96 (16)
C8—N2—C7—O3	-2.8 (2)	C13—N3—C14—O5	-89.75 (16)
C5—N2—C7—N1	-1.52 (16)	C12—N3—C14—O5	78.38 (18)

Symmetry code: (i)  $-x+1, y, -z+1/2$ .

#### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1A $\cdots$ O4	0.82	1.99	2.8003 (16)	169

---

O2—H2A···O5 <sup>ii</sup>	0.82	1.93	2.7503 (18)	176
O5—H5A···O3 <sup>iii</sup>	0.82	1.84	2.6551 (17)	175
C3—H3···O2 <sup>iv</sup>	0.93	2.58	3.489 (2)	164
C14—H14B···O1 <sup>iii</sup>	0.97	2.54	3.364 (2)	143

---

Symmetry codes: (ii)  $x, -y+1, z-1/2$ ; (iii)  $-x+1/2, y-1/2, -z+1/2$ ; (iv)  $-x+3/2, -y+3/2, -z$ .