

1-[2-(2,4-Dichlorobenzyl)oxy]-2-(2-furyl)-ethyl]-1*H*-1,2,4-triazole

Özden Özel Güven,^a Hakan Tahtacı,^a Simon J. Coles^b and Tuncer Hökelek^{c*}

^aDepartment of Chemistry, Zonguldak Karaelmas University, 67100 Zonguldak, Turkey, ^bDepartment of Chemistry, Southampton University, SO17 1BJ Southampton, England, and ^cDepartment of Physics, Hacettepe University, 06800 Beytepe, Ankara, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 21 October 2009; accepted 23 October 2009

Key indicators: single-crystal X-ray study; $T = 120\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.052; wR factor = 0.161; data-to-parameter ratio = 17.8.

In the molecule of the title compound, $\text{C}_{15}\text{H}_{13}\text{Cl}_2\text{N}_3\text{O}_2$, the triazole ring is oriented at dihedral angles of $14.8(2)$ and $81.5(1)^\circ$ to the furan and dichlorobenzene rings, respectively. The dihedral angle between the dichlorobenzene and furan rings is $86.3(2)^\circ$. An intramolecular $\text{C}-\text{H}\cdots\text{O}$ hydrogen bond results in the formation of a planar [maximum deviation $0.012(2)\text{ \AA}$] five-membered ring, which is oriented at a dihedral angle of $0.90(7)^\circ$ with respect to the dichlorobenzene ring. There is an intermolecular $\text{C}-\text{H}\cdots\pi$ contact between the methylene group and the dichlorobenzene ring.

Related literature

For general background to the use of ether structures containing $1H$ -imidazole and $1H$ -1,2,4-triazole rings as anti-fungal agents, see: Caira *et al.* (2004); Godefroi *et al.* (1969); Özel Güven *et al.* (2007a,b); Paulvannan *et al.* (2001); Peeters *et al.* (1996); Wahbi *et al.* (1995). For related structures, see: Freer *et al.* (1986); Özel Güven *et al.* (2008a,b,c,d,e,f); Peeters *et al.* (1979).



Experimental

Crystal data

$\text{C}_{15}\text{H}_{13}\text{Cl}_2\text{N}_3\text{O}_2$
 $M_r = 338.18$

Monoclinic, $P2_1/n$
 $a = 10.6057(2)\text{ \AA}$

$b = 13.3560(3)\text{ \AA}$
 $c = 11.1919(2)\text{ \AA}$
 $\beta = 101.170(1)^\circ$
 $V = 1555.30(5)\text{ \AA}^3$
 $Z = 4$

Mo $K\alpha$ radiation
 $\mu = 0.43\text{ mm}^{-1}$
 $T = 120\text{ K}$
 $0.50 \times 0.35 \times 0.20\text{ mm}$

Data collection

Bruker-Nonius Kappa CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Sheldrick, 2007)
 $R_{\text{min}} = 0.815$, $T_{\text{max}} = 0.920$

6687 measured reflections
3547 independent reflections
2522 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.161$
 $S = 1.06$
3547 reflections

199 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.77\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.33\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
C11—H11 \cdots O1	0.93	2.35	2.702 (3)	102
C9—H9B \cdots Cg1 ⁱ	0.97	2.90	3.775 (3)	151

Symmetry code: (i) $-x + 1, -y, -z + 1$. Cg1 is the centroid of the C10—C15 ring.

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *DENZO* (Otwinowski & Minor, 1997) and *COLLECT*; data reduction: *DENZO* and *COLLECT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999) and *PLATON* (Spek, 2009).

The authors acknowledge the Zonguldak Karaelmas University Research Fund (Project No. 2008-13-02-06) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CI2951).

References

- Caira, M. R., Alkhamis, K. A. & Obaidat, R. M. (2004). *J. Pharm. Sci.* **93**, 601–611.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Freer, A. A., Pearson, A. & Salole, E. G. (1986). *Acta Cryst.* **C42**, 1350–1352.
- Godefroi, E. F., Heeres, J., Van Cutsem, J. & Janssen, P. A. J. (1969). *J. Med. Chem.* **12**, 784–791.
- Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Özel Güven, Ö., Erdoğan, T., Coles, S. J. & Hökelek, T. (2008a). *Acta Cryst.* **E64**, o1437.
- Özel Güven, Ö., Erdoğan, T., Coles, S. J. & Hökelek, T. (2008b). *Acta Cryst.* **E64**, o1496–o1497.
- Özel Güven, Ö., Erdoğan, T., Coles, S. J. & Hökelek, T. (2008c). *Acta Cryst.* **E64**, o1588–o1589.
- Özel Güven, Ö., Erdoğan, T., Coles, S. J. & Hökelek, T. (2008d). *Acta Cryst.* **E64**, o1655–o1656.
- Özel Güven, Ö., Erdoğan, T., Göker, H. & Yıldız, S. (2007a). *Bioorg. Med. Chem. Lett.* **17**, 2233–2236.

- Özel Güven, Ö., Erdogan, T., Göker, H. & Yıldız, S. (2007b). *J. Heterocycl. Chem.* **44**, 731–734.
- Özel Güven, Ö., Tahtaci, H., Coles, S. J. & Hökelek, T. (2008e). *Acta Cryst. E* **64**, o1914–o1915.
- Özel Güven, Ö., Tahtaci, H., Tahir, M. N. & Hökelek, T. (2008f). *Acta Cryst. E* **64**, o2465.
- Paulvannan, K., Hale, R., Sedehi, D. & Chen, T. (2001). *Tetrahedron*, **57**, 9677–9682.
- Peeters, O. M., Blaton, N. M. & De Ranter, C. J. (1979). *Acta Cryst. B* **35**, 2461–2464.
- Peeters, O. M., Blaton, N. M. & De Ranter, C. J. (1996). *Acta Cryst. C* **52**, 2225–2229.
- Sheldrick, G. M. (2007). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Wahbi, Y., Caujolle, R., Tournaire, C., Payard, M., Linas, M. D. & Seguela, J. P. (1995). *Eur. J. Med. Chem.* **30**, 955–962.

supporting information

Acta Cryst. (2009). E65, o2868–o2869 [https://doi.org/10.1107/S1600536809044018]

1-[2-(2,4-Dichlorobenzyl)oxy]-2-(2-furyl)ethyl]-1*H*-1,2,4-triazole

Özden Özel Güven, Hakan Tahtacı, Simon J. Coles and Tuncer Hökelek

S1. Comment

In recent years, among antifungal agents, azole derivatives still have an important place in the class of systemic antifungal drugs. Some ether structures containing 1*H*-imidazole ring like micozanole, ecozanole and sulconazole have been synthesized and developed for clinical uses as antifungal agents (Godefroi *et al.*, 1969). The crystal structures of these ether derivatives like miconazole (Peeters *et al.*, 1979), econazole (Freer *et al.*, 1986) have been reported previously. Also, antifungal activity of aromatic ethers possessing 1*H*-1,2,4-triazole ring have been reported (Wahbi *et al.*, 1995). Itraconazole (Peeters *et al.*, 1996) and fluconazole (Caira *et al.*, 2004) are 1*H*-1,2,4-triazole ring containing azole derivatives. 1,2,4-Triazoles are biologically interesting molecules and their chemistry is receiving considerable attention due to antihypertensive, antifungal and antibacterial properties (Paulvannan *et al.*, 2001). Ether structures possessing 1*H*-benzimidazole ring have been reported to show antibacterial activity more than antifungal activity (Özel Güven *et al.*, 2007a,b). The crystal structures of 1*H*-benzimidazole ring containing ether derivatives (Özel Güven *et al.*, 2008a,b,c,d) and also, 1*H*-1,2,4-triazole ring containing ether derivatives have been reported recently (Özel Güven *et al.*, 2008e,f). Now, we report herein the crystal structure of 2,4-dichloro- derivative of 1*H*-1,2,4-triazole and furyl rings containing ether structure.

In the molecule of the title compound (Fig. 1) the bond lengths and angles are generally within normal ranges. The planar triazole ring is oriented with respect to the furan and dichlorobenzene rings at dihedral angles of 14.8 (2) $^{\circ}$ and 81.5 (1) $^{\circ}$, respectively. Atoms C3, C4 and C9 are -0.021 (2), 0.029 (2) and 0.034 (4) Å away from the planes of the triazole, furan and dichlorobenzene, respectively. So, they are nearly coplanar with the adjacent rings. The dichlorobenzene ring is oriented with respect to the furan ring at a dihedral angle of 86.3 (2) $^{\circ}$. An intramolecular C—H \cdots O hydrogen bond (Table 1) results in the formation of a planar five-membered ring (O1/H11/C9—C11), which is oriented with respect to dichlorobenzene ring at a dihedral angle of 0.90 (7) $^{\circ}$. So, they are coplanar.


In the crystal, an intermolecular C—H \cdots π interaction (Table 1) is observed between the methylene group and the dichlorobenzene ring. A view of the molecular packing in the crystal is shown in Fig. 2.

S2. Experimental


The title compound was synthesized by the reaction of 1-(furan-2-yl)-2-(1*H*-1,2,4-triazol-1-yl)ethanol (unpublished results) with NaH and appropriate benzyl halide. To a solution of alcohol (400 mg, 2.232 mmol) in DMF (4 ml) was added NaH (89 mg, 2.232 mmol) in small fractions. The appropriate benzyl halide (436 mg, 2.232 mmol) was added dropwise. The mixture was stirred at room temperature for 3 h, and excess hydride was decomposed with methyl alcohol (5 ml). After evaporation to dryness under reduced pressure, the crude residue was suspended with water and extracted with methylene chloride. The organic layer was dried over anhydrous sodium sulfate and then evaporated to dryness. The crude residue was purified by chromatography on a silica-gel column using chloroform as eluent. Crystals suitable for X-ray analysis were obtained by the recrystallization of the ether from 2-propanol (yield; 355 mg, 47%).

S3. Refinement

H atoms were positioned geometrically, with C-H = 0.93, 0.98 and 0.97 Å for aromatic, methine and methylene H, respectively, and constrained to ride on their parent atoms with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

**Figure 1**


The molecular structure of the title compound with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. The dashed line indicates a hydrogen bond.

**Figure 2**

Part of a packing diagram. Hydrogen atoms have been omitted for clarity.

1-[2-(2,4-Dichlorobenzyl)oxy]-2-(2-furyl)ethyl]-1*H*-1,2,4-triazole

Crystal data



$M_r = 338.18$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 10.6057 (2) \text{ \AA}$

$b = 13.3560 (3) \text{ \AA}$

$c = 11.1919 (2) \text{ \AA}$

$\beta = 101.170 (1)^\circ$

$V = 1555.30 (5) \text{ \AA}^3$

$Z = 4$

$F(000) = 696$

$D_x = 1.444 \text{ Mg m}^{-3}$

$\text{Mo } K\alpha \text{ radiation, } \lambda = 0.71073 \text{ \AA}$

Cell parameters from 3274 reflections

$\theta = 2.9\text{--}27.5^\circ$

$\mu = 0.43 \text{ mm}^{-1}$

$T = 120 \text{ K}$

Plate, colourless

$0.50 \times 0.35 \times 0.20 \text{ mm}$

Data collection

Bruker–Nonius Kappa CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 2007)
 $T_{\min} = 0.815$, $T_{\max} = 0.920$

6687 measured reflections
3547 independent reflections
2522 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$
 $\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.0^\circ$
 $h = -13 \rightarrow 13$
 $k = -17 \rightarrow 17$
 $l = -14 \rightarrow 14$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.052$
 $wR(F^2) = 0.161$
 $S = 1.06$
3547 reflections
199 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0852P)^2 + 0.4221P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.77 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.33 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl1	0.48306 (8)	0.29223 (6)	0.76970 (7)	0.0823 (3)
Cl2	0.66147 (6)	0.12991 (6)	0.39547 (7)	0.0719 (3)
O1	0.26246 (14)	0.02290 (13)	0.27553 (13)	0.0523 (4)
O2	0.20740 (18)	0.07592 (15)	0.01684 (16)	0.0673 (5)
N1	0.06121 (18)	-0.11249 (15)	0.27806 (16)	0.0498 (5)
N2	0.0878 (2)	-0.20960 (17)	0.3093 (2)	0.0648 (6)
N3	0.0001 (2)	-0.13162 (18)	0.45167 (19)	0.0644 (6)
C1	0.0492 (3)	-0.2156 (2)	0.4135 (3)	0.0657 (7)
H1	0.0556	-0.2746	0.4583	0.079*
C2	0.0106 (3)	-0.0688 (2)	0.3648 (2)	0.0598 (6)
H2	-0.0142	-0.0019	0.3637	0.072*
C3	0.0914 (2)	-0.0710 (2)	0.16712 (19)	0.0550 (6)
H3A	0.0425	-0.0100	0.1464	0.066*
H3B	0.0658	-0.1183	0.1011	0.066*
C4	0.2337 (2)	-0.04811 (18)	0.17976 (19)	0.0495 (5)
H4	0.2831	-0.1093	0.2036	0.059*

C5	0.2625 (2)	-0.01101 (19)	0.0610 (2)	0.0505 (5)
C6	0.2450 (3)	0.0932 (3)	-0.0918 (2)	0.0760 (8)
H6	0.2218	0.1486	-0.1416	0.091*
C7	0.3176 (4)	0.0206 (3)	-0.1140 (3)	0.0926 (11)
H7	0.3551	0.0147	-0.1823	0.111*
C8	0.3303 (3)	-0.0490 (3)	-0.0155 (3)	0.0884 (10)
H8	0.3765	-0.1086	-0.0071	0.106*
C9	0.3934 (2)	0.05052 (18)	0.30581 (19)	0.0460 (5)
H9A	0.4171	0.0885	0.2396	0.055*
H9B	0.4468	-0.0089	0.3191	0.055*
C10	0.4136 (2)	0.11323 (16)	0.41992 (18)	0.0430 (5)
C11	0.3143 (2)	0.13351 (19)	0.4814 (2)	0.0528 (6)
H11	0.2325	0.1087	0.4508	0.063*
C12	0.3349 (2)	0.1896 (2)	0.5868 (2)	0.0590 (6)
H12	0.2674	0.2025	0.6267	0.071*
C13	0.4554 (2)	0.22648 (19)	0.6326 (2)	0.0547 (6)
C14	0.5566 (2)	0.20958 (18)	0.5743 (2)	0.0532 (6)
H14	0.6379	0.2351	0.6052	0.064*
C15	0.5333 (2)	0.15318 (17)	0.4678 (2)	0.0470 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.1025 (6)	0.0819 (6)	0.0616 (4)	-0.0194 (4)	0.0135 (4)	-0.0290 (4)
Cl2	0.0461 (4)	0.0894 (6)	0.0834 (5)	-0.0112 (3)	0.0204 (3)	-0.0183 (4)
O1	0.0435 (8)	0.0694 (11)	0.0444 (8)	-0.0108 (7)	0.0093 (6)	-0.0156 (7)
O2	0.0722 (12)	0.0746 (13)	0.0579 (10)	0.0045 (10)	0.0191 (8)	0.0056 (9)
N1	0.0488 (10)	0.0542 (12)	0.0468 (10)	-0.0102 (9)	0.0101 (8)	-0.0052 (8)
N2	0.0735 (14)	0.0536 (13)	0.0701 (14)	-0.0059 (11)	0.0206 (11)	-0.0055 (10)
N3	0.0621 (13)	0.0746 (16)	0.0615 (12)	0.0017 (11)	0.0249 (10)	0.0094 (11)
C1	0.0627 (16)	0.0617 (17)	0.0743 (17)	-0.0078 (13)	0.0171 (13)	0.0120 (13)
C2	0.0644 (15)	0.0610 (16)	0.0577 (13)	0.0051 (12)	0.0213 (11)	0.0013 (12)
C3	0.0521 (13)	0.0703 (16)	0.0415 (11)	-0.0148 (11)	0.0069 (9)	-0.0046 (10)
C4	0.0505 (12)	0.0546 (14)	0.0431 (11)	-0.0075 (10)	0.0086 (9)	-0.0083 (10)
C5	0.0482 (12)	0.0583 (14)	0.0466 (11)	-0.0061 (11)	0.0133 (9)	-0.0116 (10)
C6	0.081 (2)	0.095 (2)	0.0525 (14)	-0.0173 (18)	0.0141 (13)	0.0096 (15)
C7	0.101 (2)	0.125 (3)	0.0621 (17)	-0.008 (2)	0.0413 (17)	-0.0092 (18)
C8	0.099 (2)	0.100 (3)	0.0766 (19)	0.0192 (19)	0.0428 (18)	-0.0125 (17)
C9	0.0430 (11)	0.0520 (13)	0.0437 (10)	-0.0073 (10)	0.0101 (8)	-0.0030 (9)
C10	0.0439 (11)	0.0428 (12)	0.0416 (10)	-0.0036 (9)	0.0067 (8)	0.0020 (8)
C11	0.0455 (12)	0.0633 (16)	0.0495 (12)	-0.0082 (10)	0.0094 (9)	-0.0073 (10)
C12	0.0594 (14)	0.0677 (16)	0.0518 (13)	-0.0041 (12)	0.0157 (11)	-0.0110 (11)
C13	0.0670 (15)	0.0499 (14)	0.0456 (12)	-0.0063 (11)	0.0072 (10)	-0.0063 (10)
C14	0.0548 (13)	0.0495 (14)	0.0508 (12)	-0.0109 (10)	-0.0009 (10)	0.0008 (10)
C15	0.0434 (11)	0.0479 (13)	0.0493 (11)	-0.0030 (9)	0.0083 (9)	0.0026 (9)

Geometric parameters (\AA , $^\circ$)

C11—C13	1.742 (2)	C5—C8	1.321 (3)
C12—C15	1.740 (2)	C6—H6	0.93
O1—C4	1.419 (3)	C7—C6	1.293 (5)
O1—C9	1.413 (2)	C7—H7	0.93
O2—C5	1.350 (3)	C8—C7	1.428 (5)
O2—C6	1.370 (3)	C8—H8	0.93
N1—N2	1.358 (3)	C9—H9A	0.97
N1—C2	1.332 (3)	C9—H9B	0.97
N1—C3	1.451 (3)	C10—C15	1.386 (3)
N2—C1	1.311 (3)	C10—C11	1.391 (3)
N3—C1	1.341 (4)	C10—C9	1.508 (3)
N3—C2	1.305 (3)	C11—C12	1.379 (3)
C1—H1	0.93	C11—H11	0.93
C2—H2	0.93	C12—H12	0.93
C3—H3A	0.97	C13—C12	1.373 (3)
C3—H3B	0.97	C14—C13	1.378 (4)
C4—C3	1.519 (3)	C14—C15	1.391 (3)
C4—C5	1.504 (3)	C14—H14	0.93
C4—H4	0.98		
C9—O1—C4	114.40 (16)	C6—C7—C8	108.0 (3)
C5—O2—C6	106.9 (2)	C6—C7—H7	126.0
N2—N1—C3	121.0 (2)	C8—C7—H7	126.0
C2—N1—N2	108.9 (2)	C5—C8—C7	105.6 (3)
C2—N1—C3	130.1 (2)	C5—C8—H8	127.2
C1—N2—N1	101.6 (2)	C7—C8—H8	127.2
C2—N3—C1	101.9 (2)	O1—C9—C10	108.67 (17)
N2—C1—N3	116.1 (2)	O1—C9—H9A	110.0
N2—C1—H1	121.9	O1—C9—H9B	110.0
N3—C1—H1	121.9	C10—C9—H9A	110.0
N1—C2—H2	124.3	C10—C9—H9B	110.0
N3—C2—N1	111.4 (2)	H9A—C9—H9B	108.3
N3—C2—H2	124.3	C11—C10—C9	122.02 (19)
N1—C3—C4	112.17 (18)	C15—C10—C9	120.72 (19)
N1—C3—H3A	109.2	C15—C10—C11	117.3 (2)
N1—C3—H3B	109.2	C10—C11—H11	119.4
C4—C3—H3A	109.2	C12—C11—C10	121.3 (2)
C4—C3—H3B	109.2	C12—C11—H11	119.4
H3A—C3—H3B	107.9	C11—C12—H12	120.1
O1—C4—C5	113.35 (19)	C13—C12—C11	119.8 (2)
O1—C4—C3	105.63 (18)	C13—C12—H12	120.1
O1—C4—H4	109.0	C12—C13—Cl1	119.7 (2)
C3—C4—H4	109.0	C12—C13—C14	121.2 (2)
C5—C4—C3	110.63 (18)	C14—C13—Cl1	119.08 (19)
C5—C4—H4	109.0	C13—C14—C15	118.0 (2)
O2—C5—C4	117.3 (2)	C13—C14—H14	121.0

C8—C5—O2	110.1 (3)	C15—C14—H14	121.0
C8—C5—C4	132.5 (3)	C10—C15—Cl2	119.41 (17)
O2—C6—H6	125.3	C10—C15—C14	122.5 (2)
C7—C6—O2	109.3 (3)	C14—C15—Cl2	118.09 (18)
C7—C6—H6	125.3		
C9—O1—C4—C3	177.01 (19)	C3—C4—C5—C8	-113.9 (3)
C9—O1—C4—C5	-61.7 (3)	O2—C5—C8—C7	0.9 (4)
C4—O1—C9—C10	-171.48 (18)	C4—C5—C8—C7	178.2 (3)
C6—O2—C5—C4	-178.8 (2)	C8—C7—C6—O2	-0.2 (4)
C6—O2—C5—C8	-1.0 (3)	C5—C8—C7—C6	-0.4 (4)
C5—O2—C6—C7	0.8 (3)	C11—C10—C9—O1	2.1 (3)
C2—N1—N2—C1	0.4 (3)	C15—C10—C9—O1	-177.9 (2)
C3—N1—N2—C1	178.7 (2)	C9—C10—C11—C12	178.9 (2)
N2—N1—C2—N3	-0.9 (3)	C15—C10—C11—C12	-1.1 (4)
C3—N1—C2—N3	-179.0 (2)	C9—C10—C15—Cl2	-0.3 (3)
N2—N1—C3—C4	-77.0 (3)	C9—C10—C15—C14	-178.6 (2)
C2—N1—C3—C4	100.9 (3)	C11—C10—C15—Cl2	179.67 (18)
N1—N2—C1—N3	0.2 (3)	C11—C10—C15—C14	1.4 (3)
C2—N3—C1—N2	-0.7 (3)	C10—C11—C12—C13	0.0 (4)
C1—N3—C2—N1	1.0 (3)	C11—C13—C12—C11	-176.9 (2)
O1—C4—C3—N1	-60.3 (3)	C14—C13—C12—C11	0.9 (4)
C5—C4—C3—N1	176.7 (2)	C15—C14—C13—Cl1	177.24 (18)
O1—C4—C5—O2	-55.2 (3)	C15—C14—C13—C12	-0.6 (4)
O1—C4—C5—C8	127.7 (3)	C13—C14—C15—C10	-0.6 (4)
C3—C4—C5—O2	63.3 (3)	C13—C14—C15—Cl2	-178.91 (19)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
C11—H11···O1	0.93	2.35	2.702 (3)	102
C9—H9B···Cg1 ⁱ	0.97	2.90	3.775 (3)	151

Symmetry code: (i) $-x+1, -y, -z+1$.