

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Dimethyl(2,2':6',2''-terpyridine- κ^3N,N',N'')zinc(II)

Tamila Shalumova and Joseph M. Tanski*

 Department of Chemistry, Vassar College, Poughkeepsie, NY 12604, USA
 Correspondence e-mail: jotanski@vassar.edu


Received 21 September 2009; accepted 1 October 2009

 Key indicators: single-crystal X-ray study; $T = 125$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.022; wR factor = 0.061; data-to-parameter ratio = 18.7.

The title compound, $[\text{Zn}(\text{CH}_3)_2(\text{C}_{15}\text{H}_{11}\text{N}_3)]$, was synthesized by the addition of dimethylzinc to 2,2':6',2''-terpyridine and was crystallized by the slow evaporation of THF. The pentacoordinate Zn^{II} atom, lying on a twofold rotation axis, displays a distorted trigonal-bipyramidal geometry, with two terminal N atoms at the axial positions and the central N atom and two methyl C atoms at the equatorial positions.

Related literature

For the crystal structures of terpyridine dichloridozinc(II) compounds, see: Corbridge & Cox (1956); Einstein & Penfold (1966); Vlasse *et al.* (1983). For examples of other substituted terpyridine zinc(II) compounds, see: Harrison *et al.* (1986); Hou *et al.* (2004). The structure of a bipyridine dimethylzinc(II) compound was reported by Wissing *et al.* (1994).



Experimental

Crystal data

 $[\text{Zn}(\text{CH}_3)_2(\text{C}_{15}\text{H}_{11}\text{N}_3)]$
 $M_r = 328.71$

 Monoclinic, $C2/c$
 $a = 17.4250$ (11) Å
 $b = 9.1083$ (6) Å
 $c = 11.7595$ (14) Å
 $\beta = 127.193$ (1)°
 $V = 1486.8$ (2) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 1.65$ mm⁻¹
 $T = 125$ K
 $0.23 \times 0.13 \times 0.06$ mm

Data collection

 Bruker APEXII CCD
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.703$, $T_{\text{max}} = 0.908$

 9483 measured reflections
 1837 independent reflections
 1710 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.022$
 $wR(F^2) = 0.061$
 $S = 1.09$
 1837 reflections

 98 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.41$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.27$ e Å⁻³
Table 1

Selected bond lengths (Å).

Zn—C1	2.0282 (15)	Zn—N2	2.2603 (16)
Zn—N1	2.3381 (12)		

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINTE* (Bruker, 2007); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

This work was supported by Vassar College. X-ray facilities were provided by the US National Science Foundation (grant No. 0521237 to JMT).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HY2233).

References

- Bruker (2007). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Corbridge, D. E. C. & Cox, E. G. (1956). *J. Chem. Soc.* pp. 594–603.
- Einstein, F. W. B. & Penfold, B. R. (1966). *Acta Cryst.* **20**, 924–926.
- Harrison, P. G., Begley, M. J., Kikabhai, T. & Killer, F. (1986). *J. Chem. Soc. Dalton Trans.* pp. 929–938.
- Hou, L., Li, D. & Ng, S. W. (2004). *Acta Cryst.* **E60**, m1734–m1735.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Vlasse, M., Rojo, T. & Beltran-Porter, D. (1983). *Acta Cryst.* **C39**, 560–563.
- Wissing, E., Kaupp, M., Boersma, J., Spek, A. L. & van Koten, G. (1994). *Organometallics*, **13**, 2349–2356.

supporting information

Acta Cryst. (2009). E65, m1325 [https://doi.org/10.1107/S1600536809040136]

Dimethyl(2,2':6',2''-terpyridine- κ^3N,N',N'')zinc(II)**Tamila Shalumova and Joseph M. Tanski****S1. Comment**

The chelating ligand 2,2':6',2''-terpyridine coordinates to a variety of first-row transition metal Lewis acids. Specifically, it can coordinate to disubstituted zinc compounds, allowing for the formation of trigonal bipyramidal zinc(II) complexes (Harrison *et al.*, 1986). The structure of terpyridine dichlorozinc(II) has been reported a number of times with different results (Corbridge & Cox, 1956; Einstein & Penfold, 1966; Vlasse *et al.*, 1983). In addition to substituents on the zinc(II) center, substituents on the terpyridine are also known (Hou *et al.*, 2004). The structure presented here is the first known structure in a class of terpyridine zinc(II) compounds with two alkyl groups bound directly to the metal center.

The title compound (Fig. 1) was obtained by the reaction of dimethylzinc with 2,2':6',2''-terpyridine. It exhibits a distorted trigonal bipyramidal geometry about the metal center and has the two terminal N atoms in the axial positions, with a bond length of 2.3381 (12) Å (Table 1). The central N atom, coordinated to the zinc *via* the equatorial position, has a slightly smaller bond length, 2.2603 (16) Å, due to the size of the ligand, which is not able to wrap around the metal 180°. The N1—Zn—N1ⁱ bond angle is 140.52 (6)° [symmetry code: (i) -x, y, 1/2-z], illustrating the degree to which the compound is distorted from a perfectly trigonal bipyramid. The Zn—C bond length is 2.0282 (15) Å, and the C1—Zn—C1ⁱ bond angle is 133.21 (9)°: slightly greater than the expected 120° between equatorial atoms. Interestingly, the Zn—N bond lengths shown here are about 0.2 Å longer than those reported for similar terpyridine zinc(II) complexes, while the Zn—C length is expectedly shorter than the Zn—Cl and Zn—S lengths (Harrison *et al.*, 1986; Hou *et al.*, 2004; Vlasse *et al.*, 1983). However, the Zn—C bond length is similar to that reported for a bipyridine dimethylzinc(II) complex, characterized by Wessing *et al.* (1994). The terpyridyl N1—Zn—N1ⁱ and N1—Zn—N2 bond angles are in agreement with those reported in the literature (Harrison *et al.*, 1986; Hou *et al.*, 2004; Vlasse *et al.*, 1983).

S2. Experimental

Under a nitrogen atmosphere, dimethylzinc (2 M in toluene, 1.07 ml, 2.14 mmol) was added to a stirring solution of terpyridine (0.511 g, 2.19 mmol) in toluene (3.5 ml). The resulting orange precipitate, which is extremely sensitive to hydrolysis, was filtered and dried *in vacuo* (yield 65%, 0.46 g). Crystallization was achieved by slow evaporation of THF in a nitrogen filled glovebox, which yielded yellow plates within five days. Analysis, calculated for C₁₇H₁₇N₃Zn: C 62.11, H 5.21, N 12.78%; found: C 59.18, H 5.09, N 12.12%. ¹H NMR (300 MHz, C₆D₆): δ -0.529 (s, 6H, -CH₃).

S3. Refinement

A suitable crystal was mounted in a nylon loop with Paratone-*N* cryoprotectant oil and data was collected on a Bruker APEXII CCD platform diffractometer. H atoms were included in calculated positions and were refined using a riding model, with C—H = 0.95 (aromatic) and 0.98 (methyl) Å and with $U_{\text{iso}}(\text{H}) = 1.2(1.5 \text{ for methyl})U_{\text{eq}}(\text{C})$.




Figure 1

Molecular structure of the title compound, with displacement ellipsoids shown at the 50% probability level. [Symmetry code: (i) $-x, y, 1/2-z$.]

Dimethyl(2,2':6',2''-terpyridine- κ^3N,N',N'')zinc(II)

Crystal data

$[\text{Zn}(\text{CH}_3)_2(\text{C}_{15}\text{H}_{11}\text{N}_3)]$

$M_r = 328.71$

Monoclinic, $C2/c$

Hall symbol: $-C\ 2yc$

$a = 17.4250(11)\ \text{\AA}$

$b = 9.1083(6)\ \text{\AA}$

$c = 11.7595(14)\ \text{\AA}$

$\beta = 127.193(1)^\circ$

$V = 1486.8(2)\ \text{\AA}^3$

$Z = 4$

$F(000) = 680$

$D_x = 1.469\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 6987 reflections

$\theta = 2.7\text{--}28.3^\circ$

$\mu = 1.65\ \text{mm}^{-1}$

$T = 125\ \text{K}$

Plate, yellow

$0.23 \times 0.13 \times 0.06\ \text{mm}$

Data collection

Bruker APEXII CCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 φ and ω scans
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.703$, $T_{\max} = 0.908$

9483 measured reflections
1837 independent reflections
1710 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$
 $\theta_{\max} = 28.3^\circ$, $\theta_{\min} = 2.7^\circ$
 $h = -23 \rightarrow 23$
 $k = -12 \rightarrow 12$
 $l = -15 \rightarrow 15$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.022$
 $wR(F^2) = 0.061$
 $S = 1.09$
1837 reflections
98 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0347P)^2 + 0.7204P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.41 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Zn	0.0000	0.27994 (2)	0.2500	0.02138 (9)
N1	0.09270 (8)	0.19322 (13)	0.18122 (12)	0.0217 (2)
N2	0.0000	0.03178 (17)	0.2500	0.0181 (3)
C1	0.11086 (11)	0.36835 (17)	0.43838 (16)	0.0282 (3)
H1A	0.1292	0.4625	0.4207	0.042*
H1B	0.1660	0.3013	0.4858	0.042*
H1C	0.0909	0.3840	0.4996	0.042*
C2	0.13813 (11)	0.28148 (17)	0.14866 (16)	0.0260 (3)
H2A	0.1337	0.3846	0.1557	0.031*
C3	0.19141 (11)	0.22995 (19)	0.10513 (17)	0.0291 (3)
H3A	0.2237	0.2959	0.0845	0.035*
C4	0.19631 (10)	0.0801 (2)	0.09259 (16)	0.0298 (3)
H4A	0.2317	0.0413	0.0621	0.036*
C5	0.14921 (10)	-0.01296 (17)	0.12488 (15)	0.0261 (3)
H5A	0.1513	-0.1163	0.1160	0.031*
C6	0.09862 (9)	0.04762 (15)	0.17079 (13)	0.0197 (3)
C7	0.04737 (9)	-0.04323 (15)	0.21108 (13)	0.0188 (3)
C8	0.04827 (10)	-0.19665 (15)	0.20904 (15)	0.0239 (3)
H8A	0.0814	-0.2475	0.1801	0.029*
C9	0.0000	-0.2732 (2)	0.2500	0.0260 (4)
H9A	0.0000	-0.3775	0.2500	0.031*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn	0.02557 (13)	0.01758 (13)	0.02424 (13)	0.000	0.01676 (11)	0.000

N1	0.0219 (5)	0.0236 (6)	0.0222 (6)	-0.0008 (4)	0.0147 (5)	-0.0004 (4)
N2	0.0182 (7)	0.0178 (7)	0.0176 (7)	0.000	0.0105 (6)	0.000
C1	0.0330 (7)	0.0234 (7)	0.0303 (7)	-0.0055 (6)	0.0202 (7)	-0.0032 (6)
C2	0.0259 (7)	0.0279 (7)	0.0261 (7)	-0.0032 (6)	0.0167 (6)	0.0009 (6)
C3	0.0230 (7)	0.0420 (9)	0.0243 (7)	-0.0018 (6)	0.0154 (6)	0.0048 (6)
C4	0.0236 (7)	0.0461 (9)	0.0247 (7)	0.0082 (6)	0.0173 (6)	0.0050 (6)
C5	0.0252 (7)	0.0299 (8)	0.0253 (7)	0.0066 (6)	0.0164 (6)	0.0020 (6)
C6	0.0172 (6)	0.0242 (7)	0.0161 (6)	0.0021 (5)	0.0093 (5)	0.0006 (5)
C7	0.0177 (6)	0.0195 (6)	0.0167 (6)	0.0016 (5)	0.0091 (5)	-0.0005 (5)
C8	0.0230 (7)	0.0208 (7)	0.0237 (7)	0.0031 (5)	0.0119 (6)	-0.0023 (5)
C9	0.0262 (10)	0.0171 (9)	0.0276 (10)	0.000	0.0126 (9)	0.000

Geometric parameters (Å, °)

Zn—C1 ⁱ	2.0282 (15)	C2—H2A	0.9500
Zn—C1	2.0282 (15)	C3—C4	1.381 (2)
Zn—N1	2.3381 (12)	C3—H3A	0.9500
Zn—N2	2.2603 (16)	C4—C5	1.383 (2)
Zn—N1 ⁱ	2.3382 (12)	C4—H4A	0.9500
N1—C2	1.3364 (19)	C5—C6	1.3958 (18)
N1—C6	1.3416 (18)	C5—H5A	0.9500
N2—C7 ⁱ	1.3470 (15)	C6—C7	1.4893 (18)
N2—C7	1.3470 (15)	C7—C8	1.3979 (19)
C1—H1A	0.9800	C8—C9	1.3838 (18)
C1—H1B	0.9800	C8—H8A	0.9500
C1—H1C	0.9800	C9—C8 ⁱ	1.3838 (18)
C2—C3	1.385 (2)	C9—H9A	0.9500
C1 ⁱ —Zn—C1	133.21 (9)	N1—C2—H2A	118.4
C1 ⁱ —Zn—N2	113.39 (5)	C3—C2—H2A	118.4
C1—Zn—N2	113.39 (5)	C4—C3—C2	118.22 (14)
C1 ⁱ —Zn—N1	99.05 (5)	C4—C3—H3A	120.9
C1—Zn—N1	96.37 (5)	C2—C3—H3A	120.9
N2—Zn—N1	70.26 (3)	C3—C4—C5	119.41 (14)
C1 ⁱ —Zn—N1 ⁱ	96.37 (5)	C3—C4—H4A	120.3
C1—Zn—N1 ⁱ	99.05 (5)	C5—C4—H4A	120.3
N2—Zn—N1 ⁱ	70.26 (3)	C4—C5—C6	118.83 (14)
N1—Zn—N1 ⁱ	140.52 (6)	C4—C5—H5A	120.6
C2—N1—C6	118.48 (12)	C6—C5—H5A	120.6
C2—N1—Zn	123.28 (10)	N1—C6—C5	121.84 (13)
C6—N1—Zn	118.22 (9)	N1—C6—C7	115.22 (11)
C7 ⁱ —N2—C7	119.05 (16)	C5—C6—C7	122.93 (13)
C7 ⁱ —N2—Zn	120.48 (8)	N2—C7—C8	121.89 (13)
C7—N2—Zn	120.47 (8)	N2—C7—C6	115.77 (12)
Zn—C1—H1A	109.5	C8—C7—C6	122.34 (12)
Zn—C1—H1B	109.5	C9—C8—C7	118.82 (14)
H1A—C1—H1B	109.5	C9—C8—H8A	120.6
Zn—C1—H1C	109.5	C7—C8—H8A	120.6

H1A—C1—H1C	109.5	C8—C9—C8 ⁱ	119.53 (19)
H1B—C1—H1C	109.5	C8—C9—H9A	120.2
N1—C2—C3	123.19 (14)	C8 ⁱ —C9—H9A	120.2
C1 ⁱ —Zn—N1—C2	68.69 (12)	C2—C3—C4—C5	-0.6 (2)
C1—Zn—N1—C2	-66.99 (12)	C3—C4—C5—C6	-0.6 (2)
N2—Zn—N1—C2	-179.60 (12)	C2—N1—C6—C5	-1.1 (2)
N1 ⁱ —Zn—N1—C2	-179.60 (12)	Zn—N1—C6—C5	177.37 (10)
C1 ⁱ —Zn—N1—C6	-109.70 (11)	C2—N1—C6—C7	179.08 (12)
C1—Zn—N1—C6	114.62 (11)	Zn—N1—C6—C7	-2.45 (15)
N2—Zn—N1—C6	2.01 (9)	C4—C5—C6—N1	1.5 (2)
N1 ⁱ —Zn—N1—C6	2.01 (9)	C4—C5—C6—C7	-178.65 (12)
C1 ⁱ —Zn—N2—C7 ⁱ	-89.74 (8)	C7 ⁱ —N2—C7—C8	0.42 (9)
C1—Zn—N2—C7 ⁱ	90.26 (8)	Zn—N2—C7—C8	-179.58 (9)
N1—Zn—N2—C7 ⁱ	178.72 (7)	C7 ⁱ —N2—C7—C6	-179.48 (12)
N1 ⁱ —Zn—N2—C7 ⁱ	-1.28 (7)	Zn—N2—C7—C6	0.52 (12)
C1 ⁱ —Zn—N2—C7	90.26 (8)	N1—C6—C7—N2	1.31 (16)
C1—Zn—N2—C7	-89.74 (8)	C5—C6—C7—N2	-178.51 (11)
N1—Zn—N2—C7	-1.28 (7)	N1—C6—C7—C8	-178.59 (13)
N1 ⁱ —Zn—N2—C7	178.72 (7)	C5—C6—C7—C8	1.6 (2)
C6—N1—C2—C3	-0.3 (2)	N2—C7—C8—C9	-0.83 (19)
Zn—N1—C2—C3	-178.64 (11)	C6—C7—C8—C9	179.07 (10)
N1—C2—C3—C4	1.1 (2)	C7—C8—C9—C8 ⁱ	0.39 (9)

Symmetry code: (i) $-x, y, -z+1/2$.