

Dimethyl 2,2'-[butane-1,4-diylbis(sulfanediy)]dibenzoate

Iván Brito,^{a*} Alejandro Cárdenas,^b Joselyn Albanez,^a Michael Bolte^c and Matías López-Rodríguez^d

^aDepartamento de Química, Facultad de Ciencias Básicas, Universidad de Antofagasta, Casilla 170, Antofagasta - Chile, ^bDepartamento de Física, Facultad de Ciencias Básicas, Universidad de Antofagasta, Casilla 170, Antofagasta-Chile, ^cInstitut für Anorganische Chemie der Goethe-Universität Frankfurt, Max-von-Laue-Str. 7, D-60438 Frankfurt am Main, Germany, and ^dInstituto de Bio-Orgánica 'Antonio González', Universidad de La Laguna, Astrofísico Francisco, Sánchez N°2, La Laguna, Tenerife, Spain.

Correspondence e-mail: ivanbritob@yahoo.com


Received 14 October 2009; accepted 18 October 2009

Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.039; wR factor = 0.065; data-to-parameter ratio = 14.4.

The title compound, $\text{C}_{20}\text{H}_{22}\text{O}_4\text{S}_2$, was synthesized by the reaction of 1,4-dibromobutene with methyl thiosalicylate. The aliphatic segment of this ligand is in an all-*trans* conformation. The bridging chain, $-\text{S}-(\text{CH}_2)_4-\text{S}-$, is almost planar (r.m.s. deviation for all non-H atoms: 0.056 Å) and its mean plane forms dihedral angles of 16.60 (7) and 5.80 (2)° with the aromatic rings. In the crystal, the molecules are linked by weak $\text{C}-\text{H}\cdots\text{O}$ interactions into chains with graph-set notation $C(14)$ along $[0\ 0\ 1]$. The crystal studied was a racemic twin, the ratio of the twin components being 0.27 (9):0.73 (9).

Related literature

For the potential of coordination polymers based on multi-topic bridging ligands and metal centers as functional materials, see: Guo *et al.* (2002); Melcer *et al.* (2001). For the use of flexible ligands in such structures, see: Bu *et al.* (2001); Withersby *et al.* (1997). For our studies on the synthesis and structural characterization of divalent sulfur compounds, see: Brito *et al.* (2004, 2005, 2006). For a related compound, see: Awaleh *et al.* (2005). For graph-set notation of hydrogen bonds, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{20}\text{H}_{22}\text{O}_4\text{S}_2$
 $M_r = 390.5$
 Orthorhombic, $P2_12_12_1$
 $a = 7.4027$ (7) Å
 $b = 14.2976$ (11) Å
 $c = 17.7396$ (13) Å

$V = 1877.6$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.31$ mm⁻¹
 $T = 173$ K
 $0.28 \times 0.25 \times 0.23$ mm

Data collection

STOE IPDS II two-circle-diffractometer
 Absorption correction: multi-scan (MULABS; Spek, 2003; Blessing, 1995)
 $T_{\min} = 0.919$, $T_{\max} = 0.933$

8050 measured reflections
 3437 independent reflections
 2244 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.069$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.065$
 $S = 0.73$
 3437 reflections
 239 parameters
 H-atom parameters constrained

$\Delta\rho_{\max} = 0.16$ e Å⁻³
 $\Delta\rho_{\min} = -0.27$ e Å⁻³
 Absolute structure: Flack (1983),
 1430 Friedel pairs
 Flack parameter: 0.27 (9)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C}15-\text{H}15\cdots\text{O}3^i$	0.95	2.45	3.123 (5)	127

Symmetry code: (i) $-x + \frac{1}{2}, -y, z - \frac{1}{2}$.

Data collection: X-Area (Stoe & Cie, 2001); cell refinement: X-Area; data reduction: X-Area; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XP (Sheldrick, 2008) and PLATON (Spek, 2003); software used to prepare material for publication: WinGX (Farrugia, 1999).

This work was supported by a grant from the Universidad de Antofagasta (DI-1324-06). We thank the Spanish Research Council (CSIC) for providing us with a free-of-charge licence for the CSD system. JA thanks the Universidad de Antofagasta for PhD fellowships.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2290).

References

- Awaleh, M. O., Badia, A. & Brisse, F. (2005). *Acta Cryst.* **E61**, o2476–o2478.
- Bernstein, J., Davis, R. E., Shimon, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
- Brito, I., López-Rodríguez, M., Vargas, D. & León, Y. (2006). *Acta Cryst.* **E62**, o914–o916.
- Brito, I., Vargas, D., León, Y., Cárdenas, A., López-Rodríguez, M. & Wittke, O. (2004). *Acta Cryst.* **E60**, o1668–o1670.
- Brito, I., Vargas, D., Reyes, A., Cárdenas, A. & López-Rodríguez, M. (2005). *Acta Cryst.* **C61**, o234–o236.
- Bu, X. H., Chen, W., Lu, S. L., Zhang, R. H., Liao, D. Z., Bu, W. M., Shionoya, M., Brisse, F. & Ribas, J. (2001). *Angew. Chem. Int. Ed.* **40**, 3201–3203.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Guo, D., Pang, K. L., Duan, C. Y., He, C. & Meng, Q. (2002). *Inorg. Chem.* **41**, 5978–5985.
- Melcer, N. J., Enright, G. D., Ripmeester, J. A. & Shimizu, G. K. H. (2001). *Inorg. Chem.* **40**, 4641–4648.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Stoe & Cie (2001). *X-AREA* and *X-RED*. Stoe & Cie, Darmstadt, Germany.
- Withersby, M. A., Blake, A. J., Champness, N. R., Hubberstey, P., Li, W. S. & Schröder, M. (1997). *Angew. Chem. Int. Ed. Engl.* **36**, 2327–2329.

supporting information

Acta Cryst. (2009). E65, o2828–o2829 [https://doi.org/10.1107/S1600536809042846]

Dimethyl 2,2'-[butane-1,4-diylbis(sulfanediy)]dibenzoate

Iván Brito, Alejandro Cárdenas, Joselyn Albanez, Michael Bolte and Matías López-Rodríguez

S1. Comment

In recent years, the rational design of coordination polymers based on multitopic bridging ligands and metal centers represents one of the most rapidly developing fields owing to their potential as functional materials (Guo *et al.*, 2002; Melcer *et al.*, 2001). The use of flexible ligands in such studies has attracted increasing attention because the flexibility and conformational freedom of such ligands offer the possibility for the construction of diverse frameworks with tailored properties and functions (Bu *et al.*, 2001; Withersby *et al.*, 1997).

The structure of the title compound is described here as part of our work involving the study of the synthesis and structural characterization of divalent-sulfur compounds (Brito *et al.*, 2004, 2005, 2006). The title compound is a longer analogue of 2,2'-dithiodibenzoate with the two benzoate units interconnected by a flexible $-\text{S}-(\text{CH}_2)_4-\text{S}-$ bridge. The bridging chain moiety, $-\text{S}-(\text{CH}_2)_4-\text{S}-$ is almost planar (r.m.s. deviation for all non-H atoms: 0.056 Å). Its mean plane forms a dihedral angle of 16.60 (7) and 5.80 (2)° with the aromatic rings. The C(*sp*²)-S bond lengths [1.780 (3), 1.773 (3) Å] are significantly shorter than the C(*sp*³)-S [1.821 (3), 1.822 (3) Å] bond lengths due to *p*- π conjugation, similar to that observed in 1,6-Bis(phenylsulfanyl)hexane (Awaleh *et al.*, 2005). The torsion angles in the aliphatic segment of the title compound are all *trans*, indicating that the molecule is in the fully extended conformation. The supramolecular structure of the title compound depends solely upon C—H \cdots O hydrogen bonds: there are no significant S \cdots S nor S \cdots O contacts present in the structure and C—H \cdots π (arene) hydrogen bonds and aromatic $\pi\cdots\pi$ stacking interactions are also absent. Atom C15 in the molecule at (*x*,*y*,*z*) acts as a hydrogen-bond donor to the carbonyl O3 atom in the molecule at ($-x + 1/2$, $-y$, $z - 1/2$), thereby generating a C(14) chain (Bernstein *et al.*, 1995) running in the [0 0 1] direction (Figure 2). The molecular structure is stabilized by two C—H \cdots O intramolecular hydrogen bonds (Table 1).

S2. Experimental

The title compound was synthesized as follows: a solution of 1,4-dibromobutene (1.08 g, 5 mmol) in ethanol (10 ml) was added dropwise to a mixture of methyl thiosalicylate (1.85 g, 11 mmol), KOH (0.617 g, 11 mmol) and ethanol (10 ml). The reaction mixture was stirred for 24 h at room temperature. The precipitate was filtered off, washed with water. Yield 80%; m.p. 393–396 K. FT—IR (KBr pellets, cm⁻¹): ν (w, C—H of CH₃ (*asym.*)) 2984, ν (w, C—H of CH₃ (*sym.*)) 2842, ν (w, C—H (aliphatic chain, *sym.*)) 2947, ν (w, C—H (chain aliphatic, *asym.*)) 2919, ν (s, C=O) 1703, ν (s, C—H disubstitution 1,2) 1429, ν (s, O—C(CH₃)) 1252, ν (w, C—S) 749. Analysis calculated for C₂₀H₂₂O₄S₂, C 61.51, H 5.68, S 16.42%; found C 61.47, H, 5.72, S 15.32. Crystals suitable for single-crystals X-ray analysis were obtained by recrystallization from an acetonitrile solution.

S3. Refinement

Hydrogen atoms were located in a difference Fourier map but they were included in calculated positions [C—H = 0.95 - 0.99 Å] and refined as riding [$U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{C}_{\text{methyl}})$]. The crystal was refined as a racemic twin with a

ratio of the twin components of 0.27 (9)/0.73 (9).




Figure 1

A view of the molecular structure of the title compound, with the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level and H atoms are shown as small spheres of arbitrary radii.




Figure 2

Part of the crystal structure of the title compound viewed along the a axis, showing the formation of a C(14) chain along the $[0\ 0\ 1]$ direction. [Symmetry code for O3: $1/2 - x, -y, -1/2 + z$.]

Dimethyl 2,2'-[butane-1,4-diylbis(sulfanediyl)]dibenzoate

Crystal data

$C_{20}H_{22}O_4S_2$

$M_r = 390.5$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 7.4027$ (7) Å

$b = 14.2976$ (11) Å

$c = 17.7396$ (13) Å

$V = 1877.6$ (3) Å³

$Z = 4$

$F(000) = 824$

$D_x = 1.381$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 4482 reflections

$\theta = 3.4\text{--}25.9^\circ$

$\mu = 0.31$ mm⁻¹

$T = 173$ K

Block, colourless

$0.28 \times 0.25 \times 0.23$ mm

*Data collection*STOE IPDS II two-circle-
diffractometer

Graphite monochromator

 ω scansAbsorption correction: multi-scan
(*MULABS*; Spek, 2003; Blessing, 1995) $T_{\min} = 0.919$, $T_{\max} = 0.933$

8050 measured reflections

3437 independent reflections

2244 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.069$ $\theta_{\max} = 25.6^\circ$, $\theta_{\min} = 3.3^\circ$ $h = -8 \rightarrow 7$ $k = -17 \rightarrow 13$ $l = -21 \rightarrow 20$ *Refinement*Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.039$ $wR(F^2) = 0.065$ $S = 0.73$

3437 reflections

239 parameters

0 restraints

Primary atom site location: structure-invariant
direct methodsSecondary atom site location: difference Fourier
mapHydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0001P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} < 0.001$ $\Delta\rho_{\max} = 0.16 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$ Extinction correction: *SHELXL*

Extinction coefficient: 0.0030 (3)

Absolute structure: Flack (1983), 1430 Friedel
pairs

Absolute structure parameter: 0.27 (9)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.26565 (13)	0.29690 (6)	0.42902 (3)	0.0257 (2)
S2	0.30831 (13)	0.20540 (6)	0.80343 (4)	0.0292 (2)
O1	0.2673 (5)	0.41362 (15)	0.30785 (10)	0.0449 (8)
O2	0.2727 (4)	0.37231 (15)	0.18638 (9)	0.0340 (6)
O3	0.3720 (4)	0.08601 (17)	0.91935 (11)	0.0359 (7)
O4	0.4816 (4)	0.12269 (18)	1.03289 (10)	0.0369 (7)
C1	0.2650 (6)	0.2191 (2)	0.51027 (13)	0.0249 (8)
H1A	0.1554	0.1794	0.5099	0.03*
H1B	0.3725	0.1779	0.5092	0.03*
C2	0.2678 (6)	0.2809 (2)	0.58124 (13)	0.0264 (8)
H2A	0.3754	0.322	0.5799	0.032*
H2B	0.1591	0.3211	0.5819	0.032*
C3	0.2725 (6)	0.2212 (2)	0.65279 (13)	0.0261 (9)
H3A	0.1607	0.1834	0.6561	0.031*

H3B	0.3766	0.1779	0.6506	0.031*
C4	0.2885 (6)	0.2832 (2)	0.72250 (13)	0.0280 (9)
H4A	0.1803	0.3234	0.7276	0.034*
H4B	0.3964	0.3238	0.7187	0.034*
C11	0.2723 (5)	0.2174 (2)	0.35190 (13)	0.0232 (9)
C12	0.2798 (6)	0.2517 (2)	0.27679 (15)	0.0244 (9)
C13	0.2887 (5)	0.1872 (2)	0.21715 (14)	0.0288 (9)
H13	0.2957	0.2098	0.1669	0.035*
C14	0.2875 (6)	0.0934 (2)	0.22956 (15)	0.0350 (10)
H14	0.2954	0.0511	0.1883	0.042*
C15	0.2747 (6)	0.0593 (2)	0.30303 (15)	0.0339 (9)
H15	0.2696	-0.0061	0.3121	0.041*
C16	0.2695 (6)	0.1219 (2)	0.36288 (14)	0.0282 (9)
H16	0.2638	0.0983	0.4129	0.034*
C17	0.2741 (6)	0.3527 (2)	0.26050 (14)	0.0276 (9)
C18	0.2495 (7)	0.4698 (2)	0.16715 (15)	0.0389 (10)
H18A	0.3472	0.5066	0.1897	0.058*
H18B	0.1331	0.4919	0.1864	0.058*
H18C	0.2525	0.477	0.1122	0.058*
C21	0.3729 (5)	0.2822 (3)	0.87713 (15)	0.0249 (9)
C22	0.4253 (6)	0.2465 (3)	0.94829 (16)	0.0266 (9)
C23	0.4802 (5)	0.3083 (3)	1.00471 (16)	0.0308 (9)
H23	0.5158	0.2841	1.0523	0.037*
C24	0.4840 (6)	0.4036 (3)	0.99282 (17)	0.0349 (10)
H24	0.5217	0.4446	1.0319	0.042*
C25	0.4321 (5)	0.4390 (2)	0.92306 (17)	0.0319 (9)
H25	0.435	0.5045	0.914	0.038*
C26	0.3758 (6)	0.3778 (3)	0.86655 (16)	0.0294 (10)
H26	0.3385	0.4027	0.8194	0.035*
C27	0.4227 (5)	0.1458 (3)	0.96319 (16)	0.0287 (9)
C28	0.4834 (6)	0.0230 (3)	1.04913 (17)	0.0392 (11)
H28A	0.5551	-0.0095	1.0108	0.059*
H28B	0.3594	-0.001	1.0485	0.059*
H28C	0.5368	0.0125	1.099	0.059*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0329 (6)	0.0275 (4)	0.0166 (3)	-0.0011 (5)	-0.0012 (4)	-0.0023 (3)
S2	0.0443 (6)	0.0259 (5)	0.0175 (3)	-0.0023 (5)	-0.0008 (3)	0.0001 (4)
O1	0.084 (3)	0.0299 (14)	0.0212 (9)	-0.0002 (16)	-0.0042 (15)	-0.0037 (9)
O2	0.0500 (19)	0.0330 (13)	0.0189 (9)	0.0006 (14)	0.0028 (12)	0.0030 (9)
O3	0.053 (2)	0.0289 (15)	0.0253 (11)	-0.0018 (13)	-0.0031 (12)	-0.0013 (10)
O4	0.047 (2)	0.0387 (16)	0.0253 (10)	0.0049 (14)	-0.0123 (11)	0.0070 (11)
C1	0.028 (2)	0.0275 (19)	0.0195 (12)	-0.0018 (19)	0.0011 (14)	0.0010 (12)
C2	0.033 (2)	0.027 (2)	0.0193 (12)	0.0011 (19)	-0.0009 (16)	-0.0032 (12)
C3	0.029 (3)	0.027 (2)	0.0224 (14)	-0.002 (2)	-0.0004 (14)	0.0007 (11)
C4	0.035 (2)	0.031 (2)	0.0182 (12)	0.005 (2)	-0.0012 (14)	0.0028 (13)

C11	0.019 (2)	0.032 (2)	0.0186 (13)	0.001 (2)	-0.0006 (14)	-0.0067 (12)
C12	0.026 (2)	0.0257 (18)	0.0211 (13)	0.0007 (18)	0.0009 (15)	-0.0021 (12)
C13	0.033 (3)	0.034 (2)	0.0190 (12)	0.001 (2)	0.0014 (14)	-0.0066 (13)
C14	0.045 (3)	0.034 (2)	0.0265 (14)	0.002 (2)	-0.0003 (18)	-0.0135 (14)
C15	0.039 (3)	0.0284 (18)	0.0339 (14)	0.0011 (19)	0.0039 (19)	-0.0061 (13)
C16	0.039 (3)	0.0237 (19)	0.0223 (13)	0.000 (2)	-0.0014 (16)	0.0005 (12)
C17	0.028 (3)	0.034 (2)	0.0207 (14)	-0.003 (2)	-0.0004 (16)	0.0007 (13)
C18	0.049 (3)	0.038 (2)	0.0294 (14)	-0.002 (2)	-0.0001 (18)	0.0080 (13)
C21	0.029 (2)	0.027 (2)	0.0190 (13)	0.0006 (18)	0.0024 (13)	-0.0059 (14)
C22	0.026 (3)	0.030 (2)	0.0235 (16)	0.0050 (17)	0.0031 (14)	0.0010 (13)
C23	0.028 (2)	0.036 (2)	0.0275 (15)	0.0040 (18)	-0.0051 (15)	-0.0031 (17)
C24	0.039 (3)	0.036 (2)	0.0298 (16)	-0.0001 (19)	-0.0028 (17)	-0.0144 (17)
C25	0.035 (3)	0.026 (2)	0.0343 (16)	-0.0007 (17)	0.0052 (19)	-0.0028 (16)
C26	0.038 (3)	0.029 (2)	0.0214 (14)	-0.0001 (19)	0.0012 (15)	0.0031 (15)
C27	0.030 (3)	0.036 (2)	0.0202 (14)	0.005 (2)	0.0016 (15)	0.0023 (15)
C28	0.050 (3)	0.035 (2)	0.0326 (18)	0.005 (2)	-0.0053 (17)	0.0124 (15)

Geometric parameters (Å, °)

S1—C11	1.780 (3)	C13—C14	1.360 (5)
S1—C1	1.821 (3)	C13—H13	0.95
S2—C21	1.773 (3)	C14—C15	1.395 (4)
S2—C4	1.822 (3)	C14—H14	0.95
O1—C17	1.211 (4)	C15—C16	1.389 (4)
O2—C17	1.344 (3)	C15—H15	0.95
O2—C18	1.445 (4)	C16—H16	0.95
O3—C27	1.215 (4)	C18—H18A	0.98
O4—C27	1.352 (4)	C18—H18B	0.98
O4—C28	1.454 (4)	C18—H18C	0.98
C1—C2	1.538 (4)	C21—C26	1.380 (5)
C1—H1A	0.99	C21—C22	1.416 (4)
C1—H1B	0.99	C22—C23	1.395 (4)
C2—C3	1.529 (3)	C22—C27	1.465 (5)
C2—H2A	0.99	C23—C24	1.379 (5)
C2—H2B	0.99	C23—H23	0.95
C3—C4	1.526 (4)	C24—C25	1.392 (4)
C3—H3A	0.99	C24—H24	0.95
C3—H3B	0.99	C25—C26	1.395 (5)
C4—H4A	0.99	C25—H25	0.95
C4—H4B	0.99	C26—H26	0.95
C11—C16	1.379 (5)	C28—H28A	0.98
C11—C12	1.421 (4)	C28—H28B	0.98
C12—C13	1.404 (4)	C28—H28C	0.98
C12—C17	1.474 (4)		
C11—S1—C1	102.59 (14)	C16—C15—H15	120.3
C21—S2—C4	102.96 (15)	C14—C15—H15	120.3
C17—O2—C18	115.6 (2)	C11—C16—C15	122.0 (3)

C27—O4—C28	115.1 (3)	C11—C16—H16	119
C2—C1—S1	107.3 (2)	C15—C16—H16	119
C2—C1—H1A	110.3	O1—C17—O2	121.9 (3)
S1—C1—H1A	110.3	O1—C17—C12	124.8 (2)
C2—C1—H1B	110.3	O2—C17—C12	113.3 (3)
S1—C1—H1B	110.3	O2—C18—H18A	109.5
H1A—C1—H1B	108.5	O2—C18—H18B	109.5
C3—C2—C1	111.1 (2)	H18A—C18—H18B	109.5
C3—C2—H2A	109.4	O2—C18—H18C	109.5
C1—C2—H2A	109.4	H18A—C18—H18C	109.5
C3—C2—H2B	109.4	H18B—C18—H18C	109.5
C1—C2—H2B	109.4	C26—C21—C22	118.3 (3)
H2A—C2—H2B	108	C26—C21—S2	121.2 (2)
C4—C3—C2	110.5 (2)	C22—C21—S2	120.5 (3)
C4—C3—H3A	109.5	C23—C22—C21	119.4 (3)
C2—C3—H3A	109.5	C23—C22—C27	119.8 (3)
C4—C3—H3B	109.5	C21—C22—C27	120.8 (3)
C2—C3—H3B	109.5	C24—C23—C22	121.4 (3)
H3A—C3—H3B	108.1	C24—C23—H23	119.3
C3—C4—S2	106.9 (2)	C22—C23—H23	119.3
C3—C4—H4A	110.3	C23—C24—C25	119.4 (3)
S2—C4—H4A	110.3	C23—C24—H24	120.3
C3—C4—H4B	110.3	C25—C24—H24	120.3
S2—C4—H4B	110.3	C24—C25—C26	119.5 (3)
H4A—C4—H4B	108.6	C24—C25—H25	120.2
C16—C11—C12	118.3 (3)	C26—C25—H25	120.2
C16—C11—S1	121.6 (2)	C21—C26—C25	121.9 (3)
C12—C11—S1	120.1 (3)	C21—C26—H26	119
C13—C12—C11	118.8 (3)	C25—C26—H26	119
C13—C12—C17	119.8 (3)	O3—C27—O4	120.9 (3)
C11—C12—C17	121.4 (3)	O3—C27—C22	125.5 (3)
C14—C13—C12	121.7 (3)	O4—C27—C22	113.6 (3)
C14—C13—H13	119.2	O4—C28—H28A	109.5
C12—C13—H13	119.2	O4—C28—H28B	109.5
C13—C14—C15	119.8 (3)	H28A—C28—H28B	109.5
C13—C14—H14	120.1	O4—C28—H28C	109.5
C15—C14—H14	120.1	H28A—C28—H28C	109.5
C16—C15—C14	119.4 (3)	H28B—C28—H28C	109.5
C11—S1—C1—C2	177.5 (3)	C13—C12—C17—O2	2.0 (6)
S1—C1—C2—C3	-178.5 (3)	C11—C12—C17—O2	-176.6 (4)
C1—C2—C3—C4	176.1 (4)	C4—S2—C21—C26	7.4 (4)
C2—C3—C4—S2	-176.2 (3)	C4—S2—C21—C22	-171.7 (3)
C21—S2—C4—C3	168.3 (3)	C26—C21—C22—C23	-0.9 (6)
C1—S1—C11—C16	2.2 (4)	S2—C21—C22—C23	178.2 (3)
C1—S1—C11—C12	-178.1 (4)	C26—C21—C22—C27	179.2 (4)
C16—C11—C12—C13	-1.6 (7)	S2—C21—C22—C27	-1.6 (5)
S1—C11—C12—C13	178.7 (3)	C21—C22—C23—C24	0.3 (6)

C16—C11—C12—C17	177.0 (4)	C27—C22—C23—C24	-179.9 (4)
S1—C11—C12—C17	-2.7 (6)	C22—C23—C24—C25	-0.1 (6)
C11—C12—C13—C14	1.0 (7)	C23—C24—C25—C26	0.5 (6)
C17—C12—C13—C14	-177.6 (4)	C22—C21—C26—C25	1.3 (6)
C12—C13—C14—C15	0.9 (7)	S2—C21—C26—C25	-177.8 (3)
C13—C14—C15—C16	-2.2 (7)	C24—C25—C26—C21	-1.2 (6)
C12—C11—C16—C15	0.3 (7)	C28—O4—C27—O3	1.6 (5)
S1—C11—C16—C15	-180.0 (3)	C28—O4—C27—C22	-179.1 (3)
C14—C15—C16—C11	1.6 (7)	C23—C22—C27—O3	177.5 (3)
C18—O2—C17—O1	-4.3 (6)	C21—C22—C27—O3	-2.7 (7)
C18—O2—C17—C12	174.2 (4)	C23—C22—C27—O4	-1.8 (6)
C13—C12—C17—O1	-179.6 (4)	C21—C22—C27—O4	178.1 (3)
C11—C12—C17—O1	1.8 (7)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C13—H13 \cdots O2	0.95	2.36	2.705 (4)	101
C23—H23 \cdots O4	0.95	2.35	2.701 (5)	101
C15—H15 \cdots O3 ⁱ	0.95	2.45	3.123 (5)	127

Symmetry code: (i) $-x+1/2, -y, z-1/2$.