

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Poly[(6-carboxypicolinato- $\kappa^3\text{O}^2, \text{N}, \text{O}^6$)-(μ_3 -pyridine-2,6-dicarboxylato- $\kappa^5\text{O}^2, \text{N}, \text{O}^6: \text{O}^{2'}: \text{O}^{6'}$)dysprosium(III)]

Xu Li,^a Qing-Yang Lian,^a Qiu-Hui Meng,^a Yi-Fan Luo^{a*} and Rong-Hua Zeng^{a,b}

^aSchool of Chemistry and Environment, South China Normal University, Guangzhou 510006, People's Republic of China., and ^bSouth China Normal University, Key Laboratory of Technology of Electrochemical Energy Storage and Power Generation in Guangdong Universities, Guangzhou 510006, People's Republic of China
Correspondence e-mail: luoyf2004@yahoo.com.cn


Received 15 September 2009; accepted 26 September 2009

Key indicators: single-crystal X-ray study; $T = 296$ K; mean $\sigma(\text{C}-\text{C}) = 0.005$ Å; R factor = 0.021; wR factor = 0.051; data-to-parameter ratio = 10.5.

In the title complex, $[\text{Dy}(\text{C}_7\text{H}_3\text{NO}_4)(\text{C}_7\text{H}_4\text{NO}_4)]_n$, one of the ligands is fully deprotonated while the second has lost only one H atom. Each Dy^{III} ion is coordinated by six O atoms and two N atoms from two pyridine-2,6-dicarboxylate and two 6-carboxypicolinate ligands, displaying a bicapped trigonal-prismatic geometry. The average Dy—O bond distance is 2.40 Å, some 0.1 Å longer than the corresponding Ho—O distance in the isotopic holmium complex. Adjacent Dy^{III} ions are linked by the pyridine-2,6-dicarboxylate ligands, forming a layer in (100). These layers are further connected by π - π stacking interactions between neighboring pyridyl rings [centroid-centroid distance = 3.827 (3) Å] and C—H \cdots O hydrogen-bonding interactions, assembling a three-dimensional supramolecular network. Within each layer, there are other π - π stacking interactions between neighboring pyridyl rings [centroid-centroid distance = 3.501 (2) Å] and O—H \cdots O and C—H \cdots O hydrogen-bonding interactions, which further stabilize the structure.

Related literature

For the isotopic holmium analogue, see: Fernandes *et al.* (2001). For other related structures, see: Hong (2007); Huang *et al.* (2008); Idrees *et al.* (2009); Rafizadeh & Amani (2006); Thallapally *et al.* (2008).



Experimental

Crystal data

$[\text{Dy}(\text{C}_7\text{H}_3\text{NO}_4)(\text{C}_7\text{H}_4\text{NO}_4)]$
 $M_r = 493.72$
 Monoclinic, $P2_1/c$
 $a = 12.2151$ (14) Å
 $b = 8.3703$ (10) Å
 $c = 13.4698$ (16) Å
 $\beta = 102.332$ (1)°

$V = 1345.4$ (3) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 5.61$ mm⁻¹
 $T = 296$ K
 $0.23 \times 0.21 \times 0.19$ mm

Data collection

Bruker APEXII area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2004)
 $T_{\text{min}} = 0.359$, $T_{\text{max}} = 0.415$

6670 measured reflections
 2413 independent reflections
 2305 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.024$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$
 $wR(F^2) = 0.051$
 $S = 1.12$
 2413 reflections
 229 parameters
 1 restraint

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.58$ e Å⁻³
 $\Delta\rho_{\text{min}} = -1.43$ e Å⁻³

Table 1

Selected bond lengths (Å).

Dy1—O8 ⁱ	2.266 (2)	Dy1—O7	2.405 (2)
Dy1—O6 ⁱⁱ	2.314 (3)	Dy1—N2	2.469 (3)
Dy1—O1	2.328 (2)	Dy1—N1	2.488 (3)
Dy1—O5	2.385 (3)	Dy1—O4	2.513 (2)

Symmetry codes: (i) $-x, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $x, -y + \frac{1}{2}, z - \frac{1}{2}$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O2—H2A \cdots O4 ⁱⁱⁱ	0.811 (19)	1.73 (2)	2.518 (3)	164 (5)
O2—H2A \cdots O3 ⁱⁱⁱ	0.811 (19)	2.65 (3)	3.303 (4)	139 (4)
C9—H9 \cdots O1 ^{iv}	0.93	2.42	3.332 (4)	165
C1—H1 \cdots O5 ^v	0.93	2.42	3.123 (5)	133
C2—H2 \cdots O3 ⁱⁱⁱ	0.93	2.47	3.393 (5)	174

Symmetry code: (iii) $x, y + 1, z$; (iv) $x, -y + \frac{1}{2}, z + \frac{1}{2}$; (v) $-x + 1, -y + 1, -z + 2$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINTE* (Bruker, 2004); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP III* (Burnett & Johnson, 1996) and *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

The authors acknowledge the Chan Xue Yan Cooperative Special Project of Guangdong Province and the Ministry of Science and Technology of the People's Republic of China (Project No. 2007A090302046), the Project of Science and Technology of Guangdong Province (Project No. 2007A020200002-4) and the Natural Science Foundation of Guangdong Province (No. 9151063101000037) for supporting this work.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ2654).

References

- Bruker (2004). *APEX2*, *SAINTE* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burnett, M. N. & Johnson, C. K. (1996). *ORTEP III*. Report ORNL-6895. Oak Ridge National Laboratory, Tennessee, USA.
- Fernandes, A., Jaud, J., Dexpert-Ghys, J. & Brouca-Cabarrecq, C. (2001). *Polyhedron*, **20**, 2385–2391.
- Hong, M. C. (2007). *Cryst. Growth Des.* **7**, 10–14.
- Huang, Y.-G., Yuan, D.-Q., Gong, Y.-Q., Jiang, F.-L. & Hong, M.-C. (2008). *J. Mol. Struct.* **872**, 99–104.
- Idrees, M., Shadab, S. M. & Kumar, S. (2009). *Acta Cryst.* **E65**, o1492–o1493.
- Rafizadeh, M. & Amani, V. (2006). *Acta Cryst.* **E62**, m90–m91.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Thallapally, P. K., Tian, J., Kishan, M. R., Fernandez, C. A., Dalgarno, S. J., McGrail, P. B., Warren, J. E. & Atwood, J. L. (2008). *J. Am. Chem. Soc.* **130**, 16842–16843.

supporting information

Acta Cryst. (2009). E65, m1288–m1289 [https://doi.org/10.1107/S1600536809039075]

Poly[(6-carboxypicolinato- κ^3O^2,N,O^6)(μ_3 -pyridine-2,6-dicarboxylato- $\kappa^5O^2,N,O^6:O^2':O^6'$)dysprosium(III)]

Xu Li, Qing-Yang Lian, Qiu-Hui Meng, Yi-Fan Luo and Rong-Hua Zeng

S1. Comment

Research on the design and synthesis of metal-organic frameworks (MOFs) in recent years has become an active area in the fields of crystal engineering and supramolecular chemistry, not only because of their tremendous potential applications in gas storage, chemical separations, ion exchange, microelectronics, nonlinear optics, and heterogeneous catalysis, but also because of their intriguing variety of architectures and topologies (Hong, 2007; Thallapally *et al.*, 2008). The synthesis of such species is often based on the self-assembly of suitable building blocks to give supramolecular networks constructed by coordination or/and hydrogen bonds or other weaker supramolecular interactions, such as π - π stacking interactions. As a building block, the pyridine-2,6-dicarboxylic acid is a good ligand with multifunctional coordination sites providing intriguing architectures and topologies (Fernandes *et al.*, 2001; Rafizadeh & Amani, 2006; Huang *et al.*, 2008; Idrees *et al.*, 2009). Recently, we obtained the title coordination polymer, which was synthesized under hydrothermal conditions.

In the structure of the title compound (Fig. 1), one of the ligands is fully deprotonated while the second carries an OH group. Each Dy^{III} centre is eight -coordinated by six oxygen atoms two N atoms from two pyridine-2,6-dicarboxylato and two 6-carboxypicolinato ligands, and can be described as having a bicapped trigonal prismatic geometry with Dy—O distances and O—Dy—O angles ranging from 2.266 (2) Å to 2.513 (2) Å (Table 1) and 76.75.30 (8) °A to 153.60 (9) °A, respectively. It is of interest that the Dy \cdots O and Dy \cdots N distances are slightly longer than the corresponding values in the isostructural Ho complex (Fernandes *et al.*, 2001). Indeed the average Dy—O bond distance is 2.396 Å, some 0.1 Å longer than the corresponding distance (Ho—O = 2.385 Å) in the isomorphous holmium complex as anticipated due to the lanthanide contraction. The pyridine-2,6-dicarboxylato ligands act as bridges linking adjacent Dy^{III} metal centres into a layer parallel to the (1 0 0) plane. Within the layer, the 6-carboxypicolinato ligands are both hydrogen bond donors and hydrogen bond acceptors with O—H \cdots O and C—H \cdots O hydrogen bonding interactions stabilizing the crystal structure (Table 2). π - π stacking interactions (the centroid-centroid distance between neighboring pyridyl rings is 3.827 (3) Å) and C—H \cdots O hydrogen bonding interactions connect those layers to produce a three-dimensional supramolecular motif (Fig. 2). Other π - π stacking interactions between neighboring pyridyl rings are also present in each layer, the centroid-centroid distance is 3.501 (2) Å.

S2. Experimental

A mixture of Dy₂O₃ (0.375 g; 1 mmol), pyridine-2,6-dicarboxylic acid (0.167 g; 1 mmol), water (10 ml) and HNO₃ (0.024 g; 0.385 mmol) was stirred vigorously for 20 min and then sealed in a Teflon-lined stainless-steel autoclave (20 ml, capacity). The autoclave was heated and maintained at 433 K for 3 days, and then cooled to room temperature at 5 K h⁻¹ to yield colorless block-like crystals.

S3. Refinement

The H atom of the 6-carboxypicolinate ligand was located in a difference Fourier map and refined with a distance restraint of O–H = 0.82 Å, and with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$. H atoms attached to C were placed at calculated positions and were treated as riding on their parent atoms with C–H = 0.93 Å, and with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$.




Figure 1

The molecular structure showing the atomic-numbering scheme, with displacement ellipsoids drawn at the 30% probability level. Symmetry codes: (#1) $x, 0.5-y, -0.5+z$; (#2) $-x, 0.5+y, 1.5-z$.




Figure 2

A view of the three-dimensional supramolecular network. Hydrogen bonds are shown as dashed lines.

Poly[(6-carboxypicolinato- κ^3O^2,N,O^6)(μ_3 -pyridine-2,6-dicarboxylato- $\kappa^5O^2,N,O^6:O^2:O^6$)dysprosium(III)]

Crystal data

[Dy(C₇H₃NO₄)(C₇H₄NO₄)]

$M_r = 493.72$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 12.2151 (14) \text{ \AA}$

$b = 8.3703 (10) \text{ \AA}$

$c = 13.4698 (16) \text{ \AA}$

$\beta = 102.332 (1)^\circ$

$V = 1345.4 (3) \text{ \AA}^3$

$Z = 4$

$F(000) = 940$

$D_x = 2.437 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5626 reflections

$\theta = 2.4\text{--}27.8^\circ$

$\mu = 5.61 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Block, colourless

$0.23 \times 0.21 \times 0.19 \text{ mm}$

Data collection

Bruker APEXII area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2004)

$T_{\min} = 0.359$, $T_{\max} = 0.415$

6670 measured reflections

2413 independent reflections

2305 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.024$

$\theta_{\max} = 25.2^\circ$, $\theta_{\min} = 1.7^\circ$

$h = -14 \rightarrow 12$

$k = -8 \rightarrow 10$

$l = -16 \rightarrow 16$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.021$
 $wR(F^2) = 0.051$
 $S = 1.12$
 2413 reflections
 229 parameters
 1 restraint
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0241P)^2 + 1.92P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.002$
 $\Delta\rho_{\max} = 0.58 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -1.43 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Dy1	0.197015 (13)	0.232131 (18)	0.818183 (11)	0.00903 (7)
C2	0.5158 (3)	0.5574 (5)	0.8746 (3)	0.0243 (9)
H2	0.5247	0.6677	0.8750	0.029*
C3	0.4095 (3)	0.4901 (4)	0.8567 (3)	0.0184 (8)
C4	0.4836 (3)	0.2361 (4)	0.8735 (3)	0.0187 (8)
C1	0.6084 (3)	0.4572 (5)	0.8919 (3)	0.0282 (9)
H1	0.6805	0.4990	0.9030	0.034*
C5	0.5912 (3)	0.2933 (5)	0.8922 (3)	0.0274 (9)
H5	0.6518	0.2233	0.9049	0.033*
N1	0.3933 (2)	0.3321 (3)	0.8555 (2)	0.0136 (6)
C6	0.3019 (3)	0.5872 (4)	0.8367 (3)	0.0201 (8)
O1	0.2131 (2)	0.5091 (3)	0.82690 (18)	0.0180 (6)
O2	0.3085 (3)	0.7328 (3)	0.8293 (3)	0.0452 (10)
C7	0.4611 (3)	0.0608 (4)	0.8716 (3)	0.0203 (8)
O4	0.3508 (2)	0.0270 (3)	0.8468 (2)	0.0202 (6)
O3	0.5333 (2)	-0.0385 (3)	0.8889 (3)	0.0368 (8)
C13	0.2072 (3)	0.1845 (4)	1.0625 (2)	0.0128 (7)
C8	0.1581 (3)	0.0254 (4)	1.0222 (2)	0.0122 (7)
O5	0.2310 (2)	0.2782 (3)	0.9969 (2)	0.0200 (6)
N2	0.1355 (2)	0.0186 (3)	0.9205 (2)	0.0107 (6)
C11	0.0664 (3)	-0.2478 (4)	0.9275 (3)	0.0171 (8)
H11	0.0339	-0.3386	0.8937	0.021*
C12	0.0876 (3)	-0.1133 (4)	0.8745 (2)	0.0115 (7)
O7	0.1092 (2)	0.0071 (3)	0.72203 (17)	0.0148 (5)

O6	0.2198 (2)	0.2145 (3)	1.15526 (19)	0.0184 (6)
C14	0.0590 (3)	-0.0988 (4)	0.7607 (2)	0.0115 (7)
C9	0.1386 (3)	-0.1020 (4)	1.0812 (3)	0.0152 (7)
H9	0.1539	-0.0945	1.1517	0.018*
C10	0.0953 (3)	-0.2420 (4)	1.0324 (3)	0.0183 (8)
H10	0.0858	-0.3317	1.0703	0.022*
O8	-0.0153 (2)	-0.1898 (3)	0.71176 (18)	0.0171 (5)
H2A	0.334 (3)	0.822 (3)	0.838 (3)	0.026*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Dy1	0.00974 (11)	0.00762 (10)	0.01008 (11)	-0.00032 (5)	0.00291 (7)	0.00052 (5)
C2	0.026 (2)	0.0179 (19)	0.030 (2)	-0.0111 (16)	0.0069 (17)	-0.0045 (16)
C3	0.022 (2)	0.0131 (17)	0.0203 (19)	-0.0011 (15)	0.0048 (15)	0.0002 (14)
C4	0.015 (2)	0.0168 (18)	0.024 (2)	-0.0005 (14)	0.0039 (16)	-0.0011 (14)
C1	0.019 (2)	0.025 (2)	0.040 (3)	-0.0099 (16)	0.0056 (18)	-0.0064 (17)
C5	0.015 (2)	0.023 (2)	0.042 (3)	-0.0006 (16)	0.0021 (18)	-0.0035 (18)
N1	0.0122 (15)	0.0120 (14)	0.0171 (15)	-0.0013 (12)	0.0043 (12)	0.0005 (11)
C6	0.025 (2)	0.0145 (18)	0.0207 (19)	0.0017 (15)	0.0048 (16)	0.0005 (14)
O1	0.0183 (14)	0.0111 (13)	0.0251 (14)	-0.0001 (10)	0.0057 (11)	0.0010 (9)
O2	0.039 (2)	0.0065 (14)	0.088 (3)	-0.0048 (12)	0.009 (2)	0.0015 (15)
C7	0.016 (2)	0.0165 (18)	0.029 (2)	0.0006 (15)	0.0053 (16)	-0.0004 (15)
O4	0.0121 (13)	0.0105 (12)	0.0378 (16)	-0.0023 (10)	0.0048 (11)	0.0001 (10)
O3	0.0185 (15)	0.0197 (15)	0.070 (2)	0.0055 (12)	0.0043 (15)	0.0011 (14)
C13	0.0114 (17)	0.0152 (17)	0.0122 (18)	0.0025 (13)	0.0031 (14)	0.0011 (13)
C8	0.0097 (17)	0.0157 (17)	0.0124 (17)	0.0037 (13)	0.0049 (13)	-0.0021 (13)
O5	0.0295 (16)	0.0178 (13)	0.0137 (14)	-0.0094 (11)	0.0066 (12)	-0.0016 (10)
N2	0.0102 (14)	0.0123 (14)	0.0102 (14)	-0.0009 (11)	0.0036 (11)	-0.0004 (11)
C11	0.024 (2)	0.0125 (17)	0.017 (2)	-0.0039 (13)	0.0096 (17)	-0.0024 (13)
C12	0.0103 (17)	0.0112 (16)	0.0143 (17)	-0.0011 (13)	0.0053 (13)	0.0000 (13)
O7	0.0177 (13)	0.0146 (12)	0.0131 (12)	-0.0036 (10)	0.0054 (10)	-0.0004 (9)
O6	0.0262 (15)	0.0182 (13)	0.0119 (13)	-0.0017 (11)	0.0064 (11)	-0.0037 (10)
C14	0.0093 (17)	0.0120 (16)	0.0139 (17)	0.0019 (13)	0.0043 (13)	-0.0033 (13)
C9	0.0166 (18)	0.0191 (18)	0.0107 (17)	0.0001 (14)	0.0044 (14)	0.0015 (13)
C10	0.019 (2)	0.0191 (18)	0.018 (2)	-0.0012 (14)	0.0068 (16)	0.0061 (13)
O8	0.0159 (13)	0.0198 (13)	0.0163 (13)	-0.0061 (11)	0.0052 (10)	-0.0045 (10)

Geometric parameters (Å, °)

Dy1—O8 ⁱ	2.266 (2)	O2—H2A	0.811 (19)
Dy1—O6 ⁱⁱ	2.314 (3)	C7—O3	1.198 (5)
Dy1—O1	2.328 (2)	C7—O4	1.347 (4)
Dy1—O5	2.385 (3)	C13—O6	1.251 (4)
Dy1—O7	2.405 (2)	C13—O5	1.261 (4)
Dy1—N2	2.469 (3)	C13—C8	1.513 (5)
Dy1—N1	2.488 (3)	C8—N2	1.341 (4)
Dy1—O4	2.513 (2)	C8—C9	1.381 (5)

C2—C1	1.388 (6)	N2—C12	1.338 (4)
C2—C3	1.389 (6)	C11—C10	1.382 (6)
C2—H2	0.9300	C11—C12	1.387 (5)
C3—N1	1.336 (4)	C11—H11	0.9300
C3—C6	1.519 (5)	C12—C14	1.502 (5)
C4—N1	1.344 (5)	O7—C14	1.253 (4)
C4—C5	1.371 (6)	O6—Dy1 ⁱⁱⁱ	2.314 (3)
C4—C7	1.492 (5)	C14—O8	1.257 (4)
C1—C5	1.388 (6)	C9—C10	1.392 (5)
C1—H1	0.9300	C9—H9	0.9300
C5—H5	0.9300	C10—H10	0.9300
C6—O2	1.226 (4)	O8—Dy1 ^{iv}	2.266 (2)
C6—O1	1.250 (5)		
O8 ⁱ —Dy1—O6 ⁱⁱ	95.01 (9)	C1—C5—H5	120.5
O8 ⁱ —Dy1—O1	77.91 (9)	C3—N1—C4	118.4 (3)
O6 ⁱⁱ —Dy1—O1	80.18 (9)	C3—N1—Dy1	117.9 (2)
O8 ⁱ —Dy1—O5	94.91 (9)	C4—N1—Dy1	123.6 (2)
O6 ⁱⁱ —Dy1—O5	153.60 (9)	O2—C6—O1	125.5 (4)
O1—Dy1—O5	78.08 (8)	O2—C6—C3	118.5 (4)
O8 ⁱ —Dy1—O7	79.79 (9)	O1—C6—C3	115.9 (3)
O6 ⁱⁱ —Dy1—O7	76.74 (8)	C6—O1—Dy1	126.0 (2)
O1—Dy1—O7	146.15 (8)	C6—O2—H2A	159 (3)
O5—Dy1—O7	129.18 (8)	O3—C7—O4	124.0 (3)
O8 ⁱ —Dy1—N2	84.53 (9)	O3—C7—C4	123.6 (4)
O6 ⁱⁱ —Dy1—N2	141.40 (9)	O4—C7—C4	112.4 (3)
O1—Dy1—N2	136.47 (9)	C7—O4—Dy1	124.8 (2)
O5—Dy1—N2	64.02 (8)	O6—C13—O5	125.3 (3)
O7—Dy1—N2	65.15 (8)	O6—C13—C8	119.3 (3)
O8 ⁱ —Dy1—N1	143.55 (10)	O5—C13—C8	115.4 (3)
O6 ⁱⁱ —Dy1—N1	79.66 (9)	N2—C8—C9	122.2 (3)
O1—Dy1—N1	65.64 (9)	N2—C8—C13	112.5 (3)
O5—Dy1—N1	77.84 (10)	C9—C8—C13	125.3 (3)
O7—Dy1—N1	132.22 (9)	C13—O5—Dy1	126.2 (2)
N2—Dy1—N1	121.72 (9)	C12—N2—C8	118.8 (3)
O8 ⁱ —Dy1—O4	153.64 (9)	C12—N2—Dy1	119.6 (2)
O6 ⁱⁱ —Dy1—O4	92.28 (9)	C8—N2—Dy1	121.3 (2)
O1—Dy1—O4	128.35 (8)	C10—C11—C12	117.6 (3)
O5—Dy1—O4	89.58 (9)	C10—C11—H11	121.2
O7—Dy1—O4	77.30 (8)	C12—C11—H11	121.2
N2—Dy1—O4	74.15 (9)	N2—C12—C11	122.9 (3)
N1—Dy1—O4	62.74 (8)	N2—C12—C14	112.9 (3)
C1—C2—C3	118.9 (3)	C11—C12—C14	124.2 (3)
C1—C2—H2	120.6	C14—O7—Dy1	122.1 (2)
C3—C2—H2	120.6	C13—O6—Dy1 ⁱⁱⁱ	166.3 (2)
N1—C3—C2	122.2 (3)	O7—C14—O8	125.0 (3)
N1—C3—C6	114.1 (3)	O7—C14—C12	116.9 (3)
C2—C3—C6	123.7 (3)	O8—C14—C12	118.1 (3)

N1—C4—C5	122.8 (3)	C8—C9—C10	118.2 (3)
N1—C4—C7	116.3 (3)	C8—C9—H9	120.9
C5—C4—C7	120.8 (3)	C10—C9—H9	120.9
C2—C1—C5	118.7 (4)	C11—C10—C9	120.1 (3)
C2—C1—H1	120.7	C11—C10—H10	120.0
C5—C1—H1	120.7	C9—C10—H10	120.0
C4—C5—C1	119.0 (4)	C14—O8—Dy1 ^{iv}	146.8 (2)
C4—C5—H5	120.5		

Symmetry codes: (i) $-x, y+1/2, -z+3/2$; (ii) $x, -y+1/2, z-1/2$; (iii) $x, -y+1/2, z+1/2$; (iv) $-x, y-1/2, -z+3/2$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
O2—H2A...O4 ^v	0.81 (2)	1.73 (2)	2.518 (3)	164 (5)
O2—H2A...O3 ^v	0.81 (2)	2.65 (3)	3.303 (4)	139 (4)
C9—H9...O1 ⁱⁱⁱ	0.93	2.42	3.332 (4)	165
C1—H1...O5 ^{vi}	0.93	2.42	3.123 (5)	133
C2—H2...O3 ^v	0.93	2.47	3.393 (5)	174

Symmetry codes: (iii) $x, -y+1/2, z+1/2$; (v) $x, y+1, z$; (vi) $-x+1, -y+1, -z+2$.