

Dichloridooctakis(2-chlorobenzyl)di- μ_2 -hydroxido-di- μ_3 -oxido-tetratin(IV)

Qi-Jun Zhang, Han-Dong Yin,* Li-Yuan Wen and Da-Qi Wang

College of Chemistry and Chemical Engineering, Liaocheng University, Shandong 252059, People's Republic of China
Correspondence e-mail: handongyin@163.com


Received 21 October 2009; accepted 28 October 2009

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.009\text{ \AA}$; disorder in main residue; R factor = 0.029; wR factor = 0.074; data-to-parameter ratio = 15.1.

The title tetranuclear Sn^{IV} compound, $[\text{Sn}_4(\text{C}_7\text{H}_6\text{Cl})_8\text{Cl}_2\text{O}_2(\text{OH})_2]$, has site symmetry $\bar{1}$. Two O^{2-} and two OH^- anions bridge four Sn^{IV} cations to form the tetranuclear compound. The two independent Sn^{IV} cations assume SnO_3C_2 and $\text{SnO}_2\text{C}_2\text{Cl}$ distorted trigonal-bipyramidal coordination geometries. Intramolecular $\text{O}-\text{H}\cdots\text{Cl}$ hydrogen bonding is present in the structure. One Cl atom of a chlorobenzyl ligand is disordered over two sites with an occupancy ratio of 0.693 (2):0.307 (2).

Related literature

For a related structure, see: Li *et al.* (2006). For the corresponding bond distances in an organotin compound, see: Lo & Ng (2009).



Experimental

Crystal data

$[\text{Sn}_4(\text{C}_7\text{H}_6\text{Cl})_8\text{Cl}_2\text{O}_2(\text{OH})_2]$	$\gamma = 75.753 (2)^\circ$
$M_r = 1616.22$	$V = 1475.9 (6)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 1$
$a = 10.986 (2)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 11.227 (2)\text{ \AA}$	$\mu = 2.17\text{ mm}^{-1}$
$c = 13.573 (3)\text{ \AA}$	$T = 298\text{ K}$
$\alpha = 74.656 (2)^\circ$	$0.44 \times 0.37 \times 0.33\text{ mm}$
$\beta = 67.942 (2)^\circ$	

Data collection

Bruker SMART CCD area-detector diffractometer	7669 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Sheldrick, 1996)	5112 independent reflections
$T_{\min} = 0.449$, $T_{\max} = 0.535$	3865 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.015$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.029$	338 parameters
$wR(F^2) = 0.074$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\max} = 1.27\text{ e \AA}^{-3}$
5112 reflections	$\Delta\rho_{\min} = -0.57\text{ e \AA}^{-3}$

Table 1
Selected bond lengths (\AA).

$\text{Sn1}-\text{O1}$	2.148 (3)	$\text{Sn2}-\text{O1}$	2.276 (3)
$\text{Sn1}-\text{O2}$	2.050 (3)	$\text{Sn2}-\text{O2}$	2.025 (3)
$\text{Sn1}-\text{O2}^i$	2.146 (3)	$\text{Sn2}-\text{C15}$	2.147 (5)
$\text{Sn1}-\text{C1}$	2.126 (5)	$\text{Sn2}-\text{C22}$	2.149 (5)
$\text{Sn1}-\text{C8}$	2.146 (5)	$\text{Sn2}-\text{Cl1}$	2.4376 (13)

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Table 2
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O1}-\text{H1}\cdots\text{Cl2}$	0.86	2.80	3.386 (4)	127

Data collection: *SMART* (Siemens, 1996); cell refinement: *SAINT* (Siemens, 1996); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL*.

We acknowledge the National Natural Science Foundation of China (20771053) and the Natural Science Foundation of Shandong Province, China (Y2008B48) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2652).

References

- Li, G., Yin, H. & Wang, D. (2006). *Acta Cryst. E* **62**, m366–m368.
- Lo, K. M. & Ng, S. W. (2009). *Acta Cryst. E* **65**, m593.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siemens (1996). *SMART* and *SAINT*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

supporting information

Acta Cryst. (2009). E65, m1494 [https://doi.org/10.1107/S1600536809045176]

Dichloridooctakis(2-chlorobenzyl)di- μ_2 -hydroxido-di- μ_3 -oxido-tetratin(IV)

Qi-Jun Zhang, Han-Dong Yin, Li-Yuan Wen and Da-Qi Wang

S1. Comment


The title compound, (I), was obtained as an adventitious product of the partial hydrolysis of di(2-chlorobenzyl)-dichloridotin(IV) during the attempted preparation of adducts of this tin precursor complex with 5-chlorosalicylaldehyde benzoyldrazone in benzene and ethanol. It crystallizes from dichloromethane and ethanol. From Fig. 1, it can be seen that complex (I) contains two independent penta-coordinated Sn atoms. It is a centrosymmetric complex, where one half of the molecule comprises the crystallographic asymmetric unit and the other half is generated by an inversion centre. Each of the two independent Sn atoms is five-coordinate, adopting approximate trigonal bipyramidal coordination (Table 1). These are similar to those in the related organotin compound (Li *et al.*, 2006). The molecular conformation is stabilized by O1—H1 \cdots Cl2 hydrogen bond (Table 2). The Sn—C distances lie in the rather narrow range 2.126 (5)—2.149 (5) Å, which are closed to the corresponding distances reported in the organotin compound (Lo & Ng, 2009).

S2. Experimental

Di(2-chlorobenzyl)dichloridotin(IV) (2 mmol) and 5-chlorosalicylaldehyde benzoyldrazone (2 mmol) was added to a solution of sodium methoxide (3 mmol) in benzene (15 ml) and ethanol (15 ml, 95%). The mixture was then heated under reflux with stirring for 5 h and the solvent was removed by evaporation in *vacuo*. The crude adduct was recrystallized from dichloromethane/ethanol and colourless crystals suitable for X-ray diffraction were obtained.

S3. Refinement


The H atoms were positioned geometrically, with methylene C—H distances of 0.97 Å, aromatic C—H distances of 0.93 Å, O—H distances of 0.862 Å and refined as riding on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C}, \text{O})$. The Cl2 atom is disordered, the C2-phenyl part was refined as a rigid hexagon and the temperature factors of the carbon atoms were restrained to be nearly isotropic. The highest peak in the difference map is 1.21 Å apart from Cl1 atom.

**Figure 1**

The molecular structure of the compound showing 50% probability displacement ellipsoids [symmetry code: (A) = 1-x, 1-y, 1-z].

Dichloridooctakis(2-chlorobenzyl)di- μ_2 -hydroxido-di- μ_3 -oxido-tetratin(IV)

Crystal data



$$M_r = 1616.22$$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$$a = 10.986 (2) \text{ \AA}$$

$$b = 11.227 (2) \text{ \AA}$$

$$c = 13.573 (3) \text{ \AA}$$

$$\alpha = 74.656 (2)^\circ$$

$$\beta = 67.942 (2)^\circ$$

$$\gamma = 75.753 (2)^\circ$$

$$V = 1475.9 (6) \text{ \AA}^3$$

$$Z = 1$$

$$F(000) = 788$$

$$D_x = 1.818 \text{ Mg m}^{-3}$$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4036 reflections

$$\theta = 2.6\text{--}27.4^\circ$$

$$\mu = 2.17 \text{ mm}^{-1}$$

$$T = 298 \text{ K}$$

Block, colourless

$$0.44 \times 0.37 \times 0.33 \text{ mm}$$

Data collection

Bruker SMART CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(SADABS; Sheldrick, 1996)

$$T_{\min} = 0.449, T_{\max} = 0.535$$

7669 measured reflections

5112 independent reflections

3865 reflections with $I > 2\sigma(I)$

$$R_{\text{int}} = 0.015$$

$$\theta_{\max} = 25.0^\circ, \theta_{\min} = 1.7^\circ$$

$$h = -13 \rightarrow 13$$

$$k = -13 \rightarrow 13$$

$$l = -16 \rightarrow 10$$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.029$
 $wR(F^2) = 0.074$
 $S = 1.04$
 5112 reflections
 338 parameters
 0 restraints
 Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
 Hydrogen site location: inferred from neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0262P)^2 + 2.2448P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 1.27 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.57 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Sn1	0.59257 (3)	0.37338 (3)	0.44839 (2)	0.03438 (10)	
Sn2	0.49384 (3)	0.58881 (3)	0.25330 (2)	0.03713 (10)	
Cl1	0.35586 (14)	0.79133 (11)	0.28147 (11)	0.0516 (3)	
Cl2	0.6318 (2)	0.0780 (2)	0.34868 (19)	0.0704 (6)	0.693 (2)
Cl2'	0.2326 (5)	0.3325 (5)	0.5648 (4)	0.0704 (6)	0.307 (2)
Cl3	0.96390 (18)	0.40745 (16)	0.17259 (15)	0.0897 (6)	
Cl4	0.09931 (19)	0.5877 (2)	0.36984 (14)	0.0927 (6)	
Cl5	0.6057 (2)	0.72167 (16)	-0.03870 (14)	0.0908 (6)	
O1	0.6121 (3)	0.3933 (3)	0.2815 (2)	0.0444 (8)	
H1	0.6408	0.3297	0.2503	0.053*	
O2	0.4776 (3)	0.5414 (3)	0.4119 (2)	0.0365 (7)	
C1	0.4860 (7)	0.2203 (6)	0.5202 (5)	0.077 (2)	
H1A	0.5486	0.1446	0.5314	0.092*	
H1B	0.4238	0.2336	0.5908	0.092*	
C2	0.4103 (7)	0.2003 (5)	0.4555 (5)	0.0597 (16)	
C3	0.4702 (6)	0.1302 (5)	0.3739 (5)	0.0646 (17)	
H3A	0.5605	0.0982	0.3588	0.077*	0.307 (2)
C4	0.4013 (7)	0.1078 (6)	0.3160 (5)	0.0697 (18)	
H4	0.4440	0.0592	0.2629	0.084*	
C5	0.2691 (8)	0.1581 (7)	0.3376 (6)	0.078 (2)	
H5	0.2210	0.1431	0.2998	0.093*	
C6	0.2087 (7)	0.2301 (6)	0.4146 (6)	0.0775 (19)	
H6	0.1197	0.2663	0.4277	0.093*	
C7	0.2776 (8)	0.2497 (6)	0.4731 (5)	0.0724 (18)	
H7A	0.2335	0.2980	0.5263	0.087*	0.693 (2)
C8	0.8018 (5)	0.3527 (5)	0.4200 (4)	0.0541 (14)	
H8A	0.8337	0.4296	0.3766	0.065*	
H8B	0.8166	0.3377	0.4887	0.065*	
C9	0.8788 (5)	0.2460 (5)	0.3625 (4)	0.0469 (12)	
C10	0.9489 (5)	0.2595 (5)	0.2531 (4)	0.0510 (13)	
C11	1.0141 (5)	0.1589 (6)	0.2019 (5)	0.0630 (16)	
H11	1.0598	0.1719	0.1278	0.076*	
C12	1.0104 (6)	0.0399 (6)	0.2620 (6)	0.080 (2)	
H12	1.0524	-0.0287	0.2285	0.096*	

C13	0.9448 (7)	0.0222 (6)	0.3713 (7)	0.086 (2)
H13	0.9450	-0.0587	0.4121	0.103*
C14	0.8791 (6)	0.1218 (6)	0.4213 (5)	0.0666 (16)
H14	0.8337	0.1074	0.4955	0.080*
C15	0.3720 (6)	0.5022 (5)	0.2102 (4)	0.0541 (14)
H15A	0.4308	0.4523	0.1568	0.065*
H15B	0.3287	0.4446	0.2740	0.065*
C16	0.2673 (6)	0.5833 (5)	0.1664 (4)	0.0488 (13)
C17	0.1427 (6)	0.6287 (5)	0.2294 (5)	0.0596 (15)
C18	0.0470 (7)	0.7033 (6)	0.1868 (6)	0.0728 (18)
H18	-0.0357	0.7343	0.2323	0.087*
C19	0.0748 (7)	0.7312 (6)	0.0775 (6)	0.080 (2)
H19	0.0110	0.7812	0.0480	0.096*
C20	0.1997 (8)	0.6846 (6)	0.0096 (5)	0.0760 (19)
H20	0.2185	0.7022	-0.0650	0.091*
C21	0.2924 (6)	0.6141 (5)	0.0526 (5)	0.0632 (16)
H21	0.3754	0.5846	0.0065	0.076*
C22	0.6829 (5)	0.6492 (5)	0.1629 (4)	0.0546 (14)
H22A	0.7318	0.6346	0.2126	0.066*
H22B	0.7322	0.5957	0.1103	0.066*
C23	0.6833 (5)	0.7819 (4)	0.1041 (4)	0.0445 (12)
C24	0.6538 (5)	0.8249 (5)	0.0094 (4)	0.0491 (13)
C25	0.6586 (6)	0.9448 (5)	-0.0481 (4)	0.0566 (14)
H25	0.6375	0.9697	-0.1116	0.068*
C26	0.6952 (6)	1.0277 (5)	-0.0102 (5)	0.0672 (17)
H26	0.7000	1.1093	-0.0487	0.081*
C27	0.7244 (7)	0.9914 (6)	0.0830 (5)	0.0712 (18)
H27	0.7487	1.0480	0.1085	0.085*
C28	0.7178 (6)	0.8692 (6)	0.1404 (5)	0.0666 (16)
H28	0.7370	0.8454	0.2047	0.080*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.0413 (2)	0.02862 (17)	0.03475 (19)	-0.00165 (14)	-0.01569 (15)	-0.00788 (13)
Sn2	0.0458 (2)	0.03431 (18)	0.03224 (19)	-0.00562 (15)	-0.01628 (15)	-0.00401 (14)
Cl1	0.0610 (8)	0.0372 (6)	0.0546 (8)	0.0030 (6)	-0.0240 (7)	-0.0085 (6)
Cl2	0.0727 (14)	0.0590 (12)	0.0858 (15)	-0.0032 (10)	-0.0315 (12)	-0.0237 (10)
Cl2'	0.0727 (14)	0.0590 (12)	0.0858 (15)	-0.0032 (10)	-0.0315 (12)	-0.0237 (10)
Cl3	0.0743 (11)	0.0692 (11)	0.0932 (13)	-0.0020 (9)	-0.0141 (10)	0.0080 (9)
Cl4	0.0917 (13)	0.1304 (16)	0.0571 (10)	-0.0280 (12)	-0.0216 (9)	-0.0155 (10)
Cl5	0.1487 (18)	0.0662 (10)	0.0763 (12)	-0.0408 (11)	-0.0494 (12)	-0.0049 (8)
O1	0.054 (2)	0.0381 (18)	0.0397 (19)	-0.0030 (16)	-0.0209 (16)	-0.0015 (14)
O2	0.0440 (19)	0.0304 (16)	0.0341 (17)	0.0006 (14)	-0.0169 (15)	-0.0051 (13)
C1	0.137 (6)	0.064 (4)	0.056 (4)	-0.059 (4)	-0.052 (4)	0.014 (3)
C2	0.097 (5)	0.048 (3)	0.048 (3)	-0.043 (3)	-0.034 (3)	0.011 (3)
C3	0.081 (5)	0.060 (4)	0.067 (4)	-0.028 (3)	-0.041 (4)	0.003 (3)
C4	0.099 (5)	0.068 (4)	0.063 (4)	-0.032 (4)	-0.038 (4)	-0.013 (3)

C5	0.093 (6)	0.082 (5)	0.084 (5)	-0.041 (4)	-0.051 (4)	-0.002 (4)
C6	0.074 (5)	0.074 (4)	0.090 (5)	-0.029 (4)	-0.031 (4)	-0.003 (4)
C7	0.101 (6)	0.065 (4)	0.061 (4)	-0.042 (4)	-0.027 (4)	0.000 (3)
C8	0.047 (3)	0.063 (4)	0.064 (4)	0.000 (3)	-0.022 (3)	-0.033 (3)
C9	0.035 (3)	0.054 (3)	0.055 (3)	-0.001 (2)	-0.019 (2)	-0.016 (3)
C10	0.039 (3)	0.058 (3)	0.058 (4)	-0.003 (3)	-0.020 (3)	-0.013 (3)
C11	0.045 (3)	0.075 (4)	0.068 (4)	0.005 (3)	-0.015 (3)	-0.031 (3)
C12	0.062 (4)	0.068 (4)	0.108 (6)	0.004 (3)	-0.016 (4)	-0.044 (4)
C13	0.069 (5)	0.052 (4)	0.112 (6)	-0.008 (3)	-0.016 (4)	0.003 (4)
C14	0.051 (4)	0.064 (4)	0.069 (4)	0.004 (3)	-0.014 (3)	-0.007 (3)
C15	0.065 (4)	0.048 (3)	0.060 (4)	-0.013 (3)	-0.029 (3)	-0.013 (3)
C16	0.061 (4)	0.045 (3)	0.055 (3)	-0.013 (3)	-0.030 (3)	-0.014 (2)
C17	0.063 (4)	0.067 (4)	0.057 (4)	-0.013 (3)	-0.026 (3)	-0.015 (3)
C18	0.064 (4)	0.079 (4)	0.085 (5)	-0.005 (3)	-0.038 (4)	-0.020 (4)
C19	0.082 (5)	0.077 (5)	0.095 (6)	-0.004 (4)	-0.057 (5)	-0.005 (4)
C20	0.106 (6)	0.083 (5)	0.058 (4)	-0.030 (4)	-0.045 (4)	-0.005 (3)
C21	0.066 (4)	0.063 (4)	0.074 (4)	-0.011 (3)	-0.030 (3)	-0.025 (3)
C22	0.047 (3)	0.046 (3)	0.058 (3)	-0.004 (2)	-0.016 (3)	0.005 (3)
C23	0.033 (3)	0.041 (3)	0.048 (3)	-0.006 (2)	-0.006 (2)	-0.002 (2)
C24	0.049 (3)	0.044 (3)	0.047 (3)	-0.008 (2)	-0.009 (3)	-0.006 (2)
C25	0.064 (4)	0.046 (3)	0.048 (3)	-0.005 (3)	-0.013 (3)	-0.002 (3)
C26	0.072 (4)	0.039 (3)	0.070 (4)	-0.002 (3)	-0.007 (3)	-0.007 (3)
C27	0.089 (5)	0.065 (4)	0.062 (4)	-0.033 (4)	-0.011 (4)	-0.016 (3)
C28	0.065 (4)	0.076 (4)	0.061 (4)	-0.020 (3)	-0.024 (3)	-0.005 (3)

Geometric parameters (\AA , $^{\circ}$)

Sn1—O1	2.148 (3)	C9—C14	1.412 (7)
Sn1—O2	2.050 (3)	C10—C11	1.385 (8)
Sn1—O2 ⁱ	2.146 (3)	C11—C12	1.370 (8)
Sn1—C1	2.126 (5)	C11—H11	0.9300
Sn1—C8	2.146 (5)	C12—C13	1.368 (9)
Sn2—O1	2.276 (3)	C12—H12	0.9300
Sn2—O2	2.025 (3)	C13—C14	1.367 (9)
Sn2—C15	2.147 (5)	C13—H13	0.9300
Sn2—C22	2.149 (5)	C14—H14	0.9300
Sn2—Cl1	2.4376 (13)	C15—C16	1.489 (7)
Cl2—C3	1.658 (7)	C15—H15A	0.9700
Cl2—H3A	0.7301	C15—H15B	0.9700
Cl2'—C7	1.606 (8)	C16—C17	1.368 (8)
Cl2'—H7A	0.7242	C16—C21	1.424 (7)
Cl3—C10	1.737 (6)	C17—C18	1.381 (8)
Cl4—C17	1.742 (6)	C18—C19	1.361 (9)
Cl5—C24	1.735 (5)	C18—H18	0.9300
O1—H1	0.8590	C19—C20	1.399 (9)
C1—C2	1.503 (7)	C19—H19	0.9300
C1—H1A	0.9700	C20—C21	1.345 (8)
C1—H1B	0.9700	C20—H20	0.9300

C2—C7	1.379 (9)	C21—H21	0.9300
C2—C3	1.391 (8)	C22—C23	1.492 (6)
C3—C4	1.379 (7)	C22—H22A	0.9700
C3—H3A	0.9301	C22—H22B	0.9700
C4—C5	1.371 (9)	C23—C24	1.377 (7)
C4—H4	0.9300	C23—C28	1.388 (7)
C5—C6	1.359 (9)	C24—C25	1.369 (7)
C5—H5	0.9300	C25—C26	1.374 (8)
C6—C7	1.371 (8)	C25—H25	0.9300
C6—H6	0.9300	C26—C27	1.355 (8)
C7—H7A	0.9300	C26—H26	0.9300
C8—C9	1.502 (7)	C27—C28	1.391 (8)
C8—H8A	0.9700	C27—H27	0.9300
C8—H8B	0.9700	C28—H28	0.9300
C9—C10	1.379 (7)		
O2—Sn1—C1	114.2 (2)	C10—C9—C8	124.6 (5)
O2—Sn1—O2 ⁱ	73.49 (12)	C14—C9—C8	119.6 (5)
C1—Sn1—O2 ⁱ	97.25 (19)	C9—C10—C11	123.1 (5)
O2—Sn1—C8	123.37 (17)	C9—C10—Cl3	120.6 (4)
C1—Sn1—C8	122.4 (3)	C11—C10—Cl3	116.3 (5)
O2 ⁱ —Sn1—C8	97.54 (16)	C12—C11—C10	119.1 (6)
O2—Sn1—O1	74.51 (12)	C12—C11—H11	120.5
C1—Sn1—O1	100.54 (17)	C10—C11—H11	120.5
O2 ⁱ —Sn1—O1	147.55 (11)	C13—C12—C11	119.8 (6)
C8—Sn1—O1	95.50 (17)	C13—C12—H12	120.1
O2—Sn2—C15	114.44 (17)	C11—C12—H12	120.1
O2—Sn2—C22	108.14 (17)	C14—C13—C12	120.9 (6)
C15—Sn2—C22	131.7 (2)	C14—C13—H13	119.6
O2—Sn2—O1	72.21 (11)	C12—C13—H13	119.6
C15—Sn2—O1	85.91 (16)	C13—C14—C9	121.3 (6)
C22—Sn2—O1	86.30 (16)	C13—C14—H14	119.4
O2—Sn2—Cl1	90.33 (9)	C9—C14—H14	119.4
C15—Sn2—Cl1	102.16 (15)	C16—C15—Sn2	118.8 (3)
C22—Sn2—Cl1	99.22 (15)	C16—C15—H15A	107.6
O1—Sn2—Cl1	162.53 (9)	Sn2—C15—H15A	107.6
C3—Cl2—H3A	3.0	C16—C15—H15B	107.6
C7—Cl2'—H7A	15.7	Sn2—C15—H15B	107.6
Sn1—O1—Sn2	100.13 (13)	H15A—C15—H15B	107.0
Sn1—O1—H1	120.9	C17—C16—C21	115.9 (5)
Sn2—O1—H1	137.0	C17—C16—C15	124.1 (5)
Sn2—O2—Sn1	112.76 (14)	C21—C16—C15	119.9 (5)
Sn2—O2—Sn1 ⁱ	139.33 (14)	C16—C17—C18	123.0 (6)
Sn1—O2—Sn1 ⁱ	106.51 (12)	C16—C17—Cl4	119.2 (4)
C2—C1—Sn1	114.9 (3)	C18—C17—Cl4	117.8 (5)
C2—C1—H1A	108.5	C19—C18—C17	119.4 (6)
Sn1—C1—H1A	108.5	C19—C18—H18	120.3
C2—C1—H1B	108.5	C17—C18—H18	120.3

Sn1—C1—H1B	108.5	C18—C19—C20	119.8 (6)
H1A—C1—H1B	107.5	C18—C19—H19	120.1
C7—C2—C3	115.8 (5)	C20—C19—H19	120.1
C7—C2—C1	122.3 (6)	C21—C20—C19	119.8 (6)
C3—C2—C1	121.9 (6)	C21—C20—H20	120.1
C4—C3—C2	122.6 (6)	C19—C20—H20	120.1
C4—C3—Cl2	121.9 (6)	C20—C21—C16	122.0 (6)
C2—C3—Cl2	115.4 (5)	C20—C21—H21	119.0
C4—C3—H3A	119.9	C16—C21—H21	119.0
C2—C3—H3A	117.5	C23—C22—Sn2	118.1 (3)
Cl2—C3—H3A	2.4	C23—C22—H22A	107.8
C5—C4—C3	119.1 (6)	Sn2—C22—H22A	107.8
C5—C4—H4	120.5	C23—C22—H22B	107.8
C3—C4—H4	120.5	Sn2—C22—H22B	107.8
C6—C5—C4	119.7 (6)	H22A—C22—H22B	107.1
C6—C5—H5	120.2	C24—C23—C28	115.7 (5)
C4—C5—H5	120.2	C24—C23—C22	122.7 (5)
C5—C6—C7	120.7 (7)	C28—C23—C22	121.5 (5)
C5—C6—H6	119.7	C25—C24—C23	123.7 (5)
C7—C6—H6	119.7	C25—C24—Cl5	118.1 (4)
C6—C7—C2	122.1 (7)	C23—C24—Cl5	118.1 (4)
C6—C7—Cl2'	130.8 (7)	C24—C25—C26	118.6 (5)
C2—C7—Cl2'	107.1 (5)	C24—C25—H25	120.7
C6—C7—H7A	119.0	C26—C25—H25	120.7
C2—C7—H7A	119.0	C27—C26—C25	120.4 (5)
Cl2'—C7—H7A	12.1	C27—C26—H26	119.8
C9—C8—Sn1	111.3 (3)	C25—C26—H26	119.8
C9—C8—H8A	109.4	C26—C27—C28	119.8 (6)
Sn1—C8—H8A	109.4	C26—C27—H27	120.1
C9—C8—H8B	109.4	C28—C27—H27	120.1
Sn1—C8—H8B	109.4	C23—C28—C27	121.7 (6)
H8A—C8—H8B	108.0	C23—C28—H28	119.2
C10—C9—C14	115.9 (5)	C27—C28—H28	119.2
O2—Sn1—O1—Sn2	4.51 (11)	Sn1—C8—C9—C10	100.7 (5)
C1—Sn1—O1—Sn2	116.9 (2)	Sn1—C8—C9—C14	-77.6 (5)
O2 ⁱ —Sn1—O1—Sn2	-5.2 (3)	C14—C9—C10—C11	1.2 (8)
C8—Sn1—O1—Sn2	-118.63 (17)	C8—C9—C10—C11	-177.1 (5)
O2—Sn2—O1—Sn1	-4.61 (11)	C14—C9—C10—Cl3	-176.6 (4)
C15—Sn2—O1—Sn1	-121.91 (19)	C8—C9—C10—Cl3	5.0 (7)
C22—Sn2—O1—Sn1	105.78 (19)	C9—C10—C11—C12	-0.5 (9)
Cl1—Sn2—O1—Sn1	-3.5 (4)	Cl3—C10—C11—C12	177.4 (5)
C15—Sn2—O2—Sn1	82.0 (2)	C10—C11—C12—C13	-1.2 (10)
C22—Sn2—O2—Sn1	-74.7 (2)	C11—C12—C13—C14	2.0 (11)
O1—Sn2—O2—Sn1	5.17 (13)	C12—C13—C14—C9	-1.3 (10)
Cl1—Sn2—O2—Sn1	-174.49 (13)	C10—C9—C14—C13	-0.3 (8)
C15—Sn2—O2—Sn1 ⁱ	-114.1 (3)	C8—C9—C14—C13	178.1 (6)
C22—Sn2—O2—Sn1 ⁱ	89.3 (3)	O2—Sn2—C15—C16	118.1 (4)

O1—Sn2—O2—Sn1 ⁱ	169.1 (3)	C22—Sn2—C15—C16	−92.2 (5)
C11—Sn2—O2—Sn1 ⁱ	−10.5 (2)	O1—Sn2—C15—C16	−173.6 (4)
C1—Sn1—O2—Sn2	−100.2 (2)	C11—Sn2—C15—C16	22.1 (5)
O2 ⁱ —Sn1—O2—Sn2	169.2 (2)	Sn2—C15—C16—C17	−83.9 (6)
C8—Sn1—O2—Sn2	81.0 (2)	Sn2—C15—C16—C21	98.5 (5)
O1—Sn1—O2—Sn2	−5.41 (13)	C21—C16—C17—C18	−2.0 (8)
C1—Sn1—O2—Sn1 ⁱ	90.6 (2)	C15—C16—C17—C18	−179.7 (5)
O2 ⁱ —Sn1—O2—Sn1 ⁱ	0.000 (1)	C21—C16—C17—Cl4	176.0 (4)
C8—Sn1—O2—Sn1 ⁱ	−88.2 (2)	C15—C16—C17—Cl4	−1.6 (7)
O1—Sn1—O2—Sn1 ⁱ	−174.57 (16)	C16—C17—C18—C19	1.9 (9)
O2—Sn1—C1—C2	55.6 (6)	Cl4—C17—C18—C19	−176.2 (5)
O2 ⁱ —Sn1—C1—C2	130.8 (5)	C17—C18—C19—C20	−0.2 (10)
C8—Sn1—C1—C2	−125.5 (5)	C18—C19—C20—C21	−1.1 (10)
O1—Sn1—C1—C2	−22.0 (6)	C19—C20—C21—C16	0.9 (9)
Sn1—C1—C2—C7	−94.6 (6)	C17—C16—C21—C20	0.6 (8)
Sn1—C1—C2—C3	85.5 (6)	C15—C16—C21—C20	178.4 (5)
C7—C2—C3—C4	−1.9 (8)	O2—Sn2—C22—C23	−116.2 (4)
C1—C2—C3—C4	178.0 (5)	C15—Sn2—C22—C23	92.7 (5)
C7—C2—C3—Cl2	176.7 (4)	O1—Sn2—C22—C23	173.9 (4)
C1—C2—C3—Cl2	−3.4 (7)	Cl1—Sn2—C22—C23	−22.8 (4)
C2—C3—C4—C5	1.3 (9)	Sn2—C22—C23—C24	−74.9 (6)
Cl2—C3—C4—C5	−177.2 (5)	Sn2—C22—C23—C28	107.0 (5)
C3—C4—C5—C6	0.7 (9)	C28—C23—C24—C25	0.8 (8)
C4—C5—C6—C7	−2.0 (10)	C22—C23—C24—C25	−177.4 (5)
C5—C6—C7—C2	1.3 (9)	C28—C23—C24—Cl5	−178.4 (4)
C5—C6—C7—Cl2'	177.9 (6)	C22—C23—C24—Cl5	3.5 (7)
C3—C2—C7—C6	0.6 (8)	C23—C24—C25—C26	0.2 (9)
C1—C2—C7—C6	−179.3 (5)	Cl5—C24—C25—C26	179.3 (4)
C3—C2—C7—Cl2'	−176.7 (4)	C24—C25—C26—C27	−0.8 (9)
C1—C2—C7—Cl2'	3.3 (7)	C25—C26—C27—C28	0.3 (10)
O2—Sn1—C8—C9	−131.3 (3)	C24—C23—C28—C27	−1.3 (8)
C1—Sn1—C8—C9	50.0 (5)	C22—C23—C28—C27	176.9 (5)
O2 ⁱ —Sn1—C8—C9	153.6 (4)	C26—C27—C28—C23	0.7 (10)
O1—Sn1—C8—C9	−56.2 (4)		

Symmetry code: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (\AA , °)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O1—H1 \cdots Cl2	0.86	2.80	3.386 (4)	127