

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2-Bromo-4-methylbenzonitrile

Muhammad Shahid, Munawar Ali Munawar,\* Sohail Nadeem, Waqar Nasir and Muhammad Salim

Institute of Chemistry, University of the Punjab, New Campus, Lahore, Pakistan  
Correspondence e-mail: munawaralimunawar@yahoo.com


Received 16 November 2009; accepted 17 November 2009

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.033;  $wR$  factor = 0.084; data-to-parameter ratio = 20.9.

The title molecule,  $\text{C}_8\text{H}_6\text{BrN}$ , is almost planar (r.m.s. deviation for the non-H atoms = 0.008 Å). In the crystal, weak  $\pi-\pi$  stacking interactions [centroid-centroid separations = 3.782 (2) and 3.919 (2) Å] generate [100] columns of molecules.

## Related literature

For the synthesis, see: Johnson & Sandborn (1941). 2-Bromo-4-methylbenzonitrile derivatives are used as intermediates in the synthesis of phthalocyanine dyes. For applications of phthalocyanine dyes in photo redox reactions and photo-dynamic cancer therapy, see: Simon & Sirlin (1989); Simon *et al.* (1989).



## Experimental

## Crystal data

 $\text{C}_8\text{H}_6\text{BrN}$  $M_r = 196.05$ 

Triclinic,  $P\bar{1}$   
 $a = 7.5168$  (11) Å  
 $b = 7.8383$  (11) Å  
 $c = 7.9428$  (11) Å  
 $\alpha = 69.243$  (7)°  
 $\beta = 64.375$  (8)°  
 $\gamma = 87.567$  (8)°

$V = 391.14$  (10) Å<sup>3</sup>  
 $Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 5.17$  mm<sup>-1</sup>  
 $T = 296$  K  
 $0.41 \times 0.28 \times 0.19$  mm

## Data collection

Bruker Kappa APEXII CCD diffractometer  
Absorption correction: multi-scan (SADABS; Bruker, 2007)  
 $T_{\min} = 0.226$ ,  $T_{\max} = 0.440$

8084 measured reflections  
1921 independent reflections  
1244 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.025$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.033$   
 $wR(F^2) = 0.084$   
 $S = 1.01$   
1921 reflections

92 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.44$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.49$  e Å<sup>-3</sup>

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON.

Muhammad Shahid acknowledges the Higher Education Commission of Pakistan for providing funds, the Institute of Chemistry, University of the Punjab, for providing research facilities and the Education Department, Government of the Punjab, for their co-operation.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5232).

## References

- Bruker (2007). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.  
Johnson, J. R. & Sandborn, L. T. (1941). *Org. Synth. Coll.* **1**, 111–116.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Simon, J., Bassoul, P. & Norvez, S. (1989). *New. J. Chem.* **13**, 13–31.  
Simon, J. & Sirlin, C. (1989). *Pure Appl. Chem.* **61**, 1625–1629.  
Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## supporting information

*Acta Cryst.* (2009). E65, o3166 [doi:10.1107/S1600536809048983]

**2-Bromo-4-methylbenzotrile**

**Muhammad Shahid, Munawar Ali Munawar, Sohail Nadeem, Waqar Nasir and Muhammad Salim**

**S1. Comment**

Synthesis of 2-bromo-4-methylbenzotrile derivatives are important compounds due to their use as intermediates in the synthesis of phthalocyanine dyes. The substituted phthalocyanine dyes have been used for photo redox reactions (Simon & Sirlin, 1989) and photodynamic cancer therapy (Simon *et al.*, 1989).


The title compound(I) is almost planar. The cyano plane (C4/C8/N1) is oriented at a dihedral angle of 79.7 (3)° with respect to aromatic ring (C1/C2/C3/C4/C5/C6). The dihedral angle between the plane containing the methyl carbon (C1/C2/C6/C7) and aromatic ring plane is 0.22 (0.18)°. No significant intermolecular or intramolecular hydrogen bonding interaction has been observed in the molecule.

**S2. Experimental**

3-Bromo-4-amino toluene (10 g, 54 mmol) (Johnson & Sandborn, 1941) was dissolved in HCl (30 ml, 17%). The mixture was cooled to 273 K in an ice-salt mixture. Over 5 min, an aqueous solution (9 ml) of NaNO<sub>2</sub> (4.3 g) was added to the above mixture. The temperature was maintained at 273-278 K. A mixture of aqueous solution (6%) of Cu(I)cyanide and KCN (40%) was heated to 333 K and added to the above cold neutralized diazonium salt solution. After work up of reaction, colourless blocks of (I) were obtained by the slow evaporation of water.


**S3. Refinement**

The H atoms were geometrically placed (C—H = 0.93–0.96Å) and refined as riding with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$  or  $1.5U_{\text{eq}}(\text{methyl C})$ .



**Figure 1**

The molecular structure of (I) with 50% displacement ellipsoids.

**Figure 2**

Unit cell packing diagram.

**2-bromo-4-methylbenzonitrile***Crystal data* $C_8H_7BrN$  $M_r = 196.05$ Triclinic,  $P\bar{1}$ Hall symbol:  $-P\ 1$  $a = 7.5168\ (11)\ \text{\AA}$  $b = 7.8383\ (11)\ \text{\AA}$  $c = 7.9428\ (11)\ \text{\AA}$  $\alpha = 69.243\ (7)^\circ$  $\beta = 64.375\ (8)^\circ$  $\gamma = 87.567\ (8)^\circ$  $V = 391.14\ (10)\ \text{\AA}^3$  $Z = 2$  $F(000) = 192$  $D_x = 1.665\ \text{Mg m}^{-3}$ Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$ 

Cell parameters from 3073 reflections

 $\theta = 2.2\text{--}21.2^\circ$  $\mu = 5.17\ \text{mm}^{-1}$  $T = 296\ \text{K}$ 

Block, colourless

 $0.41 \times 0.28 \times 0.19\ \text{mm}$ *Data collection*Bruker Kappa APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

 $\varphi$  and  $\omega$  scansAbsorption correction: multi-scan  
(*SADABS*; Bruker, 2007) $T_{\min} = 0.226$ ,  $T_{\max} = 0.440$ 

8084 measured reflections

1921 independent reflections

1244 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$   
 $\theta_{\text{max}} = 28.3^\circ$ ,  $\theta_{\text{min}} = 2.8^\circ$   
 $h = -9 \rightarrow 9$

$k = -10 \rightarrow 10$   
 $l = -10 \rightarrow 10$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.033$   
 $wR(F^2) = 0.084$   
 $S = 1.01$   
 1921 reflections  
 92 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0326P)^2 + 0.2649P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\text{max}} < 0.001$   
 $\Delta\rho_{\text{max}} = 0.44 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\text{min}} = -0.49 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.32613 (7)	0.89370 (4)	0.35907 (6)	0.07874 (19)
C1	0.2019 (4)	0.3498 (4)	0.7427 (4)	0.0525 (7)
C2	0.2395 (4)	0.5395 (4)	0.6606 (4)	0.0512 (7)
H2	0.2419	0.6033	0.7387	0.061*
C3	0.2732 (4)	0.6352 (4)	0.4660 (4)	0.0457 (6)
C4	0.2711 (4)	0.5439 (4)	0.3461 (4)	0.0445 (6)
C5	0.2349 (5)	0.3537 (4)	0.4270 (5)	0.0532 (7)
H5	0.2337	0.2899	0.3485	0.064*
C6	0.2008 (5)	0.2588 (4)	0.6222 (5)	0.0582 (8)
H6	0.1765	0.1310	0.6747	0.070*
C7	0.1639 (6)	0.2446 (5)	0.9569 (5)	0.0764 (10)
H7A	0.2191	0.3191	1.0006	0.115*
H7B	0.2253	0.1339	0.9654	0.115*
H7C	0.0231	0.2136	1.0415	0.115*
C8	0.3088 (5)	0.6413 (4)	0.1400 (5)	0.0540 (7)
N1	0.3389 (5)	0.7126 (4)	-0.0230 (5)	0.0771 (9)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.1226 (4)	0.0413 (2)	0.0893 (3)	0.01752 (18)	-0.0609 (3)	-0.02676 (18)
C1	0.0514 (18)	0.0578 (18)	0.0451 (17)	0.0061 (14)	-0.0231 (14)	-0.0136 (14)

C2	0.0571 (18)	0.0587 (18)	0.0515 (18)	0.0159 (14)	-0.0288 (15)	-0.0308 (15)
C3	0.0525 (17)	0.0403 (14)	0.0491 (17)	0.0096 (12)	-0.0251 (14)	-0.0193 (13)
C4	0.0449 (16)	0.0480 (16)	0.0410 (16)	0.0047 (12)	-0.0188 (13)	-0.0174 (13)
C5	0.0641 (19)	0.0478 (16)	0.0525 (18)	0.0029 (14)	-0.0249 (15)	-0.0249 (14)
C6	0.069 (2)	0.0437 (16)	0.056 (2)	0.0005 (14)	-0.0258 (16)	-0.0151 (15)
C7	0.086 (3)	0.085 (3)	0.050 (2)	0.007 (2)	-0.0321 (19)	-0.0132 (18)
C8	0.0586 (19)	0.0560 (18)	0.0478 (19)	-0.0003 (14)	-0.0225 (15)	-0.0205 (15)
N1	0.098 (2)	0.077 (2)	0.0512 (18)	-0.0056 (17)	-0.0332 (17)	-0.0171 (16)

*Geometric parameters (Å, °)*

Br1—C3	1.882 (3)	C4—C8	1.440 (4)
C1—C2	1.380 (4)	C5—C6	1.371 (4)
C1—C6	1.384 (4)	C5—H5	0.9300
C1—C7	1.503 (4)	C6—H6	0.9300
C2—C3	1.368 (4)	C7—H7A	0.9600
C2—H2	0.9300	C7—H7B	0.9600
C3—C4	1.384 (4)	C7—H7C	0.9600
C4—C5	1.383 (4)	C8—N1	1.133 (4)
C2—C1—C6	118.2 (3)	C6—C5—H5	119.9
C2—C1—C7	121.0 (3)	C4—C5—H5	119.9
C6—C1—C7	120.8 (3)	C5—C6—C1	121.2 (3)
C3—C2—C1	121.0 (3)	C5—C6—H6	119.4
C3—C2—H2	119.5	C1—C6—H6	119.4
C1—C2—H2	119.5	C1—C7—H7A	109.5
C2—C3—C4	120.8 (3)	C1—C7—H7B	109.5
C2—C3—Br1	119.6 (2)	H7A—C7—H7B	109.5
C4—C3—Br1	119.6 (2)	C1—C7—H7C	109.5
C5—C4—C3	118.6 (3)	H7A—C7—H7C	109.5
C5—C4—C8	119.6 (3)	H7B—C7—H7C	109.5
C3—C4—C8	121.8 (3)	N1—C8—C4	177.7 (3)
C6—C5—C4	120.3 (3)		