

7-Fluoro-6-nitroquinazolin-4(3H)-one

Yundeng Wu,^{a,b} Ancheng Ji,^b Aihua Zhang^{b*} and Yipeng Shen^b

^aSchool of Pharmaceutical Sciences, Nanjing University of Technology, No. 5 Xinmofan Road, Nanjing 210009, People's Republic of China, and ^bDepartment of Medicinal Chemistry, Jiangsu Provincial Institute of Materia Medica, Nanjing University of Technology, No. 26 Majia Street, Nanjing 210009, People's Republic of China

Correspondence e-mail: fafazhang928@yahoo.com.cn


Received 30 October 2009; accepted 6 November 2009

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.050; wR factor = 0.160; data-to-parameter ratio = 10.7.

The quinazolinone unit of the title compound, $\text{C}_8\text{H}_4\text{FN}_3\text{O}_3$, is essentially planar, with a maximum deviation of $0.0538(14)\text{ \AA}$ for the O atom. The nitro group is twisted by $12.0(3)^\circ$ from the mean plane of the quinazolinone ring system. The crystal structure is stabilized by intermolecular N—H···O, C—H···N and C—H···O hydrogen bonds.

Related literature

The title compound is used as an intermediate for the production of several multi-targeted Raf kinase inhibitors, such as 4(3*H*)-quinazolinone and its derivatives, see: Bridges *et al.* (1996); Kim *et al.* (2008). For the antitumor activities of quinolines, see: Labuda *et al.* (2009). For synthetic aspects, see: Newcastle *et al.* (1996). For bond-length data, see: Allen *et al.* (1987).

**Experimental***Crystal data*

$\text{C}_8\text{H}_4\text{FN}_3\text{O}_3$	$\gamma = 83.83(3)^\circ$
$M_r = 209.14$	$V = 401.70(16)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 5.6360(11)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 8.409(2)\text{ \AA}$	$\mu = 0.15\text{ mm}^{-1}$
$c = 8.674(2)\text{ \AA}$	$T = 293\text{ K}$
$\alpha = 79.38(3)^\circ$	$0.30 \times 0.20 \times 0.20\text{ mm}$
$\beta = 89.23(3)^\circ$	

Data collection

Enraf–Nonius CAD-4 diffractometer
Absorption correction: ψ scan (North *et al.*, 1968)
 $T_{\min} = 0.956$, $T_{\max} = 0.971$
1623 measured reflections

1461 independent reflections
1131 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.018$
3 standard reflections
every 200 reflections
intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.160$
 $S = 1.00$
1461 reflections

137 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.23\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1A···O1 ⁱ	0.86	1.98	2.815 (2)	165
C1—H1B···O2 ⁱⁱ	0.93	2.47	3.396 (3)	179
C7—H7A···N2 ⁱⁱⁱ	0.93	2.50	3.422 (3)	171

Symmetry codes: (i) $-x, -y, -z + 1$; (ii) $x + 1, y, z + 1$; (iii) $-x + 1, -y + 1, -z$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1995); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *PLATON* (Spek, 2009).

This research work was supported financially by the Research Funds of Jiangsu Provincial Institute of Materia Medica (No. SX200801).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: PV2231).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bridges, A. J., Zhou, H., Cody, D. R., Newcastle, G. W., McMichael, A., Showalter, H. D. H., Fry, D. W., Kraker, A. J. & Denny, W. A. (1996). *J. Med. Chem.* **39**, 267–276.
- Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Harms, K. & Wocadlo, S. (1995). *XCAD4*. University of Marburg, Germany.
- Kim, Y. H., Choi, H., Lee, J., Hwang, I.-C., Moon, S. K., Kim, S. J., Lee, H. W., Im, D. S., Lee, S. S., Ahn, S. K., Kim, S. W., Han, Cheol K., Yoon, J. H., Lee, K. J. & Choi, N. S. (2008). *Bioorg. Med. Chem. Lett.* **18**, 6279–6282.
- Labuda, J., Ovadekova, R. & Galandova, J. (2009). *Mikrochim. Acta*, **164**, 371–377.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst. A* **24**, 351–359.
- Newcastle, G. W., Palmer, B. D., Bridges, A. J., Showalter, H. D. H., Sun, L., Nelson, J., McMichael, A., Kraker, A. J., Fry, D. W. & Denny, W. A. (1996). *J. Med. Chem.* **39**, 918–928.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.

supporting information

Acta Cryst. (2009). E65, o3075 [doi:10.1107/S1600536809046984]

7-Fluoro-6-nitroquinazolin-4(3*H*)-one

Yundeng Wu, Ancheng Ji, Aihua Zhang and Yipeng Shen

S1. Comment

4(*3H*)-Quinazolinone and its derivatives have been investigated extensively, owing to their important role in the synthesis of several multi-kinase inhibitors and to their potentially beneficial antitumor activities in many types of malignancies (Labuda *et al.*, 2009).

As part of our studies on the synthesis of 4(*3H*)-quinazolinone and its derivatives, the title compound, (I), which is used as the key intermediate (Rewcastle *et al.*, 1996), has been synthesized in our laboratory. We report herein the crystal structure of the title compound.


The molecule of the title compound is planar (Fig. 1). The quinazolinone moiety is essentially planar with maximum deviation for any atoms being 0.0538 (14) for O1. The nitro group is twisted from the mean-plane of the quinazolinone ring by 12.0 (3) $^{\circ}$. The bond lengths and angles in (I) are within normal ranges (Allen *et al.*, 1987). The crystal structure of (I) is stabilized by classical and non-classical intermolecular hydrogen bonds of the types N—H \cdots O, C—H \cdots N and C—H \cdots O; details have been provided in Table 1 and presented as a packing diagram in Fig. 2.

S2. Experimental


The title compound, was prepared by following a reported procedure (Rewcastle *et al.*, 1996). 7-Fluoroquinazolin-4(*3H*)-one (47.4 g, 0.29 mmol) was added to a mixture of concentrated H₂SO₄ (100 ml) and fuming HNO₃ (100 ml), and heated at 373 K for 1 h. The crude product, 7-fluoro-6-nitroquinazolin-4(*3H*)-one, was obtained by pouring the reacting mixture onto ice-water (1500 ml). The crystals of (I) suitable for X-ray diffraction studies were obtained by recrystallization from acetic acid.

S3. Refinement

H atoms were positioned geometrically at distances N—H = 0.86 Å and C—H = 0.93 Å and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2$ times $U_{\text{eq}}(\text{parent atoms})$.

**Figure 1**

The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at 30% probability levels.

**Figure 2**

A packing diagram of the title compound. Hydrogen bonds are shown as dashed lines.

7-Fluoro-6-nitroquinazolin-4(3H)-one

Crystal data

$C_8H_4FN_3O_3$
 $M_r = 209.14$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 5.6360 (11) \text{ \AA}$
 $b = 8.409 (2) \text{ \AA}$
 $c = 8.674 (2) \text{ \AA}$
 $\alpha = 79.38 (3)^\circ$
 $\beta = 89.23 (3)^\circ$

$\gamma = 83.83 (3)^\circ$
 $V = 401.70 (16) \text{ \AA}^3$
 $Z = 2$
 $F(000) = 212$
 $D_x = 1.729 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Cell parameters from 25 reflections
 $\theta = 9-13^\circ$
 $\mu = 0.15 \text{ mm}^{-1}$

$T = 293\text{ K}$
Block, colorless

Data collection

Enraf–Nonius CAD-4
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
 $\omega/2\theta$ scans
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.956$, $T_{\max} = 0.971$
1623 measured reflections

0.30 × 0.20 × 0.20 mm
1461 independent reflections
1131 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.018$
 $\theta_{\max} = 25.3^\circ$, $\theta_{\min} = 2.4^\circ$
 $h = 0 \rightarrow 6$
 $k = -10 \rightarrow 10$
 $l = -10 \rightarrow 10$
3 standard reflections every 200 reflections
intensity decay: 1%

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.050$
 $wR(F^2) = 0.160$
 $S = 1.00$
1461 reflections
137 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.1P)^2 + 0.12P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.23\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.25\text{ e \AA}^{-3}$
Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{1/4}$
Extinction coefficient: 0.062 (16)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
F	0.1138 (3)	0.4301 (2)	-0.33786 (16)	0.0639 (5)
N1	0.1677 (3)	0.1637 (2)	0.3773 (2)	0.0413 (5)
H1A	0.1629	0.1209	0.4752	0.050*
O1	-0.1399 (3)	0.0293 (2)	0.32253 (18)	0.0527 (6)
C1	0.3360 (4)	0.2651 (3)	0.3291 (3)	0.0419 (6)
H1B	0.4402	0.2828	0.4049	0.050*
N2	0.3644 (3)	0.3390 (2)	0.1878 (2)	0.0406 (5)
C2	0.0044 (4)	0.1261 (3)	0.2775 (2)	0.0380 (6)
O2	-0.2795 (5)	0.3272 (3)	-0.3951 (2)	0.0932 (9)
C3	0.0261 (4)	0.2086 (2)	0.1159 (2)	0.0331 (5)
N3	-0.2740 (4)	0.2376 (3)	-0.2685 (2)	0.0486 (6)
O3	-0.4036 (3)	0.1313 (2)	-0.2327 (2)	0.0630 (6)

C4	-0.1305 (4)	0.1862 (3)	0.0011 (3)	0.0376 (6)
H4A	-0.2523	0.1200	0.0276	0.045*
C5	-0.1046 (4)	0.2618 (3)	-0.1509 (3)	0.0380 (5)
C6	0.0818 (4)	0.3590 (3)	-0.1903 (2)	0.0396 (6)
C7	0.2349 (4)	0.3832 (3)	-0.0789 (3)	0.0387 (6)
H7A	0.3567	0.4491	-0.1068	0.046*
C8	0.2091 (4)	0.3091 (2)	0.0771 (2)	0.0334 (5)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
F	0.0696 (10)	0.0883 (11)	0.0293 (8)	-0.0238 (8)	-0.0026 (6)	0.0104 (7)
N1	0.0493 (11)	0.0479 (11)	0.0260 (9)	-0.0160 (9)	-0.0022 (8)	0.0010 (8)
O1	0.0562 (11)	0.0618 (11)	0.0393 (9)	-0.0333 (8)	-0.0015 (7)	0.0086 (8)
C1	0.0434 (13)	0.0495 (13)	0.0342 (12)	-0.0154 (10)	-0.0056 (9)	-0.0051 (10)
N2	0.0413 (10)	0.0474 (11)	0.0338 (10)	-0.0172 (8)	-0.0030 (8)	-0.0019 (8)
C2	0.0401 (12)	0.0390 (12)	0.0338 (12)	-0.0107 (9)	-0.0005 (9)	-0.0001 (9)
O2	0.1106 (19)	0.123 (2)	0.0446 (12)	-0.0476 (15)	-0.0370 (12)	0.0113 (12)
C3	0.0356 (11)	0.0321 (11)	0.0306 (11)	-0.0060 (9)	-0.0001 (8)	-0.0019 (8)
N3	0.0487 (12)	0.0573 (13)	0.0421 (12)	-0.0065 (10)	-0.0087 (9)	-0.0138 (10)
O3	0.0533 (11)	0.0733 (13)	0.0676 (13)	-0.0208 (10)	-0.0126 (9)	-0.0172 (10)
C4	0.0372 (12)	0.0385 (12)	0.0385 (12)	-0.0115 (9)	-0.0015 (9)	-0.0063 (9)
C5	0.0404 (12)	0.0403 (12)	0.0333 (11)	-0.0028 (10)	-0.0060 (9)	-0.0069 (9)
C6	0.0450 (13)	0.0434 (12)	0.0273 (11)	-0.0041 (10)	0.0026 (9)	0.0009 (9)
C7	0.0374 (12)	0.0412 (12)	0.0360 (12)	-0.0111 (9)	0.0028 (9)	0.0003 (9)
C8	0.0316 (11)	0.0358 (11)	0.0322 (11)	-0.0067 (8)	-0.0004 (8)	-0.0032 (8)

Geometric parameters (\AA , $^\circ$)

F—C6	1.328 (2)	C3—C4	1.391 (3)
N1—C1	1.354 (3)	C3—C8	1.401 (3)
N1—C2	1.371 (3)	N3—O3	1.208 (3)
N1—H1A	0.8600	N3—C5	1.462 (3)
O1—C2	1.222 (3)	C4—C5	1.367 (3)
C1—N2	1.284 (3)	C4—H4A	0.9300
C1—H1B	0.9300	C5—C6	1.401 (3)
N2—C8	1.381 (3)	C6—C7	1.361 (3)
C2—C3	1.455 (3)	C7—C8	1.394 (3)
O2—N3	1.211 (3)	C7—H7A	0.9300
C1—N1—C2	123.10 (18)	C5—C4—C3	119.7 (2)
C1—N1—H1A	118.4	C5—C4—H4A	120.1
C2—N1—H1A	118.4	C3—C4—H4A	120.1
N2—C1—N1	125.7 (2)	C4—C5—C6	119.8 (2)
N2—C1—H1B	117.2	C4—C5—N3	118.4 (2)
N1—C1—H1B	117.2	C6—C5—N3	121.8 (2)
C1—N2—C8	115.98 (18)	F—C6—C7	118.2 (2)
O1—C2—N1	121.96 (19)	F—C6—C5	120.7 (2)

O1—C2—C3	124.5 (2)	C7—C6—C5	121.1 (2)
N1—C2—C3	113.51 (19)	C6—C7—C8	120.0 (2)
C4—C3—C8	120.5 (2)	C6—C7—H7A	120.0
C4—C3—C2	120.43 (19)	C8—C7—H7A	120.0
C8—C3—C2	119.07 (19)	N2—C8—C7	118.51 (19)
O3—N3—O2	123.8 (2)	N2—C8—C3	122.58 (19)
O3—N3—C5	118.1 (2)	C7—C8—C3	118.91 (19)
O2—N3—C5	118.1 (2)		
C2—N1—C1—N2	0.6 (4)	O2—N3—C5—C6	-14.0 (4)
N1—C1—N2—C8	-0.8 (4)	C4—C5—C6—F	178.24 (19)
C1—N1—C2—O1	177.0 (2)	N3—C5—C6—F	-1.5 (4)
C1—N1—C2—C3	-1.4 (3)	C4—C5—C6—C7	-1.7 (4)
O1—C2—C3—C4	3.2 (4)	N3—C5—C6—C7	178.6 (2)
N1—C2—C3—C4	-178.5 (2)	F—C6—C7—C8	-179.23 (19)
O1—C2—C3—C8	-175.9 (2)	C5—C6—C7—C8	0.7 (4)
N1—C2—C3—C8	2.5 (3)	C1—N2—C8—C7	-178.6 (2)
C8—C3—C4—C5	0.5 (3)	C1—N2—C8—C3	2.0 (3)
C2—C3—C4—C5	-178.60 (19)	C6—C7—C8—N2	-178.6 (2)
C3—C4—C5—C6	1.1 (3)	C6—C7—C8—C3	0.8 (3)
C3—C4—C5—N3	-179.19 (19)	C4—C3—C8—N2	178.01 (19)
O3—N3—C5—C4	-12.3 (3)	C2—C3—C8—N2	-2.9 (3)
O2—N3—C5—C4	166.3 (2)	C4—C3—C8—C7	-1.4 (3)
O3—N3—C5—C6	167.4 (2)	C2—C3—C8—C7	177.65 (19)

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1A···O1 ⁱ	0.86	1.98	2.815 (2)	165
C1—H1B···O2 ⁱⁱ	0.93	2.47	3.396 (3)	179
C7—H7A···N2 ⁱⁱⁱ	0.93	2.50	3.422 (3)	171

Symmetry codes: (i) $-x, -y, -z+1$; (ii) $x+1, y, z+1$; (iii) $-x+1, -y+1, -z$.