

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

4-Benzenesulfonamidobenzoic acid

Hafiz Muhammad Adeel Sharif,^a Gui-Ying Dong,^b
Muhammad Nadeem Arshad^a and Islam Ullah Khan^{a*}

^aMaterials Chemistry Laboratory, Department of Chemistry, Government College University, Lahore 54000, Pakistan, and ^bCollege of Chemical Engineering and Biotechnology, Hebei Polytechnic University, Tangshan 063009, People's Republic of China

Correspondence e-mail: iukhangcu@126.com


Received 4 November 2009; accepted 9 November 2009

Key indicators: single-crystal X-ray study; $T = 295$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.065; wR factor = 0.199; data-to-parameter ratio = 18.5.

In the molecule of the title sulfonamide compound, $\text{C}_{13}\text{H}_{11}\text{NO}_4\text{S}$, the dihedral angle between the planes of the benzene ring and the carboxyl substituent group is $6.7(4)^\circ$. The two aromatic rings are inclined at $45.36(15)^\circ$ to one another. In the crystal, adjacent molecules are linked *via* classical intermolecular $\text{N}-\text{H}\cdots\text{O}$ and $\text{O}-\text{H}\cdots\text{O}$, and non-classical $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds, which stabilize the crystal structure.

Related literature

For the biological activity and pharmaceutical applications of sulfonamide derivatives, see: Innocenti *et al.* (2004); Parai *et al.* (2008); Rathish *et al.* (2009); Selvam *et al.* (2001). For related structures of sulfonamide derivatives with 4-aminobenzoic acid, see: Arshad *et al.* (2009); Khan *et al.* (2009); Nan & Xing (2006).



Experimental

Crystal data

$\text{C}_{13}\text{H}_{11}\text{NO}_4\text{S}$
 $M_r = 277.30$
Monoclinic, $P2_1/c$
 $a = 5.2050(3)$ Å
 $b = 37.726(2)$ Å

$c = 7.3781(4)$ Å
 $\beta = 117.510(3)^\circ$
 $V = 1284.98(13)$ Å³
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.26$ mm⁻¹
 $T = 295$ K

0.26 × 0.21 × 0.19 mm

Data collection

Bruker CCD diffractometer
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.935$, $T_{\max} = 0.958$

13550 measured reflections
3185 independent reflections
2633 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.199$
 $S = 1.10$
3185 reflections

172 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.38$ e Å⁻³
 $\Delta\rho_{\min} = -0.36$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N2}-\text{H2}\cdots\text{O8}^{\text{i}}$	0.81	2.28	3.054 (4)	162
$\text{O5}-\text{H5}\cdots\text{O6}^{\text{ii}}$	0.82	1.82	2.625 (3)	168
$\text{C18}-\text{H18}\cdots\text{O5}^{\text{iii}}$	0.93	2.58	3.413 (4)	150
$\text{C19}-\text{H19}\cdots\text{O6}^{\text{iv}}$	0.93	2.48	3.348 (4)	155

Symmetry codes: (i) $x + 1, y, z$; (ii) $-x - 1, -y + 1, -z - 1$; (iii) $-x - 1, -y + 1, -z$; (iv) $x, y, z + 1$.

Data collection: *SMART* (Bruker, 1998); cell refinement: *SAINT* (Bruker, 1999); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank Government College University and the Scientific Research Fund of Hebei Provincial Education Department (project 2006114) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2179).

References

- Arshad, M. N., Tahir, M. N., Khan, I. U., Siddiqui, W. A. & Shafiq, M. (2009). *Acta Cryst.* **E65**, o230.
Bruker (1998). *SMART*. Bruker AXS Inc., Madison, Wisconsin, USA.
Bruker (1999). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Innocenti, A., Antel, J., Wurl, M., Scozzafava, A. & Supuran, C. T. (2004). *Bioorg. Med. Chem. Lett.* **14**, 5703–5707.
Khan, I. U., Mustafa, G., Arshad, M. N., Shafiq, M. & Sharif, S. (2009). *Acta Cryst.* **E65**, o1073.
Nan, Z.-H. & Xing, J.-D. (2006). *Acta Cryst.* **E62**, o1978–o1979.
Parai, K. M., Panda, G., Srivastava, K. & Puri, S. K. (2008). *Bioorg. Med. Chem. Lett.* **18**, 776–781.
Rathish, I. G., Javed, K., Ahmad, S., Bano, S., Alam, M. S., Pillai, K. K., Singh, S. & Bagchi, V. (2009). *Bioorg. Med. Chem. Lett.* **19**, 255–258.
Selvam, P., Chandramohan, M., Clercq, E. D., Witvrouw, M. & Pannecouque, C. (2001). *Eur. J. Pharm. Sci.* **14**, 313–316.
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

supporting information

Acta Cryst. (2009). E65, o3086 [doi:10.1107/S1600536809047291]

4-Benzenesulfonamidobenzoic acid

Hafiz Muhammad Adeel Sharif, Gui-Ying Dong, Muhammad Nadeem Arshad and Islam Ullah Khan

S1. Comment

Benzene sulfonamide derivative have shown antimalarial (Parai *et al.*, 2008), carbonic anhydrase inhibitors (Innocenti *et al.*, 2004), antiHIV (Selvam *et al.*, 2001) and antiinflammatory (Rathish *et al.*, 2009) activities. In continuation of synthesis and structural studies of different benzene sulfonamide derivative (Khan *et al.*, 2009; Arshad *et al.*, 2009), we report here the molecular and crystal structures of title compound. The molecular structure of the title compound, **I**, is illustrated in Fig. 1. In **I**, phenyl and *p*-aminobenzoic acid moieties are connected through the SO₂ group. The structure of **I** is comparable with 4-(tosylamino)benzoic acid, (Nan & Xing, 2006). The dihedral angle between the planes of the benzene ring and the carboxyl substituent group is 6.7 (4)°. The two aromatic rings (C20–C25 and C14–C19) are inclined at 45.36 (15)° to one another. The torsion angle C14–N2–S1–C20 in the central part of the molecule is 70 (1)°.


In the crystal, adjacent molecules are linked *via* intermolecular classical N—H···O and O—H···O and non-classical C—H···O hydrogen bonds (Tab. 1, Fig. 2), which stabilize the crystal structure.

S2. Experimental


The 4-amino benzoic acid (1 g, 7.3 mmol) was dissolved in distilled water (10 ml). The pH of the solution was adjusted at 8–9 using 1M Na₂CO₃. Benzene sulfonylchloride (1.29 g, 7.3 mmol) was added to the above solution and stirred at room temperature until all the suspended benzene sulfonyl chloride was consumed. On completion of the reaction the pH was adjusted 1–2, using 1N HCl acid. The precipitate obtained was filtered, washed with distilled water, dried and recrystallized in methanol to yield colourless crystals.

S3. Refinement

All H atoms were positioned geometrically and refined using a riding model with C—H = 0.93 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aromatic, O—H = 0.82 Å and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ for the OH group and N—H = 0.81 Å and $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{N})$ for the NH group.

**Figure 1**

The molecular structure of **I** with the atom numbering scheme. Displacement ellipsoids are drawn at 30% probability level. H atoms are presented as a small spheres of arbitrary radius.

**Figure 2**

A partial packing diagram of the title compound. Hydrogen bonds are shown as dashed lines and H atoms not involved in hydrogen bonding omitted for clarity.

4-Benzenesulfonamidobenzoic acid

Crystal data

$C_{13}H_{11}NO_4S$

$M_r = 277.30$

Monoclinic, $P2_1/c$

Hall symbol: -P 2ybc

$a = 5.2050$ (3) Å
 $b = 37.726$ (2) Å
 $c = 7.3781$ (4) Å
 $\beta = 117.510$ (3)°
 $V = 1284.98$ (13) Å³
 $Z = 4$
 $F(000) = 576$
 $D_x = 1.433$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
 Cell parameters from 5008 reflections
 $\theta = 3.0\text{--}25^\circ$
 $\mu = 0.26$ mm⁻¹
 $T = 295$ K
 Block, colourless
 $0.26 \times 0.21 \times 0.19$ mm

Data collection

Bruker CCD
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.935$, $T_{\max} = 0.958$

13550 measured reflections
 3185 independent reflections
 2633 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$
 $\theta_{\max} = 28.4^\circ$, $\theta_{\min} = 1.1^\circ$
 $h = -4 \rightarrow 6$
 $k = -50 \rightarrow 45$
 $l = -9 \rightarrow 9$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.065$
 $wR(F^2) = 0.199$
 $S = 1.10$
 3185 reflections
 172 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0878P)^2 + 1.4857P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.38$ e Å⁻³
 $\Delta\rho_{\min} = -0.36$ e Å⁻³

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.27263 (17)	0.36345 (2)	0.53929 (12)	0.0443 (3)
O5	-0.4876 (5)	0.48980 (7)	-0.2675 (3)	0.0496 (6)
H5	-0.5699	0.5003	-0.3768	0.074*
O6	-0.2171 (5)	0.47077 (6)	-0.4087 (3)	0.0464 (5)
O7	0.4834 (6)	0.35320 (8)	0.7386 (4)	0.0693 (8)
O8	-0.0109 (5)	0.37334 (7)	0.5034 (4)	0.0557 (6)
N2	0.4098 (5)	0.39697 (7)	0.4738 (4)	0.0408 (6)
H2	0.5771	0.3942	0.5010	0.049*
C14	0.2360 (6)	0.41529 (7)	0.2848 (4)	0.0344 (6)

C15	0.2845 (6)	0.41056 (8)	0.1173 (5)	0.0411 (7)
H15	0.4269	0.3949	0.1242	0.049*
C16	0.1192 (6)	0.42930 (8)	-0.0615 (5)	0.0406 (6)
H16	0.1543	0.4267	-0.1735	0.049*
C17	-0.0980 (5)	0.45182 (7)	-0.0728 (4)	0.0319 (5)
C18	-0.1453 (6)	0.45608 (8)	0.0961 (4)	0.0375 (6)
H18	-0.2921	0.4710	0.0884	0.045*
C19	0.0239 (6)	0.43826 (8)	0.2754 (4)	0.0386 (6)
H19	-0.0049	0.4417	0.3895	0.046*
C20	0.2370 (7)	0.32889 (8)	0.3691 (5)	0.0456 (7)
C21	0.4682 (10)	0.30652 (11)	0.4110 (8)	0.0731 (12)
H21	0.6402	0.3090	0.5314	0.088*
C22	0.4401 (13)	0.28042 (13)	0.2715 (11)	0.0938 (18)
H22	0.5934	0.2650	0.2995	0.113*
C23	0.1881 (15)	0.27699 (14)	0.0918 (10)	0.0938 (17)
H23	0.1703	0.2594	-0.0015	0.113*
C24	-0.0327 (15)	0.29943 (14)	0.0524 (9)	0.1009 (19)
H24	-0.2033	0.2972	-0.0692	0.121*
C25	-0.0115 (10)	0.32565 (11)	0.1883 (7)	0.0724 (12)
H25	-0.1653	0.3411	0.1574	0.087*
C26	-0.2763 (6)	0.47182 (7)	-0.2629 (4)	0.0340 (6)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0400 (4)	0.0527 (5)	0.0429 (4)	0.0102 (3)	0.0214 (3)	0.0115 (3)
O5	0.0442 (12)	0.0658 (15)	0.0422 (11)	0.0254 (11)	0.0228 (10)	0.0118 (10)
O6	0.0505 (12)	0.0567 (13)	0.0380 (11)	0.0155 (10)	0.0255 (10)	0.0054 (9)
O7	0.0665 (17)	0.087 (2)	0.0466 (14)	0.0161 (15)	0.0193 (13)	0.0233 (13)
O8	0.0467 (13)	0.0653 (15)	0.0681 (16)	0.0086 (11)	0.0376 (12)	0.0058 (12)
N2	0.0281 (11)	0.0490 (14)	0.0410 (13)	0.0053 (10)	0.0121 (10)	0.0076 (10)
C14	0.0279 (12)	0.0365 (14)	0.0357 (13)	0.0003 (10)	0.0122 (10)	0.0004 (10)
C15	0.0350 (14)	0.0452 (16)	0.0460 (15)	0.0146 (12)	0.0210 (12)	0.0041 (12)
C16	0.0410 (15)	0.0483 (16)	0.0395 (14)	0.0095 (12)	0.0247 (12)	0.0008 (12)
C17	0.0277 (12)	0.0343 (13)	0.0342 (13)	0.0010 (10)	0.0146 (10)	-0.0026 (10)
C18	0.0361 (14)	0.0393 (14)	0.0421 (14)	0.0095 (11)	0.0223 (12)	0.0009 (11)
C19	0.0416 (15)	0.0437 (15)	0.0360 (14)	0.0073 (12)	0.0226 (12)	-0.0003 (11)
C20	0.0475 (17)	0.0407 (15)	0.0560 (18)	0.0068 (13)	0.0303 (15)	0.0121 (13)
C21	0.054 (2)	0.058 (2)	0.108 (4)	0.0152 (18)	0.038 (2)	0.003 (2)
C22	0.089 (4)	0.059 (3)	0.156 (6)	0.019 (2)	0.076 (4)	-0.001 (3)
C23	0.130 (5)	0.066 (3)	0.101 (4)	0.008 (3)	0.067 (4)	-0.011 (3)
C24	0.122 (5)	0.076 (3)	0.074 (3)	0.019 (3)	0.019 (3)	-0.015 (3)
C25	0.074 (3)	0.060 (2)	0.066 (2)	0.022 (2)	0.017 (2)	0.0007 (19)
C26	0.0307 (13)	0.0361 (13)	0.0346 (13)	0.0023 (10)	0.0146 (10)	-0.0031 (10)

Geometric parameters (Å, °)

S1—O7	1.423 (3)	C17—C26	1.482 (4)
S1—O8	1.423 (2)	C18—C19	1.379 (4)
S1—N2	1.632 (3)	C18—H18	0.9300
S1—C20	1.760 (4)	C19—H19	0.9300
O5—C26	1.279 (3)	C20—C25	1.368 (5)
O5—H5	0.8186	C20—C21	1.383 (5)
O6—C26	1.249 (3)	C21—C22	1.383 (7)
N2—C14	1.440 (3)	C21—H21	0.9300
N2—H2	0.8048	C22—C23	1.374 (8)
C14—C19	1.380 (4)	C22—H22	0.9300
C14—C15	1.382 (4)	C23—C24	1.347 (8)
C15—C16	1.390 (4)	C23—H23	0.9300
C15—H15	0.9300	C24—C25	1.376 (7)
C16—C17	1.386 (4)	C24—H24	0.9300
C16—H16	0.9300	C25—H25	0.9300
C17—C18	1.386 (4)		
O7—S1—O8	120.12 (17)	C14—C19—C18	119.7 (3)
O7—S1—N2	106.43 (16)	C14—C19—H19	120.1
O8—S1—N2	107.42 (14)	C18—C19—H19	120.1
O7—S1—C20	108.24 (17)	C25—C20—C21	119.8 (4)
O8—S1—C20	107.64 (16)	C25—C20—S1	120.2 (3)
N2—S1—C20	106.21 (14)	C21—C20—S1	119.9 (3)
C26—O5—H5	109.5	C20—C21—C22	119.1 (5)
C14—N2—S1	119.58 (19)	C20—C21—H21	120.4
C14—N2—H2	115.3	C22—C21—H21	120.5
S1—N2—H2	113.4	C23—C22—C21	120.7 (5)
C19—C14—C15	120.5 (3)	C23—C22—H22	119.6
C19—C14—N2	118.7 (3)	C21—C22—H22	119.6
C15—C14—N2	120.7 (2)	C24—C23—C22	119.2 (5)
C14—C15—C16	119.6 (3)	C24—C23—H23	120.4
C14—C15—H15	120.2	C22—C23—H23	120.4
C16—C15—H15	120.2	C23—C24—C25	121.4 (5)
C17—C16—C15	119.9 (3)	C23—C24—H24	119.3
C17—C16—H16	120.0	C25—C24—H24	119.3
C15—C16—H16	120.0	C20—C25—C24	119.7 (4)
C18—C17—C16	119.7 (3)	C20—C25—H25	120.1
C18—C17—C26	119.9 (2)	C24—C25—H25	120.1
C16—C17—C26	120.4 (2)	O6—C26—O5	123.0 (3)
C19—C18—C17	120.4 (3)	O6—C26—C17	120.2 (2)
C19—C18—H18	119.8	O5—C26—C17	116.8 (2)
C17—C18—H18	119.8		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N2—H2 \cdots O8 ⁱ	0.81	2.28	3.054 (4)	162
O5—H5 \cdots O6 ⁱⁱ	0.82	1.82	2.625 (3)	168
C18—H18 \cdots O5 ⁱⁱⁱ	0.93	2.58	3.413 (4)	150
C19—H19 \cdots O6 ^{iv}	0.93	2.48	3.348 (4)	155

Symmetry codes: (i) $x+1, y, z$; (ii) $-x-1, -y+1, -z-1$; (iii) $-x-1, -y+1, -z$; (iv) $x, y, z+1$.