

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

(4-Bromo-3,5-dimethyl-1H-pyrazol-1-yl)-(2,6-difluorophenyl)methanone

Xiang-Dong Mei, Yan-Hui Liang and Zhong-Yue Wang*

Key Laboratory of Pesticide Chemistry and Applications, Ministry of Agriculture, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, People's Republic of China

Correspondence e-mail: wangzhy0301@sina.com


Received 27 November 2009; accepted 7 December 2009

 Key indicators: single-crystal X-ray study; $T = 173$ K; mean $\sigma(\text{C}-\text{C}) = 0.010$ Å; R factor = 0.077; wR factor = 0.158; data-to-parameter ratio = 13.0.

There are two molecules in the asymmetric unit of the title compound, $\text{C}_{12}\text{H}_9\text{BrF}_2\text{N}_2\text{O}$. They have very similar conformations: the dihedral angles between their pyrazole and benzene ring systems are 78.4 (3) and 78.6 (4)°. In the crystal, weak aromatic $\pi-\pi$ stacking [centroid-centroid separation = 3.696 (5) Å] helps to establish the packing.

Related literature

For background to pyrazole derivatives in agrochemical and medicinal research, see: Sabbagh *et al.* (2009); Zheng *et al.* (2009).



Experimental

Crystal data

$\text{C}_{12}\text{H}_9\text{BrF}_2\text{N}_2\text{O}$
 $M_r = 315.12$
 Monoclinic, $P2_1/c$
 $a = 7.116$ (3) Å
 $b = 29.304$ (10) Å
 $c = 11.674$ (4) Å
 $\beta = 91.533$ (5)°

$V = 2433.5$ (15) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 3.39$ mm⁻¹
 $T = 173$ K
 $0.17 \times 0.17 \times 0.17$ mm

Data collection

Rigaku Saturn724+ CCD diffractometer
 Absorption correction: numerical (*CrystalClear*; Rigaku, 2008)
 $T_{\min} = 0.596$, $T_{\max} = 0.596$

15188 measured reflections
 4276 independent reflections
 3819 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.060$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.077$
 $wR(F^2) = 0.158$
 $S = 1.27$
 4276 reflections

329 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 1.08$ e Å⁻³
 $\Delta\rho_{\min} = -0.47$ e Å⁻³

Data collection: *CrystalClear* (Rigaku, 2008); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

This work was supported by National Technology System for the Grape Industry, the Major State Basic Research Development Program of China (No. 2006CB101907 and No. 2010CB126106), and the 863 high-tech key project of China (2006AA10A203).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: HB5260).

References

- Rigaku (2008). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
 Sabbagh, O. I., Raraka, M. M., Ibrahim, S. M., Pannecouque, C., Andrei, G., Snoeck, R., Balzarini, J. & Rashad, A. A. (2009). *Eur. J. Med. Chem.* **44**, 3746–3753.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Zheng, L. W., Wu, L. L., Zhao, B. X., Dong, W. L. & Miao, J. Y. (2009). *Bioorg. Med. Chem.* **17**, 1957–1962.

supporting information

Acta Cryst. (2010). E66, o115 [doi:10.1107/S1600536809052532]

(4-Bromo-3,5-dimethyl-1*H*-pyrazol-1-yl)(2,6-difluorophenyl)methanone

Xiang-Dong Mei, Yan-Hui Liang and Zhong-Yue Wang

S1. Experimental

The title compound (0.2 g) was dissolved in ethanol (50 ml) at room temperature. Colourless crystals of compound (I) were obtained through slow evaporation after two weeks.

S2. Refinement

The H atoms were placed at calculated positions, with C—H = 0.93–0.98 Å, and refined as riding with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{ep}}(\text{C})$.




Figure 1

The molecular structure of the title compound showing 50% probability displacement ellipsoids and atom-numbering scheme.

(4-Bromo-3,5-dimethyl-1*H*-pyrazol-1-yl)(2,6-difluorophenyl)methanone

Crystal data

$\text{C}_{12}\text{H}_9\text{BrF}_2\text{N}_2\text{O}$

$M_r = 315.12$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 7.116\ (3)\ \text{\AA}$

$b = 29.304\ (10)\ \text{\AA}$

$c = 11.674\ (4)\ \text{\AA}$

$\beta = 91.533\ (5)^\circ$

$V = 2433.5\ (15)\ \text{\AA}^3$

$Z = 8$

$F(000) = 1248$

$D_x = 1.720\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 768 reflections

$\theta = 2.2\text{--}27.5^\circ$

$\mu = 3.39\ \text{mm}^{-1}$

$T = 173\ \text{K}$

Block, colourless

$0.17 \times 0.17 \times 0.17\ \text{mm}$

Data collection

Rigaku Saturn724+ CCD
diffractometer

Radiation source: sealed tube

Graphite monochromator

ω scans at fixed $\chi = 45^\circ$

Absorption correction: numerical
(*CrystalClear*; Rigaku, 2008)

$T_{\min} = 0.596$, $T_{\max} = 0.596$

15188 measured reflections

4276 independent reflections

3819 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.060$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.8^\circ$

$h = -6 \rightarrow 8$

$k = -34 \rightarrow 34$

$l = -12 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.077$

$wR(F^2) = 0.158$

$S = 1.27$

4276 reflections

329 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0402P)^2 + 7.5951P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 1.08 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.47 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.48772 (10)	0.41014 (2)	0.89703 (6)	0.0409 (2)
Br2	1.00200 (10)	0.31950 (3)	1.09739 (6)	0.0491 (3)
F1	0.7911 (6)	0.30667 (16)	1.4485 (4)	0.0586 (12)
F2	0.1573 (5)	0.32948 (15)	1.3484 (3)	0.0503 (11)
F3	0.6692 (7)	0.42324 (19)	0.6390 (5)	0.0780 (16)
F4	1.3057 (7)	0.4580 (2)	0.6345 (6)	0.0889 (18)
O1	0.5532 (8)	0.26916 (16)	1.2410 (4)	0.0534 (14)
O2	1.0003 (10)	0.48031 (17)	0.8138 (4)	0.0649 (17)
N1	0.5031 (7)	0.34346 (17)	1.1966 (4)	0.0298 (12)
N2	0.4946 (7)	0.38788 (17)	1.2406 (5)	0.0322 (12)
N3	0.9873 (7)	0.40262 (16)	0.8255 (4)	0.0290 (12)
N4	0.9923 (7)	0.36189 (17)	0.7664 (4)	0.0303 (12)
C1	0.4918 (9)	0.4144 (2)	1.1501 (6)	0.0323 (15)
C2	0.4947 (9)	0.3874 (2)	1.0474 (5)	0.0304 (14)
C3	0.5023 (8)	0.3431 (2)	1.0775 (5)	0.0287 (14)
C4	0.4855 (11)	0.4650 (2)	1.1623 (6)	0.0431 (17)

H4A	0.4898	0.4731	1.2438	0.065*
H4B	0.3690	0.4767	1.1265	0.065*
H4C	0.5936	0.4785	1.1247	0.065*
C5	0.5039 (10)	0.3009 (2)	1.0060 (6)	0.0380 (16)
H5A	0.4885	0.3092	0.9250	0.057*
H5B	0.4005	0.2809	1.0279	0.057*
H5C	0.6238	0.2850	1.0183	0.057*
C6	0.5157 (10)	0.3064 (2)	1.2725 (5)	0.0351 (15)
C7	0.4758 (9)	0.3177 (2)	1.3954 (5)	0.0302 (14)
C8	0.6131 (9)	0.3157 (2)	1.4813 (6)	0.0362 (16)
C9	0.5761 (11)	0.3226 (2)	1.5946 (6)	0.0434 (18)
H9A	0.6742	0.3219	1.6513	0.052*
C10	0.3931 (10)	0.3306 (2)	1.6239 (6)	0.0387 (16)
H10A	0.3650	0.3341	1.7025	0.046*
C11	0.2501 (10)	0.3337 (2)	1.5433 (6)	0.0409 (17)
H11A	0.1250	0.3405	1.5642	0.049*
C12	0.2950 (9)	0.3266 (2)	1.4315 (6)	0.0355 (15)
C13	0.9966 (9)	0.3303 (2)	0.8459 (6)	0.0330 (15)
C14	0.9948 (9)	0.3504 (2)	0.9562 (5)	0.0325 (15)
C15	0.9901 (9)	0.3960 (2)	0.9434 (5)	0.0339 (15)
C16	1.0047 (12)	0.2810 (2)	0.8158 (7)	0.051 (2)
H16A	1.0217	0.2777	0.7332	0.076*
H16B	0.8872	0.2661	0.8369	0.076*
H16C	1.1105	0.2666	0.8575	0.076*
C17	0.9875 (11)	0.4331 (3)	1.0304 (6)	0.0481 (19)
H17A	0.9903	0.4197	1.1074	0.072*
H17B	0.8729	0.4512	1.0195	0.072*
H17C	1.0977	0.4527	1.0217	0.072*
C18	0.9919 (10)	0.4439 (2)	0.7649 (6)	0.0378 (16)
C19	0.9876 (10)	0.4396 (2)	0.6374 (6)	0.0374 (16)
C20	0.8243 (12)	0.4300 (3)	0.5776 (7)	0.052 (2)
C21	0.8162 (19)	0.4272 (3)	0.4604 (8)	0.087 (4)
H21A	0.7018	0.4200	0.4206	0.105*
C22	0.975 (2)	0.4350 (3)	0.4030 (8)	0.100 (5)
H22A	0.9711	0.4331	0.3217	0.120*
C23	1.141 (2)	0.4456 (4)	0.4571 (10)	0.090 (4)
H23A	1.2508	0.4512	0.4148	0.108*
C24	1.1454 (13)	0.4478 (3)	0.5754 (7)	0.058 (2)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0507 (5)	0.0423 (4)	0.0296 (4)	-0.0035 (3)	0.0003 (3)	0.0051 (3)
Br2	0.0450 (4)	0.0693 (6)	0.0328 (4)	-0.0021 (4)	-0.0021 (3)	0.0171 (4)
F1	0.036 (2)	0.073 (3)	0.067 (3)	0.004 (2)	0.001 (2)	-0.013 (2)
F2	0.039 (2)	0.071 (3)	0.041 (2)	0.006 (2)	-0.0080 (19)	-0.004 (2)
F3	0.055 (3)	0.076 (4)	0.102 (4)	-0.014 (3)	-0.013 (3)	0.008 (3)
F4	0.049 (3)	0.093 (4)	0.126 (5)	-0.006 (3)	0.018 (3)	-0.001 (4)

O1	0.095 (4)	0.021 (2)	0.044 (3)	0.013 (3)	0.016 (3)	0.000 (2)
O2	0.124 (5)	0.029 (3)	0.042 (3)	0.000 (3)	0.002 (3)	-0.002 (2)
N1	0.036 (3)	0.027 (3)	0.026 (3)	-0.002 (2)	0.001 (2)	-0.003 (2)
N2	0.040 (3)	0.024 (3)	0.033 (3)	0.000 (2)	0.002 (2)	0.000 (2)
N3	0.037 (3)	0.023 (3)	0.027 (3)	0.001 (2)	0.000 (2)	-0.003 (2)
N4	0.039 (3)	0.025 (3)	0.027 (3)	-0.003 (2)	0.001 (2)	-0.005 (2)
C1	0.039 (4)	0.025 (3)	0.033 (4)	-0.001 (3)	0.004 (3)	0.000 (3)
C2	0.040 (4)	0.028 (3)	0.024 (3)	0.001 (3)	0.007 (3)	0.001 (3)
C3	0.029 (3)	0.028 (3)	0.029 (3)	-0.001 (3)	0.004 (3)	-0.002 (3)
C4	0.058 (5)	0.027 (4)	0.044 (4)	-0.007 (3)	-0.001 (4)	0.000 (3)
C5	0.047 (4)	0.035 (4)	0.032 (4)	-0.005 (3)	0.001 (3)	-0.006 (3)
C6	0.044 (4)	0.035 (4)	0.027 (4)	0.002 (3)	0.004 (3)	0.002 (3)
C7	0.041 (4)	0.020 (3)	0.030 (4)	0.000 (3)	0.002 (3)	0.000 (3)
C8	0.034 (4)	0.027 (3)	0.047 (4)	0.002 (3)	0.001 (3)	-0.004 (3)
C9	0.056 (5)	0.043 (4)	0.031 (4)	-0.006 (3)	-0.011 (3)	0.000 (3)
C10	0.053 (5)	0.029 (4)	0.033 (4)	0.002 (3)	-0.004 (3)	0.005 (3)
C11	0.038 (4)	0.040 (4)	0.045 (4)	-0.001 (3)	0.010 (3)	0.000 (3)
C12	0.036 (4)	0.037 (4)	0.034 (4)	-0.003 (3)	-0.001 (3)	-0.007 (3)
C13	0.036 (4)	0.025 (3)	0.038 (4)	-0.003 (3)	-0.002 (3)	0.000 (3)
C14	0.033 (4)	0.034 (4)	0.030 (4)	-0.003 (3)	0.001 (3)	0.002 (3)
C15	0.040 (4)	0.040 (4)	0.022 (3)	0.004 (3)	0.000 (3)	-0.004 (3)
C16	0.072 (5)	0.032 (4)	0.048 (5)	-0.010 (4)	0.002 (4)	0.005 (3)
C17	0.063 (5)	0.051 (5)	0.030 (4)	0.002 (4)	0.004 (4)	-0.014 (3)
C18	0.053 (4)	0.027 (4)	0.034 (4)	0.009 (3)	0.003 (3)	-0.004 (3)
C19	0.055 (4)	0.021 (3)	0.036 (4)	0.006 (3)	-0.002 (3)	0.003 (3)
C20	0.067 (6)	0.037 (4)	0.052 (5)	-0.003 (4)	-0.008 (4)	0.008 (4)
C21	0.167 (12)	0.046 (5)	0.047 (6)	-0.006 (6)	-0.051 (7)	0.008 (4)
C22	0.231 (18)	0.044 (6)	0.025 (5)	0.015 (8)	0.011 (8)	0.012 (4)
C23	0.147 (12)	0.062 (7)	0.065 (7)	0.016 (7)	0.052 (7)	0.016 (6)
C24	0.072 (6)	0.048 (5)	0.055 (5)	0.000 (4)	0.013 (5)	0.003 (4)

Geometric parameters (Å, °)

Br1—C2	1.877 (6)	C7—C12	1.389 (9)
Br2—C14	1.881 (6)	C8—C9	1.371 (10)
F1—C8	1.358 (8)	C9—C10	1.375 (10)
F2—C12	1.363 (8)	C9—H9A	0.9500
F3—C20	1.347 (9)	C10—C11	1.370 (10)
F4—C24	1.351 (10)	C10—H10A	0.9500
O1—C6	1.185 (8)	C11—C12	1.368 (9)
O2—C18	1.210 (8)	C11—H11A	0.9500
N1—C3	1.390 (8)	C13—C14	1.416 (9)
N1—N2	1.401 (7)	C13—C16	1.489 (9)
N1—C6	1.403 (8)	C14—C15	1.345 (9)
N2—C1	1.311 (8)	C15—C17	1.487 (9)
N3—N4	1.380 (7)	C16—H16A	0.9800
N3—C15	1.390 (8)	C16—H16B	0.9800
N3—C18	1.403 (8)	C16—H16C	0.9800

N4—C13	1.311 (8)	C17—H17A	0.9800
C1—C2	1.438 (9)	C17—H17B	0.9800
C1—C4	1.491 (8)	C17—H17C	0.9800
C2—C3	1.346 (9)	C18—C19	1.494 (9)
C3—C5	1.491 (8)	C19—C20	1.369 (10)
C4—H4A	0.9800	C19—C24	1.373 (11)
C4—H4B	0.9800	C20—C21	1.371 (12)
C4—H4C	0.9800	C21—C22	1.350 (16)
C5—H5A	0.9800	C21—H21A	0.9500
C5—H5B	0.9800	C22—C23	1.358 (17)
C5—H5C	0.9800	C22—H22A	0.9500
C6—C7	1.506 (8)	C23—C24	1.382 (13)
C7—C8	1.382 (9)	C23—H23A	0.9500
C3—N1—N2	112.0 (5)	C10—C11—H11A	121.4
C3—N1—C6	128.7 (5)	F2—C12—C11	119.2 (6)
N2—N1—C6	119.3 (5)	F2—C12—C7	116.7 (6)
C1—N2—N1	104.8 (5)	C11—C12—C7	124.1 (6)
N4—N3—C15	112.0 (5)	N4—C13—C14	110.5 (5)
N4—N3—C18	119.5 (5)	N4—C13—C16	121.2 (6)
C15—N3—C18	128.3 (5)	C14—C13—C16	128.3 (6)
C13—N4—N3	104.9 (5)	C15—C14—C13	108.3 (6)
N2—C1—C2	110.2 (5)	C15—C14—Br2	125.1 (5)
N2—C1—C4	120.9 (6)	C13—C14—Br2	126.6 (5)
C2—C1—C4	128.9 (6)	C14—C15—N3	104.4 (5)
C3—C2—C1	108.3 (5)	C14—C15—C17	130.6 (6)
C3—C2—Br1	125.9 (5)	N3—C15—C17	125.1 (6)
C1—C2—Br1	125.7 (5)	C13—C16—H16A	109.5
C2—C3—N1	104.7 (5)	C13—C16—H16B	109.5
C2—C3—C5	130.8 (6)	H16A—C16—H16B	109.5
N1—C3—C5	124.5 (6)	C13—C16—H16C	109.5
C1—C4—H4A	109.5	H16A—C16—H16C	109.5
C1—C4—H4B	109.5	H16B—C16—H16C	109.5
H4A—C4—H4B	109.5	C15—C17—H17A	109.5
C1—C4—H4C	109.5	C15—C17—H17B	109.5
H4A—C4—H4C	109.5	H17A—C17—H17B	109.5
H4B—C4—H4C	109.5	C15—C17—H17C	109.5
C3—C5—H5A	109.5	H17A—C17—H17C	109.5
C3—C5—H5B	109.5	H17B—C17—H17C	109.5
H5A—C5—H5B	109.5	O2—C18—N3	121.6 (6)
C3—C5—H5C	109.5	O2—C18—C19	123.0 (6)
H5A—C5—H5C	109.5	N3—C18—C19	115.4 (5)
H5B—C5—H5C	109.5	C20—C19—C24	117.6 (7)
O1—C6—N1	121.9 (6)	C20—C19—C18	121.3 (7)
O1—C6—C7	123.2 (6)	C24—C19—C18	121.0 (7)
N1—C6—C7	114.9 (5)	F3—C20—C19	117.2 (7)
C8—C7—C12	115.5 (6)	F3—C20—C21	120.6 (9)
C8—C7—C6	122.2 (6)	C19—C20—C21	122.2 (9)

C12—C7—C6	122.0 (6)	C22—C21—C20	118.1 (10)
F1—C8—C9	120.4 (6)	C22—C21—H21A	120.9
F1—C8—C7	116.8 (6)	C20—C21—H21A	120.9
C9—C8—C7	122.9 (6)	C21—C22—C23	122.5 (10)
C8—C9—C10	118.3 (7)	C21—C22—H22A	118.8
C8—C9—H9A	120.9	C23—C22—H22A	118.8
C10—C9—H9A	120.9	C22—C23—C24	118.1 (11)
C11—C10—C9	122.1 (7)	C22—C23—H23A	120.9
C11—C10—H10A	119.0	C24—C23—H23A	120.9
C9—C10—H10A	119.0	F4—C24—C19	117.4 (8)
C12—C11—C10	117.2 (7)	F4—C24—C23	121.2 (9)
C12—C11—H11A	121.4	C19—C24—C23	121.4 (10)
C3—N1—N2—C1	-1.0 (7)	C8—C7—C12—C11	1.0 (10)
C6—N1—N2—C1	177.4 (6)	C6—C7—C12—C11	175.0 (6)
C15—N3—N4—C13	0.5 (7)	N3—N4—C13—C14	-0.1 (7)
C18—N3—N4—C13	176.6 (6)	N3—N4—C13—C16	-179.4 (6)
N1—N2—C1—C2	1.1 (7)	N4—C13—C14—C15	-0.3 (8)
N1—N2—C1—C4	-179.2 (6)	C16—C13—C14—C15	178.9 (7)
N2—C1—C2—C3	-0.9 (8)	N4—C13—C14—Br2	-179.3 (5)
C4—C1—C2—C3	179.4 (7)	C16—C13—C14—Br2	-0.1 (11)
N2—C1—C2—Br1	178.8 (5)	C13—C14—C15—N3	0.6 (7)
C4—C1—C2—Br1	-0.9 (11)	Br2—C14—C15—N3	179.6 (4)
C1—C2—C3—N1	0.3 (7)	C13—C14—C15—C17	-179.6 (7)
Br1—C2—C3—N1	-179.4 (4)	Br2—C14—C15—C17	-0.6 (11)
C1—C2—C3—C5	178.5 (6)	N4—N3—C15—C14	-0.7 (7)
Br1—C2—C3—C5	-1.2 (11)	C18—N3—C15—C14	-176.4 (6)
N2—N1—C3—C2	0.4 (7)	N4—N3—C15—C17	179.5 (6)
C6—N1—C3—C2	-177.8 (6)	C18—N3—C15—C17	3.8 (11)
N2—N1—C3—C5	-177.9 (6)	N4—N3—C18—O2	-175.1 (7)
C6—N1—C3—C5	3.8 (10)	C15—N3—C18—O2	0.3 (11)
C3—N1—C6—O1	10.4 (11)	N4—N3—C18—C19	4.4 (9)
N2—N1—C6—O1	-167.7 (6)	C15—N3—C18—C19	179.8 (6)
C3—N1—C6—C7	-168.9 (6)	O2—C18—C19—C20	-103.7 (9)
N2—N1—C6—C7	13.0 (8)	N3—C18—C19—C20	76.9 (8)
O1—C6—C7—C8	67.7 (10)	O2—C18—C19—C24	72.5 (10)
N1—C6—C7—C8	-113.0 (7)	N3—C18—C19—C24	-107.0 (8)
O1—C6—C7—C12	-105.9 (8)	C24—C19—C20—F3	-178.2 (7)
N1—C6—C7—C12	73.4 (8)	C18—C19—C20—F3	-2.0 (10)
C12—C7—C8—F1	179.5 (6)	C24—C19—C20—C21	1.8 (11)
C6—C7—C8—F1	5.5 (9)	C18—C19—C20—C21	178.1 (7)
C12—C7—C8—C9	-1.1 (9)	F3—C20—C21—C22	179.0 (8)
C6—C7—C8—C9	-175.1 (6)	C19—C20—C21—C22	-1.1 (13)
F1—C8—C9—C10	-178.6 (6)	C20—C21—C22—C23	-0.2 (16)
C7—C8—C9—C10	2.0 (10)	C21—C22—C23—C24	0.6 (16)
C8—C9—C10—C11	-2.8 (10)	C20—C19—C24—F4	179.0 (7)
C9—C10—C11—C12	2.6 (10)	C18—C19—C24—F4	2.7 (11)
C10—C11—C12—F2	-179.8 (6)	C20—C19—C24—C23	-1.3 (12)

supporting information

C10—C11—C12—C7	-1.7 (10)	C18—C19—C24—C23	-177.6 (8)
C8—C7—C12—F2	179.0 (6)	C22—C23—C24—F4	179.9 (9)
C6—C7—C12—F2	-7.0 (9)	C22—C23—C24—C19	0.2 (14)
