

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

meso-1-[[2-(Propyl-1-sulfinyl)ethyl]-sulfinyl]propane

 Solange M. S. V. Wardell,^a James L. Wardell,^{b‡}
 Geraldo M. de Lima^c and Edward R. T. Tiekink^{d*}

^aCHEMSOL, 1 Harcourt Road, Aberdeen AB15 5NY, Scotland, ^bCentro de Desenvolvimento Tecnológico em Saúde (CDTS), Fundação Oswaldo Cruz (FIOCRUZ), Casa Amarela, Campus de Manguinhos, Av. Brasil 4365, 21040-900, Rio de Janeiro, RJ, Brazil, ^cDepartamento de Química, ICEx, Universidade Federal de Minas Gerais, 31270-901 Belo Horizonte, MG, Brazil, and ^dDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
 Correspondence e-mail: Edward.Tiekink@gmail.com


Received 11 February 2010; accepted 11 February 2010

Key indicators: single-crystal X-ray study; $T = 120$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.031; wR factor = 0.080; data-to-parameter ratio = 22.1.

The title molecule, $\text{C}_8\text{H}_{18}\text{O}_2\text{S}_2$, is disposed about a centre of inversion implying an *anti*-disposition of the sulfinyl-O atoms; the terminal *n*-propyl group has an extended conformation. The crystal packing is dominated by $\text{C}-\text{H}\cdots\text{O}$ interactions, which lead to the formation of supramolecular arrays in the *bc* plane.

Related literature

For the structures of the stereoisomers of $RS(=\text{O})\text{CH}_2\text{CH}_2\text{-S}(=\text{O})R$, see: Pelizzi *et al.* (1976); Svinning *et al.* (1976); Chu & Madden (1978); Ternay *et al.* (1978); Cattalini *et al.* (1979); Li *et al.* (2002, 2004). For the preparation and separation of the stereoisomers of the title compound, see: Hull & Bargar (1975); Li *et al.* (2005). For information on the use of bis-sulfoxides as a ligand, see: de Souza *et al.* (1995, 1997); Huang *et al.* (1986); Huang & Zhang (1986); Filgueiras & Marques (1985); Filgueiras *et al.* (1982); Bu *et al.* (2002); Li *et al.* (2005); Yapp *et al.* (1997).



Experimental

Crystal data

$\text{C}_8\text{H}_{18}\text{O}_2\text{S}_2$
 $M_r = 210.34$

Monoclinic, $P2_1/c$
 $a = 11.9794$ (9) Å

‡ Additional correspondence author, e-mail: j.wardell@abdn.ac.uk.

$b = 5.2190$ (3) Å
 $c = 8.7618$ (5) Å
 $\beta = 97.191$ (5)°
 $V = 543.48$ (6) Å³
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 0.45$ mm⁻¹
 $T = 120$ K
 $1.1 \times 0.6 \times 0.12$ mm

Data collection

Nonius KappaCCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 2007)
 $T_{\min} = 0.527$, $T_{\max} = 0.746$

5666 measured reflections
 1239 independent reflections
 1167 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.042$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.031$
 $wR(F^2) = 0.080$
 $S = 1.06$
 1239 reflections

56 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.33$ e Å⁻³
 $\Delta\rho_{\min} = -0.35$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C1}-\text{H1b}\cdots\text{O1}^i$	0.99	2.41	3.3035 (19)	150
$\text{C4}-\text{H4a}\cdots\text{O1}^i$	0.99	2.40	3.2751 (19)	147
$\text{C4}-\text{H4b}\cdots\text{O1}^{ii}$	0.99	2.57	3.5124 (18)	159

Symmetry codes: (i) $x, y + 1, z$; (ii) $-x + 1, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: COLLECT (Hooft, 1998); cell refinement: DENZO (Otwinowski & Minor, 1997) and COLLECT; data reduction: DENZO and COLLECT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 (Farrugia, 1997) and DIAMOND (Brandenburg, 2006); software used to prepare material for publication: publCIF (Westrip, 2010).

The use of the EPSRC X-ray crystallographic service at the University of Southampton, England, and the valuable assistance of the staff there is gratefully acknowledged. JLW acknowledges support from CAPES and FAPEMIG (Brazil).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: EZZ2201).

References

- Brandenburg, K. (2006). DIAMOND. Crystal Impact GbR, Bonn, Germany.
 Bu, X.-H., Weng, W., Li, J.-R., Chen, W. & Zhang, R.-H. (2002). *Inorg. Chem.* **41**, 413–415.
 Cattalini, L., Michelon, G., Marangoni, G. & Pelizzi, G. (1979). *J. Chem. Soc. Dalton Trans.* pp. 96–101.
 Chu, S. S. C. & Madden, J. (1978). *Acta Cryst.* **B34**, 841–845.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Filgueiras, C. A. L., Celso, C., Marques, E. V. & Johnson, B. F. G. (1982). *Inorg. Chim. Acta*, **59**, 71–74.
 Filgueiras, C. A. L. & Marques, E. V. (1985). *Transition Met. Chem.* **10**, 241–243.
 Hooft, R. W. W. (1998). COLLECT. Nonius BV, Delft, The Netherlands.
 Huang, Z. & Zhang, Z. (1986). *Wuji Huaxue Xuebao*, **2**, 64–71.
 Huang, Z., Zhang, R., Wang, J. & Zhang, Z. (1986). *Wuji Huaxue Xuebao*, **2**, 54–59.
 Hull, C. M. & Bargar, T. W. (1975). *J. Org. Chem.* **40**, 3152–3154.
 Li, J.-R., Bu, X.-H. & Zhang, R.-H. (2004). *Inorg. Chem.* **43**, 237–244.
 Li, J.-R., Bu, X.-H., Zhang, R.-H. & Ribas, J. (2005). *Cryst. Growth Des.* **5**, 1919–1932.
 Li, J.-R., Zhang, R.-H. & Bu, X.-H. (2002). *Acta Cryst.* **E58**, o911–o912.

- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Pelizzi, G., Michelon, G. & Bonivento, M. (1976). *Cryst. Struct. Commun.* **5**, 617–620.
- Sheldrick, G. M. (2007). *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Souza, G. F. de, Filgueiras, C. A. L., Abras, A., Carvalho, C. C., Francisco, R. H. P. & Gambardella, M. T. Do. P. (1995). *An. Ass. Bras. Quim.* **44**, 42–46.
- Souza, G. F. de, Filgueiras, C. A. L., Hitchcock, P. B. & Nixon, J. F. (1997). *Inorg. Chim. Acta*, **261**, 221–225.
- Svinning, T., Mo, F. & Bruun, T. (1976). *Acta Cryst.* **B32**, 759–764.
- Ternay, A. L. Jr, Lin, J., Sutliff, T., Chu, S. S. C. & Chung, B. (1978). *J. Org. Chem.* **43**, 3024–3031.
- Westrip, S. P. (2010). *publCIF*. In preparation.
- Yapp, D. T. T., Rettig, S. J., James, B. R. & Skov, K. A. (1997). *Inorg. Chem.* **36**, 5635–5641.

supporting information

Acta Cryst. (2010). E66, o647–o648 [doi:10.1107/S1600536810005799]

meso-1-[[2-(Propyl-1-sulfinyl)ethyl]sulfinyl]propane

Solange M. S. V. Wardell, James L. Wardell, Geraldo M. de Lima and Edward R. T. Tiekink

S1. Comment

Bis-sulfoxides such as the title compound, (I), have found use as ligands (de Souza *et al.*, 1995, 1997; Huang *et al.*, 1986; Huang & Zhang, 1986; Filgueiras & Marques, 1985; Filgueiras *et al.*, 1982; Bu *et al.*, 2002; Li *et al.*, 2005; Yapp *et al.*, 1997). Details on the preparation and the separation of stereoisomers of (I) are available in the literature (Hull & Bargar 1975; Li *et al.* 2005). Crystallography shows the molecule of (I) is disposed about a centre of inversion, Fig. 1, and the n-propyl chain has an extended conformation as seen in the value of the S1–C1–C2–C3 torsion angle of S1–C1–C2–C3 of $-179.93(11)^\circ$. The *anti*-disposition of the sulfinyl-O atoms allow for the optimisation of C–H \cdots O interactions in the crystal structure, Table 1. Thus, each sulfinyl-O1 associates with three methylene-H atoms to form a supramolecular array in the *bc* plane, Fig. 2, which stack along the *a* axis, Fig. 3.

The central core in (I), including its disposition about a centre of inversion, resembles that found in each of the reported *meso*-RS(=O)CH₂CH₂S(=O)R structures, where R = Me (Svinning *et al.*, 1976), Et (Li *et al.*, 2004), Ph (Pelizzi, *et al.* 1976; Ternay *et al.* 1978; Cattalini *et al.*, 1979), benzyl (Li *et al.* 2002), and mesityl (Chu & Madden, 1978), but not in their homo chiral stereoisomers where R = Ph (Ternay *et al.* 1978; Cattalini *et al.*, 1979) and mesityl (Chu & Madden, 1978).

S2. Experimental

Compound (I) was prepared by a published method (Li *et al.*, 2005) and had spectral and other parameters in agreement with published values (Li *et al.*, 2005; Yapp *et al.*, 1997). M.pt. 434–435 K. Lit. value 434–435 K (Hull & Bargar, 1975; Li *et al.*, 2005). The sample used in the crystallographic study was grown from an ethanol solution of (I).

S3. Refinement

The C-bound H atoms were geometrically placed (C–H = 0.98–0.99 Å) and refined as riding with $U_{iso}(H) = 1.2-1.5U_{eq}(C)$.




Figure 1

The molecular structure of (I) showing the atom-labelling scheme and displacement ellipsoids at the 70% probability level.




Figure 2

A view of a supramolecular array in (I) in the *bc* plane. The C–H···O interactions are shown as blue dashed lines. Colour code: S, yellow; O, red; C, grey; and H, green.




Figure 3

View in projection down the *c* axis of the unit cell contents in (I). Colour code: S, yellow; O, red; C, grey; and H, green.

meso-1-[[2-(Propyl-1-sulfinyl)ethyl]sulfinyl]propane

Crystal data

$C_8H_{18}O_2S_2$

$M_r = 210.34$

Monoclinic, $P2_1/c$

Hall symbol: $-P\ 2_1/c$

$a = 11.9794(9)\ \text{\AA}$

$b = 5.2190(3)\ \text{\AA}$

$c = 8.7618(5)\ \text{\AA}$

$\beta = 97.191(5)^\circ$

$V = 543.48(6)\ \text{\AA}^3$

$Z = 2$

$F(000) = 228$

$D_x = 1.285\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 19160 reflections

$\theta = 2.9\text{--}27.5^\circ$

$\mu = 0.45\ \text{mm}^{-1}$

$T = 120$ K $1.1 \times 0.6 \times 0.12$ mm
 Plate, colourless

Data collection

Nonius KappaCCD area-detector diffractometer	$T_{\min} = 0.527$, $T_{\max} = 0.746$ 5666 measured reflections
Radiation source: Enraf Nonius FR591 rotating anode	1239 independent reflections 1167 reflections with $I > 2\sigma(I)$
10 cm confocal mirrors monochromator	$R_{\text{int}} = 0.042$
Detector resolution: 9.091 pixels mm^{-1}	$\theta_{\max} = 27.5^\circ$, $\theta_{\min} = 3.4^\circ$
φ and ω scans	$h = -12 \rightarrow 15$
Absorption correction: multi-scan (SADABS; Sheldrick, 2007)	$k = -6 \rightarrow 6$ $l = -11 \rightarrow 11$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.031$	H-atom parameters constrained
$wR(F^2) = 0.080$	$w = 1/[\sigma^2(F_o^2) + (0.0344P)^2 + 0.4193P]$
$S = 1.06$	where $P = (F_o^2 + 2F_c^2)/3$
1239 reflections	$(\Delta/\sigma)_{\max} = 0.001$
56 parameters	$\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.35 \text{ e } \text{\AA}^{-3}$
Primary atom site location: structure-invariant direct methods	

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.66819 (3)	-0.09909 (6)	0.45405 (4)	0.01305 (14)
O1	0.64458 (10)	-0.37397 (19)	0.40786 (13)	0.0187 (3)
C1	0.72083 (12)	0.0560 (3)	0.29375 (16)	0.0149 (3)
H1A	0.6705	0.0192	0.1978	0.018*
H1B	0.7228	0.2438	0.3097	0.018*
C2	0.83896 (12)	-0.0425 (3)	0.27975 (17)	0.0191 (3)
H2A	0.8363	-0.2305	0.2646	0.023*
H2B	0.8885	-0.0065	0.3765	0.023*
C3	0.88818 (14)	0.0828 (3)	0.14553 (19)	0.0246 (4)
H3A	0.8398	0.0455	0.0494	0.037*
H3B	0.9638	0.0147	0.1397	0.037*
H3C	0.8925	0.2686	0.1614	0.037*
C4	0.53341 (12)	0.0583 (3)	0.44006 (15)	0.0144 (3)
H4A	0.5438	0.2443	0.4589	0.017*

H4B	0.4924	0.0343	0.3358	0.017*
-----	--------	--------	--------	--------

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0150 (2)	0.0110 (2)	0.0135 (2)	0.00060 (11)	0.00307 (13)	0.00048 (11)
O1	0.0238 (6)	0.0087 (5)	0.0244 (6)	0.0012 (4)	0.0066 (4)	0.0004 (4)
C1	0.0176 (7)	0.0133 (6)	0.0147 (6)	-0.0002 (5)	0.0048 (5)	0.0007 (5)
C2	0.0179 (7)	0.0221 (7)	0.0181 (7)	0.0014 (6)	0.0057 (5)	0.0005 (6)
C3	0.0210 (8)	0.0325 (9)	0.0217 (7)	-0.0031 (6)	0.0078 (6)	0.0010 (6)
C4	0.0167 (7)	0.0111 (6)	0.0162 (6)	0.0018 (5)	0.0049 (5)	0.0013 (5)

Geometric parameters (Å, °)

S1—O1	1.5077 (10)	C2—H2B	0.9900
S1—C4	1.8018 (14)	C3—H3A	0.9800
S1—C1	1.8021 (14)	C3—H3B	0.9800
C1—C2	1.525 (2)	C3—H3C	0.9800
C1—H1A	0.9900	C4—C4 ⁱ	1.525 (3)
C1—H1B	0.9900	C4—H4A	0.9900
C2—C3	1.527 (2)	C4—H4B	0.9900
C2—H2A	0.9900		
O1—S1—C4	106.20 (7)	H2A—C2—H2B	107.9
O1—S1—C1	106.88 (7)	C2—C3—H3A	109.5
C4—S1—C1	98.06 (6)	C2—C3—H3B	109.5
C2—C1—S1	109.27 (10)	H3A—C3—H3B	109.5
C2—C1—H1A	109.8	C2—C3—H3C	109.5
S1—C1—H1A	109.8	H3A—C3—H3C	109.5
C2—C1—H1B	109.8	H3B—C3—H3C	109.5
S1—C1—H1B	109.8	C4 ⁱ —C4—S1	108.34 (13)
H1A—C1—H1B	108.3	C4 ⁱ —C4—H4A	110.0
C1—C2—C3	111.67 (13)	S1—C4—H4A	110.0
C1—C2—H2A	109.3	C4 ⁱ —C4—H4B	110.0
C3—C2—H2A	109.3	S1—C4—H4B	110.0
C1—C2—H2B	109.3	H4A—C4—H4B	108.4
C3—C2—H2B	109.3		
O1—S1—C1—C2	-71.04 (11)	O1—S1—C4—C4 ⁱ	65.40 (14)
C4—S1—C1—C2	179.23 (10)	C1—S1—C4—C4 ⁱ	175.68 (13)
S1—C1—C2—C3	-179.93 (11)		

Symmetry code: (i) $-x+1, -y, -z+1$.*Hydrogen-bond geometry (Å, °)*

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
C1—H1b \cdots O1 ⁱⁱ	0.99	2.41	3.3035 (19)	150

C4—H4a···O1 ⁱⁱ	0.99	2.40	3.2751 (19)	147
C4—H4b···O1 ⁱⁱⁱ	0.99	2.57	3.5124 (18)	159

Symmetry codes: (ii) $x, y+1, z$; (iii) $-x+1, y+1/2, -z+1/2$.