

Bis(3-ammoniomethylpyridinium) cyclotetraphosphate

Hanène Hemissi,^{a*} Mohammed Rzaigui^a and Salem S. Al-Deyab^b

^aLaboratoire de Chimie des Matériaux, Faculté des Sciences de Bizerte, 7021 Zarzouna Bizerte, Tunisia, and ^bPetrochemical Research Chair, College of Science, King Saud University, Riyadh, Saudi Arabia

Correspondence e-mail: hanene.hemissi@fsb.rnu.tn


Received 18 February 2010; accepted 2 March 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.034; wR factor = 0.101; data-to-parameter ratio = 34.5.

In the title compound, $2\text{C}_6\text{H}_{10}\text{N}_2^{2+} \cdot \text{P}_4\text{O}_{12}^{4-}$, which involves a doubly protonated 3-ammoniomethylpyridinium cation and a cyclotetraphosphate anion, the cyclotetraphosphoric ring is arranged around an inversion center and the organic entity alternates with it, forming hybrid ribbons parallel to the b axis. The crystal structure is stabilized by a three-dimensional network of $\text{N}-\text{H} \cdots \text{O}$ and weaker $\text{C}-\text{H} \cdots \text{O}$ hydrogen bonds.

Related literature

For properties of hybrid materials, see: Aakeröy *et al.* (1989); Sankar *et al.* (1993); Teraski *et al.* (1987); Vaughan (1993); Centi (1993); Ozin (1992). For related structures containing phosphoric acid rings, see: Aloui *et al.* (2003); Hemissi *et al.* (2005); Averbuch-Pouchot & Durif (1991); Durif (1995). For bond lengths in pyridine, see: Bak *et al.* (1959). For hydrogen bonding, see: Blessing (1986); Brown (1976); Soumhi & Jouini (1996). Cyclotetraphosphoric acid was produced from $\text{Na}_4\text{P}_4\text{O}_{12} \cdot 4\text{H}_2\text{O}$, which was prepared according to the Ondik (1964) process.



Experimental

Crystal data

$2\text{C}_6\text{H}_{10}\text{N}_2^{2+} \cdot \text{P}_4\text{O}_{12}^{4-}$

$M_r = 536.20$

Triclinic, $\overline{P}1$

$a = 7.849$ (2) Å

$b = 8.384$ (2) Å

$c = 9.448$ (2) Å

$\alpha = 113.24$ (2)°

$\beta = 98.73$ (3)°

$\gamma = 108.76$ (3)°

$V = 512.4$ (2) Å³

$Z = 1$

Ag $K\alpha$ radiation

$\lambda = 0.56083$ Å

$\mu = 0.23$ mm⁻¹

$T = 293$ K

$0.35 \times 0.3 \times 0.15$ mm

Data collection

Enraf–Nonius CAD-4

diffractometer

7923 measured reflections

5005 independent reflections

3963 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.012$

2 standard reflections every 120 min

intensity decay: 1%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$

$wR(F^2) = 0.101$

$S = 1.08$

5005 reflections

145 parameters

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.49$ e Å⁻³

$\Delta\rho_{\text{min}} = -0.47$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
$\text{N1}-\text{H1} \cdots \text{O6}$	0.86	1.77	2.6294 (18)	175
$\text{N2}-\text{H2A} \cdots \text{O5}^{\text{i}}$	0.89	1.88	2.7079 (17)	154
$\text{N2}-\text{H2B} \cdots \text{O3}^{\text{ii}}$	0.89	2.02	2.7350 (17)	137
$\text{N2}-\text{H2C} \cdots \text{O1}^{\text{iii}}$	0.89	2.08	2.831 (2)	141
$\text{C1}-\text{H1A} \cdots \text{O6}^{\text{iv}}$	0.93	2.55	3.381 (2)	149
$\text{C4}-\text{H4} \cdots \text{O5}^{\text{v}}$	0.93	2.48	3.281 (2)	144
$\text{C5}-\text{H5} \cdots \text{O4}$	0.93	2.60	3.256 (2)	128
$\text{C6}-\text{H6B} \cdots \text{O1}^{\text{i}}$	0.97	2.44	3.117 (2)	127

Symmetry codes: (i) $x + 1, y, z + 1$; (ii) $x, y, z + 1$; (iii) $-x + 1, -y + 1, -z + 1$; (iv) $-x, -y, -z + 1$; (v) $x + 1, y, z$.

Data collection: *CAD-4 EXPRESS* (Enraf–Nonius, 1994); cell refinement: *CAD-4 EXPRESS*; data reduction: *XCAD4* (Harms & Wocadlo, 1996); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 1997); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2541).

References

- Aakeröy, C. B., Hitchcock, P. B., Moyle, B. D. & Seddon, K. R. (1989). *J. Chem. Soc. Chem. Commun.* pp. 1856–1859.
- Aloui, Z., Abid, S. & Rzaigui, M. (2003). *J. Phys. Chem. Solids*, **65**, 923–926.
- Averbuch-Pouchot, M. T. & Durif, A. (1991). *Eur. J. Solid State Inorg. Chem.* **28**, 9–22.
- Bak, B., Hansen-Nygaard, L. & Rastrup-Andersen, J. (1959). *J. Mol. Spectrosc.* **2**, 361–364.
- Blessing, R. H. (1986). *Acta Cryst.* **B42**, 613–621.
- Brown, I. D. (1976). *Acta Cryst.* **A32**, 24–31.
- Centi, G. (1993). *Catal. Lett.* **22**, 53–66.
- Durif, A. (1995). *Crystal Chemistry of Condensed Phosphates*. New York: Plenum Press.
- Enraf–Nonius (1994). *CAD-4 EXPRESS*. Enraf–Nonius, Delft, The Netherlands.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565–566.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Harms, K. & Wocadlo, S. (1996). *XCAD4*. University of Marburg, Germany.
- Hemissi, H., Sonia, A. & Rzaigui, M. (2005). *Z. Kristallogr.* **220**, 265–266.
- Ondik, H. M. (1964). *Acta Cryst.* **17**, 1139–1145.
- Ozin, G. A. (1992). *Adv. Mater.* **4**, 612–649.
- Sankar, G., Wright, P. A., Natarajan, S., Thomas, J. M., Greaves, G. N., Dent, A. J., Dobson, B. R., Ramsdale, C. A. & Jones, R. H. (1993). *J. Phys. Chem.* **97**, 9550–9554.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.

Soumhi, E. H. & Jouini, T. (1996). *Acta Cryst.* **C52**, 432–433.

Teraski, O., Barry, J. C. & Thomas, J. M. (1987). *Nature (London)*, **330**, 58–60

Vaughan, D. E. W. (1993). In *Multifunctional Mesoporous Inorganic Solids*, edited by A.C. Sequina & M.I. Hudson, pp. 137–156. New York: Nato ASI Series.

supporting information

Acta Cryst. (2010). E66, o779–o780 [doi:10.1107/S1600536810008056]

Bis(3-ammoniomethylpyridinium) cyclotetraphosphate

Hanène Hemissi, Mohammed Rzaigui and Salem S. Al-Deyab

S1. Comment

Hybrid materials with organic and inorganic components continue to be a focus area in solid state chemistry and material sciences due to their potential applications in various fields, such as nonlinear optics (Aakeröy *et al.*, 1989), heterogeneous catalysis (Centi, 1993), photochemical and photophysical process (Ozin, 1992), molecular sieves (Vaughan, 1993), ceramic precursors (Sankar *et al.*, 1993) and other areas that include electronic materials (Teraski *et al.*, 1987). In the present paper, the results of the x-ray structure analysis of a new organic cyclotetraphosphate, bis(3-ammoniomethylpyridinium) cyclotetraphosphate, are discussed with respect to the geometry and flexibility of the cyclotetraphosphate ring system and H-bonding interactions between the inorganic acceptor and the organic donor molecules.

The chemical composition of the title compound (I) includes two fundamental entities, the $P_4O_{12}^{4-}$ ring and the organic cations $(C_6H_{10}N_2)^{2+}$. The geometrical configuration of these entities is depicted in Figure 1, while the complete atomic arrangement is shown in Figure 2. This latter shows that the crystal structure of $(C_6H_{10}N_2)_2P_4O_{12}$ can be described by hybrid ribbons where the organic and inorganic species are alternated. These ribbons, extended in the *b*-direction, are also connected between them in the two other directions via H-bonds to develop a three-dimensional network. The P_4O_{12} rings are located around the inversion center (0, 0, 0) and are built up by only two independent PO_4 tetrahedra. The P—P—P angles are 84.43 (1) and 95.57 (1)° and show that the tetramembered phosphoric rings are distorted in comparison with the ideal value (90°). It should be noted that such deviations are commonly observed in cyclotetraphosphoric ring anions with low internal symmetry as (I) (Aloui *et al.*, 2003; Hemissi *et al.*, 2005). Nevertheless, this distortion is comparatively less important than that observed in the hexamembered P_6O_{18} rings (93.2 - 145.5°) (Averbuch-Pouchot & Durif, 1991). Consequently, P_4O_{12} is less flexible than the P_6O_{18} what could explain the pronounced distortion observed for the big rings compared with their smaller rings analogues. In spite of this distortion, the examination of the main geometrical feature of PO_4 tetrahedra (P—O distances and P—O—P angles) shows that they are in accordance with values generally observed in condensed phosphates (Durif, 1995).

One crystallographically independent organic group exists in the asymmetric unit. Inside this organic molecule, both nitrogen atoms are protonated and so it is formulated $(C_6H_{10}N_2)^{2+}$. The examination of pyridinium ring shows that this unit is essentially planar with mean deviation of ± 0.0036 Å from least-square plane defined by the six constituent atoms. The average C—N distances in pyridinium ring is 1.337 (2) Å and of the C—C bond lengths is 1.384 (2) Å. The latter value, being shorter than 1.39–1.41 Å, reported for non-substituent pyridine, may indicate some aromatic bond characters (Bak *et al.*, 1959). The pyridinium ring is non coplanar with its methylamine substituent ($-CH_2-NH_3$) which is evidenced by the torsion angle value of (C1—C2—C6—N2) equal to 96.29 (2)°. In addition to electrostatic and van der Waals interactions, the structure is further stabilized with a three-dimensional network of N—H \cdots O and the weaker C—H \cdots O hydrogen bonds (Table 1, Fig. 2)). In the hydrogen-bond scheme two main points should be noticed: (i) there is a bridging oxygen atom (O4) of the P_4O_{12} ring involved in hydrogen bond and so that is rarely observed in organic condensed phosphates. Indeed, it was only observed in $(C_6H_{10}N_2)_2P_4O_{12}\cdot 2H_2O$ (Soumhi *et al.*, 1996). (ii) Inside the structure, there

are two strong hydrogen bonds with N \cdots O distances equal to 2.629 (2) and 2.708 (2) Å. The others are weaker within N(C) \cdots O distances falling from 2.735 (2) to 3.381 (2) Å (Brown, 1976; Blessing, 1986).

S2. Experimental

Crystals of the title compound were prepared by adding ethanolic solution (5 ml) of 3-aminopicolamine (11.04 mmol) dropwise to an aqueous solution of cyclotetraphosphoric acid (5.52 mmol, 20 ml). Good quality of colourless prisms were obtained after a slow evaporation during few days at ambient temperature. The cyclotetraphosphoric acid H₄P₄O₁₂, was produced from Na₄P₄O₁₂·4H₂O, prepared according to the Ondik process (Ondik, 1964), through an ion-exchange resin in H-state (Amberlite IR 120).

S3. Refinement

All H atoms were positioned geometrically and treated as riding on their parent atoms, [N–H = 0.89, C–H = 0.96 Å (CH₃) with with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}$ and C–H = 0.96 Å (Ar–H), with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}$




Figure 1

An ORTEP view of (I) with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii. [Symmetry codes: (i) 1 - x, 1 - y, 1 - z].





Figure 2

Projection of (I) along a axis.

Bis(3-ammoniomethylpyridinium) cyclotetraphosphate

Crystal data



$M_r = 536.20$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 7.849\ (2)\ \text{\AA}$

$b = 8.384\ (2)\ \text{\AA}$

$c = 9.448\ (2)\ \text{\AA}$

$\alpha = 113.24\ (2)^\circ$

$\beta = 98.73\ (3)^\circ$

$\gamma = 108.76\ (3)^\circ$

$V = 512.4\ (2)\ \text{\AA}^3$

$Z = 1$

$F(000) = 276$

$D_x = 1.738\ \text{Mg m}^{-3}$

Ag $K\alpha$ radiation, $\lambda = 0.56083\ \text{\AA}$

Cell parameters from 25 reflections

$\theta = 9.0\text{--}10.7^\circ$

$\mu = 0.23\ \text{mm}^{-1}$

$T = 293\ \text{K}$

Prism, colourless

$0.35 \times 0.3 \times 0.15\ \text{mm}$

Data collection

Enraf–Nonius CAD-4

diffractometer

Radiation source: Enraf–Nonius FR590

Graphite monochromator

non-profiled ω scans

7923 measured reflections

5005 independent reflections

3963 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.012$

$\theta_{\text{max}} = 28.0^\circ$, $\theta_{\text{min}} = 2.2^\circ$

$h = -13 \rightarrow 13$

$k = -14 \rightarrow 14$

$l = -6 \rightarrow 15$

2 standard reflections every 120 min

intensity decay: 1%

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.034$

$wR(F^2) = 0.101$

$S = 1.08$

5005 reflections

145 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.049P)^2 + 0.1514P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$

$$\Delta\rho_{\max} = 0.49 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.47 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O4	0.14028 (12)	0.17020 (14)	0.16858 (12)	0.02591 (17)
O2	0.20484 (12)	0.04327 (13)	-0.08942 (12)	0.02816 (18)
O1	0.07961 (14)	0.30061 (15)	-0.02599 (13)	0.03096 (19)
O3	0.41615 (12)	0.37060 (14)	0.12629 (14)	0.0328 (2)
O5	-0.09804 (14)	0.30451 (14)	0.25390 (12)	0.02842 (18)
O6	0.01424 (14)	0.12073 (17)	0.37735 (13)	0.0330 (2)
P1	0.21242 (4)	0.23999 (4)	0.04410 (4)	0.02125 (7)
P2	-0.03708 (4)	0.15037 (4)	0.23497 (4)	0.02013 (7)
N2	0.71399 (14)	0.34670 (15)	1.01546 (13)	0.02438 (18)
H2A	0.7405	0.3187	1.0954	0.037*
H2B	0.6259	0.3920	1.0259	0.037*
H2C	0.8188	0.4345	1.0204	0.037*
N1	0.37134 (15)	0.21366 (16)	0.52122 (14)	0.0284 (2)
H1	0.2559	0.1902	0.4771	0.034*
C1	0.41069 (16)	0.17716 (18)	0.64460 (16)	0.0259 (2)
H1A	0.3140	0.1291	0.6824	0.031*
C4	0.6907 (2)	0.3225 (2)	0.53050 (17)	0.0320 (3)
H4	0.7847	0.3727	0.4913	0.038*
C6	0.64164 (17)	0.17070 (18)	0.85588 (16)	0.0264 (2)
H6A	0.5287	0.0744	0.8515	0.032*
H6B	0.7369	0.1203	0.8456	0.032*
C2	0.59489 (16)	0.21046 (16)	0.71693 (14)	0.02293 (19)
C5	0.5057 (2)	0.28559 (19)	0.46376 (16)	0.0309 (2)
H5	0.4735	0.3106	0.3784	0.037*
C3	0.73527 (17)	0.28345 (19)	0.65771 (16)	0.0288 (2)
H3	0.8598	0.3064	0.7034	0.035*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O4	0.0190 (3)	0.0374 (5)	0.0315 (4)	0.0132 (3)	0.0114 (3)	0.0233 (4)
O2	0.0190 (3)	0.0275 (4)	0.0306 (4)	0.0079 (3)	0.0068 (3)	0.0089 (3)

O1	0.0281 (4)	0.0376 (5)	0.0413 (5)	0.0170 (4)	0.0131 (4)	0.0284 (4)
O3	0.0174 (3)	0.0290 (4)	0.0409 (5)	0.0034 (3)	0.0105 (3)	0.0109 (4)
O5	0.0331 (4)	0.0325 (4)	0.0273 (4)	0.0199 (4)	0.0114 (4)	0.0157 (4)
O6	0.0267 (4)	0.0532 (6)	0.0328 (5)	0.0175 (4)	0.0104 (4)	0.0320 (5)
P1	0.01556 (11)	0.02324 (13)	0.02726 (14)	0.00693 (9)	0.00856 (10)	0.01444 (11)
P2	0.01707 (11)	0.02625 (13)	0.02125 (13)	0.00966 (10)	0.00632 (9)	0.01460 (11)
N2	0.0213 (4)	0.0309 (5)	0.0267 (4)	0.0122 (3)	0.0078 (3)	0.0180 (4)
N1	0.0235 (4)	0.0314 (5)	0.0285 (5)	0.0109 (4)	0.0019 (4)	0.0152 (4)
C1	0.0198 (4)	0.0305 (5)	0.0293 (5)	0.0101 (4)	0.0065 (4)	0.0167 (5)
C4	0.0303 (6)	0.0337 (6)	0.0281 (6)	0.0080 (5)	0.0117 (5)	0.0146 (5)
C6	0.0250 (5)	0.0289 (5)	0.0287 (5)	0.0123 (4)	0.0054 (4)	0.0172 (4)
C2	0.0197 (4)	0.0245 (5)	0.0235 (5)	0.0091 (4)	0.0047 (4)	0.0114 (4)
C5	0.0360 (6)	0.0298 (6)	0.0248 (5)	0.0116 (5)	0.0055 (5)	0.0143 (5)
C3	0.0201 (4)	0.0345 (6)	0.0284 (6)	0.0092 (4)	0.0073 (4)	0.0137 (5)

Geometric parameters (Å, °)

O4—P2	1.5992 (10)	N1—C5	1.3393 (19)
O4—P1	1.6057 (10)	N1—H1	0.8600
O2—P2 ⁱ	1.6044 (14)	C1—C2	1.3875 (17)
O2—P1	1.6085 (11)	C1—H1A	0.9300
O1—P1	1.4766 (10)	C4—C5	1.372 (2)
O3—P1	1.4781 (12)	C4—C3	1.389 (2)
O5—P2	1.4739 (10)	C4—H4	0.9300
O6—P2	1.4824 (10)	C6—C2	1.4990 (17)
P2—O2 ⁱ	1.6044 (14)	C6—H6A	0.9700
N2—C6	1.4894 (18)	C6—H6B	0.9700
N2—H2A	0.8900	C2—C3	1.3875 (18)
N2—H2B	0.8900	C5—H5	0.9300
N2—H2C	0.8900	C3—H3	0.9300
N1—C1	1.3345 (17)		
P2—O4—P1	136.26 (6)	C5—N1—H1	118.9
P2 ⁱ —O2—P1	134.03 (6)	N1—C1—C2	120.46 (12)
O1—P1—O3	120.41 (7)	N1—C1—H1A	119.8
O1—P1—O4	110.99 (6)	C2—C1—H1A	119.8
O3—P1—O4	106.65 (6)	C5—C4—C3	118.82 (13)
O1—P1—O2	111.42 (7)	C5—C4—H4	120.6
O3—P1—O2	105.82 (7)	C3—C4—H4	120.6
O4—P1—O2	99.36 (6)	N2—C6—C2	111.53 (10)
O5—P2—O6	119.04 (7)	N2—C6—H6A	109.3
O5—P2—O4	112.23 (6)	C2—C6—H6A	109.3
O6—P2—O4	105.42 (6)	N2—C6—H6B	109.3
O5—P2—O2 ⁱ	106.33 (6)	C2—C6—H6B	109.3
O6—P2—O2 ⁱ	109.45 (7)	H6A—C6—H6B	108.0
O4—P2—O2 ⁱ	103.28 (6)	C3—C2—C1	117.98 (11)
C6—N2—H2A	109.5	C3—C2—C6	120.87 (11)
C6—N2—H2B	109.5	C1—C2—C6	121.15 (11)

H2A—N2—H2B	109.5	N1—C5—C4	120.20 (12)
C6—N2—H2C	109.5	N1—C5—H5	119.9
H2A—N2—H2C	109.5	C4—C5—H5	119.9
H2B—N2—H2C	109.5	C2—C3—C4	120.36 (12)
C1—N1—C5	122.18 (11)	C2—C3—H3	119.8
C1—N1—H1	118.9	C4—C3—H3	119.8

Symmetry code: (i) $-x, -y, -z$.

Hydrogen-bond geometry ($\text{\AA}, ^\circ$)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
N1—H1...O6	0.86	1.77	2.6294 (18)	175
N2—H2A...O5 ⁱⁱ	0.89	1.88	2.7079 (17)	154
N2—H2B...O3 ⁱⁱⁱ	0.89	2.02	2.7350 (17)	137
N2—H2C...O1 ^{iv}	0.89	2.08	2.831 (2)	141
C1—H1A...O6 ^v	0.93	2.55	3.381 (2)	149
C4—H4...O5 ^{vi}	0.93	2.48	3.281 (2)	144
C5—H5...O4	0.93	2.60	3.256 (2)	128
C6—H6B...O1 ⁱⁱ	0.97	2.44	3.117 (2)	127

Symmetry codes: (ii) $x+1, y, z+1$; (iii) $x, y, z+1$; (iv) $-x+1, -y+1, -z+1$; (v) $-x, -y, -z+1$; (vi) $x+1, y, z$.