

Potassium (1-methoxycarbonyl-2-methyl-prop-2-en-2-ylidene)azinate

Cédric Reuter, Jörg M. Neudörfl and Hans-Günther Schmalz*

Department für Chemie der Universität zu Köln, Greinstraße 4, 50939 Köln,
 Germany

Correspondence e-mail: schmalz@uni-koeln.de


Received 18 February 2010; accepted 18 March 2010

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$;
 R factor = 0.028; wR factor = 0.061; data-to-parameter ratio = 16.3.

In the title compound, $\text{K}^+\cdot\text{C}_6\text{H}_8\text{NO}_4^-$, the K^+ cations have a coordination number of seven and are surrounded by four bidentate azinate anions. The methylene groups of the anions are always directed towards the coordinated potassium cations. The $\text{N}-\text{C}-\text{C}-\text{C}$ torsion angle is $101.2(2)^\circ$. The orthogonal non-conjugated nature of the salt confirms the supposed geometry and reactivity of this compound.

Related literature

For a short overview of peptidomimetics, see: Grauer *et al.* (2009); Vagner *et al.* (2008); Wu *et al.* (2008). For the synthesis of peptidomimetics, amino-acid-based building blocks play a key role in the assembly of these structures, see: Kemp, Boyd & Muendel (1991); Kemp, Curran *et al.* (1991); Beal *et al.* (2000); Kühne *et al.* (2008). A known deprotonation/protonation sequence (Bouveault & Wahl, 1901) was used in the synthesis of the title compound. The protonation of the title compound occurs exclusively at the α -position and no protonation of the methylene group was observed (Baldwin *et al.*, 1977).



Experimental

Crystal data

$\text{K}^+\cdot\text{C}_6\text{H}_8\text{NO}_4^-$
 $M_r = 197.23$
 Monoclinic, $Cm2/c$
 $a = 23.9269(13)\text{ \AA}$

$b = 5.2909(2)\text{ \AA}$
 $c = 14.2510(7)\text{ \AA}$
 $\beta = 113.361(2)^\circ$
 $V = 1656.21(14)\text{ \AA}^3$

$Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.62\text{ mm}^{-1}$

$T = 100\text{ K}$
 $0.20 \times 0.15 \times 0.03\text{ mm}$

Data collection

Nonius KappaCCD diffractometer
 6264 measured reflections
 1810 independent reflections

1416 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.061$
 $S = 1.01$
 1810 reflections

111 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.31\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.27\text{ e \AA}^{-3}$

Table 1
 Selected bond lengths (\AA).

K1—O1 ⁱ	2.7036 (10)	K1—O2	2.8896 (10)
K1—O2 ⁱⁱ	2.7539 (11)	K1—O3 ⁱⁱⁱ	2.8970 (12)
K1—O3 ⁱ	2.7988 (11)	K1—O1 ⁱⁱⁱ	2.9080 (11)
K1—O1	2.7994 (11)		

Symmetry codes: (i) $-x + \frac{1}{2}, -y + \frac{1}{2}, -z$; (ii) $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$; (iii) $x, y - 1, z$.

Data collection: *COLLECT* (Hooft, 1998); cell refinement: *DENZO* (Otwinowski & Minor, 1997); data reduction: *DENZO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SCHAKAL99* (Keller, 1999); software used to prepare material for publication: *PLATON* (Spek, 2009), *publCIF* (Westrip, 2010) and *ORTEP* (Davenport *et al.*, 1999).

This work was supported by the Deutsche Forschungsgemeinschaft (FG 806).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JJ2024).

References

- Baldwin, J. E., Haber, S. B., Hoskins, C. & Kruse, L. I. (1977). *J. Org. Chem.* **42**, 1239–1241.
- Beal, L. M., Liu, B., Chu, W. & Moeller, K. D. (2000). *Tetrahedron*, **56**, 10113–10125.
- Bouveault, L. & Wahl, A. (1901). *Bull. Soc. Chim. Fr.* **25**, 800–817.
- Davenport, G., Hall, S. & Dreissig, W. (1999). *ORTEP in Xtal3.7 System*. Edited by S. R. Hall, D. J. du Boulay & R. Olthof-Hazekamp. University of Western Australia, Australia.
- Grauer, A. & König, B. (2009). *Eur. J. Org. Chem.* **30**, 5099–50111.
- Hooft, R. W. W. (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Keller, E. (1999). *SCHAKAL99*. University of Freiburg, Germany.
- Kemp, D. S., Boyd, J. G. & Muendel, C. C. (1991). *Nature (London)*, **352**, 451–454.
- Kemp, D. S., Curran, T. P., Boyd, J. G. & Allen, T. J. (1991). *J. Org. Chem.* **56**, 6683–6697.
- Kühne, R., Oschkinat, H., Brockmann, C. & Schmalz, H.-G. (2008). Ger. Patent WO 2008/040332 A1.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Vagner, J., Qu, H. & Hruby, V. J. (2008). *Curr. Opin. Chem. Biol.* **12**, 292–296.
- Westrip, S. P. (2010). *publCIF*. In preparation.
- Wu, Y.-D. & Gellman, S. (2008). *Acc. Chem. Res.* **41**, 1231–1232.

supporting information

Acta Cryst. (2010). E66, m461 [doi:10.1107/S1600536810010159]

Potassium (1-methoxycarbonyl-2-methylprop-2-en-2-ylidene)azinate

Cédric Reuter, Jörg M. Neudörfl and Hans-Günther Schmalz

S1. Comment

In the last decade, interest in peptidomimetics has lead to a fast growing research field within organic chemistry (Grauer *et al.*, 2009). Artificial peptide-like compounds are used to explore the principles of protein-protein interactions and their modulation (Vagner *et al.*, 2008; Wu *et al.*, 2008). In the synthesis of different peptidomimetics, amino acid based building blocks play a key role in the assembly of these structures (Kemp, Curran *et al.*, 1991; Kemp, Boyd & Muendel, 1991; Beal *et al.*, 2000; Kühne *et al.* 2008). In the context of our work we used an already known deprotonation/protonation sequence (Bouveault *et al.*, 1901) to synthesize our compound. The protonation of the title compound occurs exclusively at the α -position whereas no protonation of the methylene group was observed (Baldwin *et al.*, 1977).


In the title compound, $C_6H_8KNO_4$, (I), (Fig. 1), the deconjugation within the molecule combined with the high basicity of the nitro enolate provides a convincing explanation for the high selectivity of this reaction (Fig. 2). The crystal structure supports the assumption made by Baldwin *et al.* about the geometry of this potassium salt. The potassium cations in the crystal structure have a coordination number of seven and are surrounded by four azinate anions with K—O distances from 2.704 (1) to 2.908 (1) Å (Fig. 3). These can either bind via the carbonyl group or the nitrogen-bonded oxygen atoms whereas both motifs can be found as bridging units. The resulting polar layers of potassium cations surrounded by oxygen atoms are perfectly shielded by the methyl and methylene residues (Fig. 4). This results in loose interactions between the different layers and explains the facile mechanical fissility of the crystals.

S2. Experimental


The title compound, $C_6H_8KNO_4$, was prepared in good yield from methyl 3-methyl-2-nitrobut-2-enoate by deprotonation with potassium hydride. In a dry, argon-flushed 50 ml flask, 30.1 mmol of potassium hydride were suspended in 15 ml of dry THF. The suspension was cooled to 0 °C and a solution of 30.1 mmol methyl-3-methyl-2-nitrobut-2-enoate in 5 ml of THF was added via syringe over 30 min. After stirring for 5 h at room temperature a small amount of *n*-octanol was added at 0 °C to destroy the excess of potassium hydride. The paste was filtrated, washed three times with THF and dried in vacuo to give 4.950 g (25.1 mmol, 83%) of an ochre powder. A portion of the salt was recrystallized from MeOH/THF to give colourless prisms.

S3. Refinement

Hydrogen atoms were located in difference Fourier maps and refined at idealized positions (C—H = 0.98 Å for methyl H atoms and 0.95 Å for all other H Atoms) using a riding model. The U values of the hydrogens are constrained relative to U_{eq} of the parent carbon atom (1.2 x $U_{eq}(C)$ for C—H₂ and 1.5 x $U_{eq}(C)$ for methyl H).

**Figure 1**

A view of (I). Displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

Protonation of the title compound (I).





Figure 3

Coordination sphere of an isolated potassium Cation.

**Figure 4**

View of the unit cell along the *b*-axis.

Potassium (1-methoxycarbonyl-2-methylprop-2-en-2-ylidene)azinate

Crystal data

$K^+ \cdot C_6H_8NO_4^-$
 $M_r = 197.23$
Monoclinic, $Cm2/c$
Hall symbol: -C 2yc
 $a = 23.9269 (13)$ Å
 $b = 5.2909 (2)$ Å
 $c = 14.2510 (7)$ Å
 $\beta = 113.361 (2)^\circ$
 $V = 1656.21 (14)$ Å³
 $Z = 8$

$F(000) = 816$
 $D_x = 1.582$ Mg m⁻³
Melting point: 180.7(10) K
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 6264 reflections
 $\theta = 1.9-27.0^\circ$
 $\mu = 0.62$ mm⁻¹
 $T = 100$ K
Platlet, colourless
0.20 × 0.15 × 0.03 mm

Data collection

Nonius KappaCCD
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
Phi/ω-Scans scans
6264 measured reflections
1810 independent reflections

1416 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.041$
 $\theta_{\text{max}} = 27.0^\circ, \theta_{\text{min}} = 1.9^\circ$
 $h = -30 \rightarrow 30$
 $k = -6 \rightarrow 6$
 $l = -18 \rightarrow 18$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.061$
 $S = 1.01$
1810 reflections
111 parameters
0 restraints

Primary atom site location: structure-invariant
direct methods
Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0266P)^2]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$

$$\Delta\rho_{\max} = 0.31 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.27 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
K1	0.269013 (15)	-0.06555 (6)	0.11206 (2)	0.01586 (11)
O1	0.22159 (5)	0.42337 (19)	0.06333 (8)	0.0183 (3)
O2	0.20272 (5)	0.26997 (19)	0.19103 (8)	0.0172 (2)
O3	0.15057 (5)	0.81224 (18)	-0.04385 (8)	0.0194 (3)
O4	0.07200 (5)	0.87761 (19)	0.00265 (8)	0.0191 (3)
N1	0.19003 (6)	0.4306 (2)	0.11777 (9)	0.0144 (3)
C1	0.14436 (7)	0.5975 (3)	0.10141 (11)	0.0142 (3)
C2	0.12553 (7)	0.7664 (3)	0.01413 (12)	0.0155 (3)
C3	0.04879 (8)	1.0606 (3)	-0.07901 (12)	0.0213 (4)
H3A	0.0089	1.1211	-0.0844	0.032*
H3B	0.0446	0.9813	-0.1436	0.032*
H3C	0.0771	1.2034	-0.0644	0.032*
C4	0.11349 (7)	0.5958 (3)	0.17409 (12)	0.0164 (3)
C5	0.12805 (8)	0.7705 (3)	0.24651 (12)	0.0221 (4)
H5A	0.1578	0.8947	0.2516	0.027*
H5B	0.1087	0.7716	0.2933	0.027*
C6	0.06556 (8)	0.3977 (3)	0.15837 (14)	0.0248 (4)
H6A	0.0312	0.4272	0.0930	0.037*
H6B	0.0514	0.4067	0.2141	0.037*
H6C	0.0828	0.2300	0.1577	0.037*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
K1	0.01789 (19)	0.0171 (2)	0.01396 (19)	0.00072 (15)	0.00778 (15)	-0.00055 (14)
O1	0.0217 (6)	0.0208 (6)	0.0198 (6)	0.0039 (5)	0.0160 (5)	0.0020 (5)
O2	0.0228 (6)	0.0153 (6)	0.0153 (6)	0.0030 (5)	0.0095 (5)	0.0046 (5)
O3	0.0205 (6)	0.0232 (6)	0.0190 (6)	0.0044 (5)	0.0125 (5)	0.0050 (5)
O4	0.0167 (6)	0.0236 (6)	0.0204 (6)	0.0070 (5)	0.0108 (5)	0.0086 (5)
N1	0.0175 (7)	0.0143 (7)	0.0130 (7)	-0.0020 (6)	0.0076 (6)	-0.0014 (6)
C1	0.0142 (8)	0.0147 (8)	0.0156 (8)	0.0008 (6)	0.0080 (7)	0.0004 (6)
C2	0.0160 (9)	0.0148 (8)	0.0159 (8)	-0.0013 (7)	0.0065 (7)	-0.0038 (7)

C3	0.0198 (9)	0.0241 (9)	0.0201 (9)	0.0057 (8)	0.0082 (8)	0.0083 (7)
C4	0.0159 (8)	0.0185 (9)	0.0169 (9)	0.0066 (7)	0.0086 (7)	0.0053 (7)
C5	0.0261 (10)	0.0238 (9)	0.0207 (9)	0.0074 (7)	0.0138 (8)	0.0046 (7)
C6	0.0237 (10)	0.0247 (9)	0.0320 (10)	0.0009 (7)	0.0175 (9)	0.0046 (8)

Geometric parameters (\AA , $^{\circ}$)

K1—O1 ⁱ	2.7036 (10)	O4—C2	1.3591 (18)
K1—O2 ⁱⁱ	2.7539 (11)	O4—C3	1.4447 (18)
K1—O3 ⁱ	2.7988 (11)	N1—C1	1.3516 (19)
K1—O1	2.7994 (11)	N1—K1 ^{iv}	3.2874 (12)
K1—O2	2.8896 (10)	C1—C2	1.451 (2)
K1—O3 ⁱⁱⁱ	2.8970 (12)	C1—C4	1.4917 (19)
K1—O1 ⁱⁱⁱ	2.9080 (11)	C1—K1 ^{iv}	3.4260 (15)
K1—C5 ⁱⁱ	3.0584 (16)	C2—K1 ^{iv}	3.2747 (16)
K1—N1	3.2542 (13)	C3—H3A	0.9800
K1—C2 ⁱⁱⁱ	3.2747 (16)	C3—H3B	0.9800
K1—N1 ⁱⁱⁱ	3.2874 (12)	C3—H3C	0.9800
K1—C4 ⁱⁱ	3.3355 (16)	C4—C5	1.325 (2)
O1—N1	1.2799 (14)	C4—C6	1.503 (2)
O1—K1 ⁱ	2.7036 (10)	C4—K1 ^v	3.3355 (16)
O1—K1 ^{iv}	2.9081 (11)	C5—K1 ^v	3.0583 (16)
O2—N1	1.2856 (15)	C5—H5A	0.9500
O2—K1 ^v	2.7539 (11)	C5—H5B	0.9500
O3—C2	1.2219 (16)	C6—H6A	0.9800
O3—K1 ⁱ	2.7989 (11)	C6—H6B	0.9800
O3—K1 ^{iv}	2.8970 (12)	C6—H6C	0.9800
O1 ⁱ —K1—O2 ⁱⁱ	162.38 (3)	N1—K1—C4 ⁱⁱ	93.33 (4)
O1 ⁱ —K1—O3 ⁱ	59.21 (3)	C2 ⁱⁱⁱ —K1—C4 ⁱⁱ	145.11 (4)
O2 ⁱⁱ —K1—O3 ⁱ	106.36 (3)	N1 ⁱⁱⁱ —K1—C4 ⁱⁱ	118.00 (3)
O1 ⁱ —K1—O1	71.80 (3)	N1—O1—K1 ⁱ	146.87 (9)
O2 ⁱⁱ —K1—O1	117.16 (3)	N1—O1—K1	98.95 (7)
O3 ⁱ —K1—O1	127.70 (3)	K1 ⁱ —O1—K1	108.20 (3)
O1 ⁱ —K1—O2	116.67 (3)	N1—O1—K1 ^{iv}	95.49 (7)
O2 ⁱⁱ —K1—O2	75.42 (2)	K1 ⁱ —O1—K1 ^{iv}	78.14 (3)
O3 ⁱ —K1—O2	168.78 (3)	K1—O1—K1 ^{iv}	135.94 (4)
O1—K1—O2	45.40 (3)	N1—O2—K1 ^v	120.14 (8)
O1 ⁱ —K1—O3 ⁱⁱⁱ	76.57 (3)	N1—O2—K1	94.53 (7)
O2 ⁱⁱ —K1—O3 ⁱⁱⁱ	118.77 (3)	K1 ^v —O2—K1	129.79 (4)
O3 ⁱ —K1—O3 ⁱⁱⁱ	103.16 (3)	C2—O3—K1 ⁱ	136.99 (9)
O1—K1—O3 ⁱⁱⁱ	80.73 (3)	C2—O3—K1 ^{iv}	96.80 (9)
O2—K1—O3 ⁱⁱⁱ	85.11 (3)	K1 ⁱ —O3—K1 ^{iv}	76.84 (3)
O1 ⁱ —K1—O1 ⁱⁱⁱ	101.86 (3)	C2—O4—C3	115.46 (12)
O2 ⁱⁱ —K1—O1 ⁱⁱⁱ	82.21 (3)	O1—N1—O2	117.81 (11)
O3 ⁱ —K1—O1 ⁱⁱⁱ	74.94 (3)	O1—N1—C1	123.09 (12)
O1—K1—O1 ⁱⁱⁱ	135.94 (4)	O2—N1—C1	119.10 (12)
O2—K1—O1 ⁱⁱⁱ	116.22 (3)	O1—N1—K1	58.18 (6)

O3 ⁱⁱⁱ —K1—O1 ⁱⁱⁱ	55.87 (3)	O2—N1—K1	62.27 (7)
O1 ⁱ —K1—C5 ⁱⁱ	96.14 (4)	C1—N1—K1	163.88 (10)
O2 ⁱⁱ —K1—C5 ⁱⁱ	72.78 (4)	O1—N1—K1 ^{iv}	61.71 (6)
O3 ⁱ —K1—C5 ⁱⁱ	90.87 (4)	O2—N1—K1 ^{iv}	127.10 (9)
O1—K1—C5 ⁱⁱ	76.55 (4)	C1—N1—K1 ^{iv}	84.21 (8)
O2—K1—C5 ⁱⁱ	79.00 (4)	K1—N1—K1 ^{iv}	107.96 (4)
O3 ⁱⁱⁱ —K1—C5 ⁱⁱ	157.29 (4)	N1—C1—C2	120.36 (13)
O1 ⁱⁱⁱ —K1—C5 ⁱⁱ	146.51 (4)	N1—C1—C4	117.74 (13)
O1 ⁱ —K1—N1	93.48 (3)	C2—C1—C4	121.88 (13)
O2 ⁱⁱ —K1—N1	98.04 (3)	N1—C1—K1 ^{iv}	72.68 (8)
O3 ⁱ —K1—N1	150.55 (3)	C2—C1—K1 ^{iv}	71.72 (8)
O1—K1—N1	22.86 (3)	C4—C1—K1 ^{iv}	129.38 (10)
O2—K1—N1	23.19 (3)	O3—C2—O4	121.61 (14)
O3 ⁱⁱⁱ —K1—N1	78.35 (3)	O3—C2—C1	129.29 (14)
O1 ⁱⁱⁱ —K1—N1	125.49 (3)	O4—C2—C1	109.10 (12)
C5 ⁱⁱ —K1—N1	80.70 (4)	O3—C2—K1 ^{iv}	61.45 (8)
O1 ⁱ —K1—C2 ⁱⁱⁱ	97.99 (4)	O4—C2—K1 ^{iv}	135.44 (9)
O2 ⁱⁱ —K1—C2 ⁱⁱⁱ	98.10 (4)	C1—C2—K1 ^{iv}	83.41 (9)
O3 ⁱ —K1—C2 ⁱⁱⁱ	118.31 (4)	O4—C3—H3A	109.5
O1—K1—C2 ⁱⁱⁱ	83.79 (3)	O4—C3—H3B	109.5
O2—K1—C2 ⁱⁱⁱ	71.83 (3)	H3A—C3—H3B	109.5
O3 ⁱⁱⁱ —K1—C2 ⁱⁱⁱ	21.75 (3)	O4—C3—H3C	109.5
O1 ⁱⁱⁱ —K1—C2 ⁱⁱⁱ	53.27 (3)	H3A—C3—H3C	109.5
C5 ⁱⁱ —K1—C2 ⁱⁱⁱ	150.79 (4)	H3B—C3—H3C	109.5
N1—K1—C2 ⁱⁱⁱ	73.06 (3)	C5—C4—C1	119.13 (14)
O1 ⁱ —K1—N1 ⁱⁱⁱ	120.52 (3)	C5—C4—C6	123.51 (14)
O2 ⁱⁱ —K1—N1 ⁱⁱⁱ	68.28 (3)	C1—C4—C6	117.33 (13)
O3 ⁱ —K1—N1 ⁱⁱⁱ	96.40 (3)	C5—C4—K1 ^v	66.48 (9)
O1—K1—N1 ⁱⁱⁱ	125.11 (3)	C1—C4—K1 ^v	99.55 (9)
O2—K1—N1 ⁱⁱⁱ	94.52 (3)	C6—C4—K1 ^v	105.93 (10)
O3 ⁱⁱⁱ —K1—N1 ⁱⁱⁱ	56.00 (3)	C4—C5—K1 ^v	90.11 (10)
O1 ⁱⁱⁱ —K1—N1 ⁱⁱⁱ	22.80 (3)	C4—C5—H5A	120.0
C5 ⁱⁱ —K1—N1 ⁱⁱⁱ	140.88 (4)	K1 ^v —C5—H5A	88.2
N1—K1—N1 ⁱⁱⁱ	107.96 (4)	C4—C5—H5B	120.0
C2 ⁱⁱⁱ —K1—N1 ⁱⁱⁱ	43.50 (4)	K1 ^v —C5—H5B	91.7
O1 ⁱ —K1—C4 ⁱⁱ	115.12 (4)	H5A—C5—H5B	120.0
O2 ⁱⁱ —K1—C4 ⁱⁱ	51.13 (3)	C4—C6—H6A	109.5
O3 ⁱ —K1—C4 ⁱⁱ	89.44 (4)	C4—C6—H6B	109.5
O1—K1—C4 ⁱⁱ	95.85 (4)	H6A—C6—H6B	109.5
O2—K1—C4 ⁱⁱ	83.16 (3)	C4—C6—H6C	109.5
O3 ⁱⁱⁱ —K1—C4 ⁱⁱ	166.33 (3)	H6A—C6—H6C	109.5
O1 ⁱⁱⁱ —K1—C4 ⁱⁱ	124.33 (4)	H6B—C6—H6C	109.5
C5 ⁱⁱ —K1—C4 ⁱⁱ	23.41 (4)		
O1 ⁱ —K1—O1—N1	160.77 (10)	O1 ⁱⁱⁱ —K1—N1—O1	-125.49 (9)
O2 ⁱⁱ —K1—O1—N1	-35.71 (9)	C5 ⁱⁱ —K1—N1—O1	77.40 (8)
O3 ⁱ —K1—O1—N1	-178.51 (7)	C2 ⁱⁱⁱ —K1—N1—O1	-115.56 (8)
O2—K1—O1—N1	-10.30 (7)	N1 ⁱⁱⁱ —K1—N1—O1	-141.94 (7)

O3 ⁱⁱⁱ —K1—O1—N1	81.97 (8)	C4 ⁱⁱ —K1—N1—O1	97.17 (8)
O1 ⁱⁱⁱ —K1—O1—N1	72.42 (10)	O1 ⁱ —K1—N1—O2	-179.41 (8)
C5 ⁱⁱ —K1—O1—N1	-98.01 (8)	O2 ⁱⁱ —K1—N1—O2	-12.78 (9)
C2 ⁱⁱⁱ —K1—O1—N1	60.23 (8)	O3 ⁱ —K1—N1—O2	-158.75 (8)
N1 ⁱⁱⁱ —K1—O1—N1	45.79 (8)	O1—K1—N1—O2	-161.15 (13)
C4 ⁱⁱ —K1—O1—N1	-84.68 (8)	O3 ⁱⁱⁱ —K1—N1—O2	105.08 (8)
O1 ⁱ —K1—O1—K1 ⁱ	0.0	O1 ⁱⁱⁱ —K1—N1—O2	73.36 (8)
O2 ⁱⁱ —K1—O1—K1 ⁱ	163.52 (4)	C5 ⁱⁱ —K1—N1—O2	-83.74 (8)
O3 ⁱ —K1—O1—K1 ⁱ	20.72 (6)	C2 ⁱⁱⁱ —K1—N1—O2	83.29 (8)
O2—K1—O1—K1 ⁱ	-171.07 (6)	N1 ⁱⁱⁱ —K1—N1—O2	56.91 (9)
O3 ⁱⁱⁱ —K1—O1—K1 ⁱ	-78.80 (4)	C4 ⁱⁱ —K1—N1—O2	-63.97 (8)
O1 ⁱⁱⁱ —K1—O1—K1 ⁱ	-88.35 (6)	O1 ⁱ —K1—N1—C1	81.4 (3)
C5 ⁱⁱ —K1—O1—K1 ⁱ	101.21 (5)	O2 ⁱⁱ —K1—N1—C1	-112.0 (3)
N1—K1—O1—K1 ⁱ	-160.77 (10)	O3 ⁱ —K1—N1—C1	102.0 (3)
C2 ⁱⁱⁱ —K1—O1—K1 ⁱ	-100.54 (4)	O1—K1—N1—C1	99.6 (4)
N1 ⁱⁱⁱ —K1—O1—K1 ⁱ	-114.98 (4)	O2—K1—N1—C1	-99.2 (3)
C4 ⁱⁱ —K1—O1—K1 ⁱ	114.55 (4)	O3 ⁱⁱⁱ —K1—N1—C1	5.9 (3)
O1 ⁱ —K1—O1—K1 ^{iv}	-91.64 (6)	O1 ⁱⁱⁱ —K1—N1—C1	-25.9 (3)
O2 ⁱⁱ —K1—O1—K1 ^{iv}	71.88 (6)	C5 ⁱⁱ —K1—N1—C1	177.0 (3)
O3 ⁱ —K1—O1—K1 ^{iv}	-70.93 (7)	C2 ⁱⁱⁱ —K1—N1—C1	-15.9 (3)
O2—K1—O1—K1 ^{iv}	97.29 (7)	N1 ⁱⁱⁱ —K1—N1—C1	-42.3 (4)
O3 ⁱⁱⁱ —K1—O1—K1 ^{iv}	-170.45 (6)	C4 ⁱⁱ —K1—N1—C1	-163.2 (3)
O1 ⁱⁱⁱ —K1—O1—K1 ^{iv}	180.0	O1 ⁱ —K1—N1—K1 ^{iv}	-56.32 (4)
C5 ⁱⁱ —K1—O1—K1 ^{iv}	9.57 (6)	O2 ⁱⁱ —K1—N1—K1 ^{iv}	110.31 (4)
N1—K1—O1—K1 ^{iv}	107.58 (10)	O3 ⁱ —K1—N1—K1 ^{iv}	-35.66 (9)
C2 ⁱⁱⁱ —K1—O1—K1 ^{iv}	167.82 (6)	O1—K1—N1—K1 ^{iv}	-38.06 (7)
N1 ⁱⁱⁱ —K1—O1—K1 ^{iv}	153.38 (4)	O2—K1—N1—K1 ^{iv}	123.09 (9)
C4 ⁱⁱ —K1—O1—K1 ^{iv}	22.91 (6)	O3 ⁱⁱⁱ —K1—N1—K1 ^{iv}	-131.83 (4)
O1 ⁱ —K1—O2—N1	0.66 (9)	O1 ⁱⁱⁱ —K1—N1—K1 ^{iv}	-163.55 (3)
O2 ⁱⁱ —K1—O2—N1	166.92 (9)	C5 ⁱⁱ —K1—N1—K1 ^{iv}	39.34 (4)
O3 ⁱ —K1—O2—N1	66.31 (19)	C2 ⁱⁱⁱ —K1—N1—K1 ^{iv}	-153.62 (4)
O1—K1—O2—N1	10.16 (7)	N1 ⁱⁱⁱ —K1—N1—K1 ^{iv}	180.0
O3 ⁱⁱⁱ —K1—O2—N1	-71.65 (8)	C4 ⁱⁱ —K1—N1—K1 ^{iv}	59.12 (4)
O1 ⁱⁱⁱ —K1—O2—N1	-119.59 (8)	O1—N1—C1—C2	5.1 (2)
C5 ⁱⁱ —K1—O2—N1	92.06 (8)	O2—N1—C1—C2	-174.70 (13)
C2 ⁱⁱⁱ —K1—O2—N1	-89.27 (8)	K1—N1—C1—C2	-84.2 (4)
N1 ⁱⁱⁱ —K1—O2—N1	-126.92 (9)	K1 ^{iv} —N1—C1—C2	55.71 (13)
C4 ⁱⁱ —K1—O2—N1	115.38 (8)	O1—N1—C1—C4	-176.54 (12)
O1 ⁱ —K1—O2—K1 ^v	-135.53 (5)	O2—N1—C1—C4	3.7 (2)
O2 ⁱⁱ —K1—O2—K1 ^v	30.73 (5)	K1—N1—C1—C4	94.1 (3)
O3 ⁱ —K1—O2—K1 ^v	-69.88 (17)	K1 ^{iv} —N1—C1—C4	-125.94 (12)
O1—K1—O2—K1 ^v	-126.03 (7)	O1—N1—C1—K1 ^{iv}	-50.60 (12)
O3 ⁱⁱⁱ —K1—O2—K1 ^v	152.16 (5)	O2—N1—C1—K1 ^{iv}	129.60 (12)
O1 ⁱⁱⁱ —K1—O2—K1 ^v	104.22 (5)	K1—N1—C1—K1 ^{iv}	-139.9 (3)
C5 ⁱⁱ —K1—O2—K1 ^v	-44.14 (6)	K1 ⁱ —O3—C2—O4	-153.88 (10)
N1—K1—O2—K1 ^v	-136.19 (11)	K1 ^{iv} —O3—C2—O4	128.14 (13)
C2 ⁱⁱⁱ —K1—O2—K1 ^v	134.54 (6)	K1 ⁱ —O3—C2—C1	25.9 (3)
N1 ⁱⁱⁱ —K1—O2—K1 ^v	96.89 (5)	K1 ^{iv} —O3—C2—C1	-52.12 (17)

C4 ⁱⁱ —K1—O2—K1 ^v	−20.81 (5)	K1 ⁱ —O3—C2—K1 ^{iv}	77.98 (12)
K1 ⁱ —O1—N1—O2	163.96 (11)	C3—O4—C2—O3	−2.9 (2)
K1—O1—N1—O2	18.87 (12)	C3—O4—C2—C1	177.36 (12)
K1 ^{iv} —O1—N1—O2	−119.37 (11)	C3—O4—C2—K1 ^{iv}	77.08 (17)
K1 ⁱ —O1—N1—C1	−15.8 (2)	N1—C1—C2—O3	−11.9 (2)
K1—O1—N1—C1	−160.93 (12)	C4—C1—C2—O3	169.81 (15)
K1 ^{iv} —O1—N1—C1	60.82 (14)	K1 ^{iv} —C1—C2—O3	44.26 (15)
K1 ⁱ —O1—N1—K1	145.09 (16)	N1—C1—C2—O4	167.86 (13)
K1 ^{iv} —O1—N1—K1	−138.24 (6)	C4—C1—C2—O4	−10.4 (2)
K1 ⁱ —O1—N1—K1 ^{iv}	−76.67 (13)	K1 ^{iv} —C1—C2—O4	−135.97 (11)
K1—O1—N1—K1 ^{iv}	138.24 (6)	N1—C1—C2—K1 ^{iv}	−56.16 (13)
K1 ^v —O2—N1—O1	123.95 (10)	C4—C1—C2—K1 ^{iv}	125.55 (13)
K1—O2—N1—O1	−18.09 (12)	N1—C1—C4—C5	101.20 (18)
K1 ^v —O2—N1—C1	−56.24 (15)	C2—C1—C4—C5	−80.5 (2)
K1—O2—N1—C1	161.72 (11)	K1 ^{iv} —C1—C4—C5	11.4 (2)
K1 ^v —O2—N1—K1	142.04 (8)	N1—C1—C4—C6	−80.71 (18)
K1 ^v —O2—N1—K1 ^{iv}	49.79 (12)	C2—C1—C4—C6	97.62 (18)
K1—O2—N1—K1 ^{iv}	−92.25 (8)	K1 ^{iv} —C1—C4—C6	−170.51 (10)
O1 ⁱ —K1—N1—O1	−18.26 (10)	N1—C1—C4—K1 ^v	32.88 (14)
O2 ⁱⁱ —K1—N1—O1	148.37 (8)	C2—C1—C4—K1 ^v	−148.79 (12)
O3 ⁱ —K1—N1—O1	2.40 (12)	K1 ^{iv} —C1—C4—K1 ^v	−56.92 (11)
O2—K1—N1—O1	161.15 (13)	C1—C4—C5—K1 ^v	−87.97 (13)
O3 ⁱⁱⁱ —K1—N1—O1	−93.78 (8)	C6—C4—C5—K1 ^v	94.06 (15)

Symmetry codes: (i) $-x+1/2, -y+1/2, -z$; (ii) $-x+1/2, y-1/2, -z+1/2$; (iii) $x, y-1, z$; (iv) $x, y+1, z$; (v) $-x+1/2, y+1/2, -z+1/2$.