

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

## Dichlorido{4-cyclohexyl-1-[1-(2-pyridyl- $\kappa N$ )ethylidene]thiosemicarbazidato- $\kappa^2 N^1, S$ ]methyltin(IV). Corrigendum

Md. Abdus Salam,<sup>a</sup> Md. Abu Affan,<sup>a</sup> Mustaffa Shamsuddin<sup>b</sup> and Seik Weng Ng<sup>c\*</sup>

<sup>a</sup>Faculty of Resource Science and Technology, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia, <sup>b</sup>Department of Chemistry, Faculty of Science, Universiti Teknologi Malaysia, 81310 UTM Skudai, Johor, Malaysia, and <sup>c</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my

Received 6 September 2010; accepted 20 November 2010

The address of one of the authors in the paper by Salam *et al.* [*Acta Cryst.* (2010), E66, m570] is corrected.

---

In the paper by Salam *et al.* (2010), the address of Mustaffa Shamsuddin is given incorrectly. The correct address is 'Department of Chemistry, Faculty of Science, Universiti Teknologi Malaysia, 81310 UTM Skudai, Johor, Malaysia', as given above.

### References

Salam, Md. A., Affan, Md. A., Shamsuddin, M. & Ng, S. W. (2010). *Acta Cryst.* E66, m570.

## Dichlorido[4-cyclohexyl-1-[1-(2-pyridyl- $\kappa N$ )ethylidene]thiosemicarbazidato- $\kappa^2 N^1, S$ ]methyltin(IV)

Md. Abdus Salam,<sup>a</sup> Md. Abu Affan,<sup>a</sup> Mustaffa Shamsuddin<sup>a</sup> and Seik Weng Ng<sup>b\*</sup>

<sup>a</sup>Faculty of Resource Science and Technology, Universiti Malaysia Sarawak, 94300 Kota Samarahan, Sarawak, Malaysia, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 19 April 2010; accepted 20 April 2010

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(C-C) = 0.005$  Å;  $R$  factor = 0.021;  $wR$  factor = 0.054; data-to-parameter ratio = 21.0.

The monodeprotonated Schiff base ligand in the title compound,  $[Sn(CH_3)(C_{14}H_{19}N_4S)Cl_2]$ ,  $N, N', S$ -chelates to the Sn atom, which is six-coordinated in an octahedral environment. The three coordinating atoms along with the methyl C atom comprise a square plane, above and below which are positioned the Cl atoms. The amino group is a hydrogen-bond donor to a Cl atom of an adjacent molecule, the hydrogen bond giving rise to a helical chain extending parallel to  $[100]$ .

### Related literature

For the crystal structures of other metal derivatives of the Schiff base, see: Joseph *et al.* (2004); Kovala-Demertzi *et al.* (2007).



### Experimental

#### Crystal data

$[Sn(CH_3)(C_{14}H_{19}N_4S)Cl_2]$   
 $M_r = 480.02$   
 Orthorhombic,  $P2_12_12_1$   
 $a = 9.2016$  (5) Å  
 $b = 12.2434$  (7) Å  
 $c = 17.4544$  (10) Å

$V = 1966.39$  (19) Å<sup>3</sup>  
 $Z = 4$   
 Mo  $K\alpha$  radiation  
 $\mu = 1.68$  mm<sup>-1</sup>  
 $T = 293$  K  
 $0.30 \times 0.25 \times 0.20$  mm

#### Data collection

Bruker SMART APEX diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{min} = 0.633$ ,  $T_{max} = 0.730$

18838 measured reflections  
 4498 independent reflections  
 4223 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.025$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$   
 $wR(F^2) = 0.054$   
 $S = 1.02$   
 4498 reflections  
 214 parameters  
 1 restraint

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{max} = 0.29$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -0.41$  e Å<sup>-3</sup>  
 Absolute structure: Flack (1983), 1493 Friedel pairs  
 Flack parameter:  $-0.020$  (17)

**Table 1**

Selected bond lengths (Å).

Sn1—C1	2.136 (3)	Sn1—S1	2.4814 (7)
Sn1—N1	2.269 (2)	Sn1—Cl1	2.4960 (7)
Sn1—N2	2.224 (2)	Sn1—Cl2	2.4701 (8)

**Table 2**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N4-H4\cdots Cl1^i$	0.86 (1)	2.36 (1)	3.219 (3)	177 (3)

Symmetry code: (i)  $x + \frac{1}{2}, -y + \frac{3}{2}, -z$ .

Data collection: APEX2 (Bruker, 2009); cell refinement: SAINT (Bruker, 2009); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

We thank MOSTI (grant No. 06-01-09-SF0046), Universiti Malaysia Sarawak and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5250).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
 Bruker (2009). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.  
 Joseph, M., Suni, V., Kurup, M. R. P., Nethaji, M., Kishore, A. & Bhat, S. G. (2004). *Polyhedron*, **23**, 3069–3080.  
 Kovala-Demertzi, D., Galani, A., Kourkoumelis, N., Miller, J. R. & Demertzi, M. A. (2007). *Polyhedron*, **26**, 2871–2879.  
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.

## supporting information

*Acta Cryst.* (2010). E66, m570 [https://doi.org/10.1107/S1600536810014443]

## Dichlorido{4-cyclohexyl-1-[1-(2-pyridyl- $\kappa$ N)ethylidene]thiosemicarbazidato- $\kappa^2$ N<sup>1</sup>,S}methyltin(IV)

Md. Abdus Salam, Md. Abu Affan, Mustaffa Shamsuddin and Seik Weng Ng

### S1. Comment

The mono-deprotonated anion of 2-acetylpyridine 4-cyclohexyl thiosemicarbazone is a ligand that  $N,N',S$ -binds to metal atoms (Joseph *et al.*, 2004). Whereas similar ligands have been complexed with diorganotin and triorganotin systems, the monoorganotin analogues have not been so extensively studied. The mono-deprotonated Schiff-base ligand in  $\text{SnCl}_2(\text{CH}_3)$  ( $\text{C}_{14}\text{H}_{19}\text{N}_4\text{S}$ )  $N,N',S$ -chelates to the tin atom, which is six-coordinate in an octahedral environment (Scheme I, Fig. 1). The three coordinating atoms along with the methyl carbon comprise a square plane, above and below which are positioned the chlorine atoms.

### S2. Experimental

2-Acetylpyridine 4-cyclohexyl thiosemicarbazone was synthesized by using a literature method (Joseph *et al.*, 2004). The compound (0.28 g, 1 mmol) was dissolved in dry methanol (10 ml) in a Schlenk apparatus under a nitrogen atmosphere. Methyltin trichloride (0.24 g, 1 mmol) dissolved in methanol (10 ml) was added. The mixture was heated for an hour. The solvent was removed and the yellow compound recrystallized from chloroform/methanol (1:1) in 70% yield, m.p. 551-553 K.

### S3. Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.93 to 0.978 Å) and were included in the refinement in the riding model approximation, with  $U(\text{H})$  set to 1.2 to 1.5 $U(\text{C})$ .

The amino H-atom was located in a difference Fourier map, and was refined isotropically with a distance restraint of N—H 0.84±0.01 Å.




Figure 1

Anisotropic displacement ellipsoid plot (Barbour, 2001) of  $\text{SnCl}_2(\text{CH}_3)(\text{C}_{14}\text{H}_{19}\text{N}_4\text{S})$  at the 50% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

### Dichlorido{4-cyclohexyl-1-[1-(2-pyridyl)- $\kappa\text{N}$ ]ethylidene}thiosemicarbazidato- $\kappa^2\text{N}^1,\text{S}$ ]methyltin(IV)

#### Crystal data

$[\text{Sn}(\text{CH}_3)(\text{C}_{14}\text{H}_{19}\text{N}_4\text{S})\text{Cl}_2]$

$M_r = 480.02$

Orthorhombic,  $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 9.2016$  (5) Å

$b = 12.2434$  (7) Å

$c = 17.4544$  (10) Å

$V = 1966.39$  (19) Å<sup>3</sup>

$Z = 4$

$F(000) = 960$

$D_x = 1.621$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 8827 reflections

$\theta = 2.3\text{--}27.7^\circ$

$\mu = 1.68$  mm<sup>-1</sup>

$T = 293$  K

Prism, yellow

$0.30 \times 0.25 \times 0.20$  mm

#### Data collection

Bruker SMART APEX  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.633$ ,  $T_{\max} = 0.730$

18838 measured reflections

4498 independent reflections

4223 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.025$

$\theta_{\max} = 27.5^\circ$ ,  $\theta_{\min} = 2.0^\circ$

$h = -11 \rightarrow 11$

$k = -15 \rightarrow 15$

$l = -22 \rightarrow 22$

#### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.021$

$wR(F^2) = 0.054$

$S = 1.02$

4498 reflections

214 parameters

1 restraint

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H atoms treated by a mixture of independent  
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.032P)^2 + 0.1524P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.29$  e Å<sup>-3</sup>

$$\Delta\rho_{\min} = -0.41 \text{ e } \text{\AA}^{-3}$$

Absolute structure: Flack (1983), 1493 Friedel  
pairs  
Absolute structure parameter:  $-0.020$  (17)

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	1.061930 (18)	0.634199 (13)	0.160119 (10)	0.03748 (6)
Cl1	0.97585 (8)	0.54078 (6)	0.04139 (4)	0.04922 (17)
Cl2	1.10680 (12)	0.73379 (8)	0.28090 (5)	0.0724 (3)
S1	1.22890 (7)	0.75142 (6)	0.08461 (5)	0.04529 (16)
N1	0.8388 (3)	0.59846 (18)	0.20943 (13)	0.0411 (5)
N2	0.9157 (2)	0.77150 (16)	0.12825 (11)	0.0351 (5)
N3	0.9636 (2)	0.85860 (18)	0.08811 (12)	0.0398 (5)
N4	1.1533 (3)	0.9409 (2)	0.02881 (19)	0.0602 (7)
H4	1.2404 (17)	0.944 (3)	0.0111 (18)	0.063 (11)*
C1	1.1749 (4)	0.4916 (3)	0.1978 (2)	0.0687 (10)
H1A	1.1755	0.4893	0.2527	0.103*
H1B	1.2731	0.4936	0.1792	0.103*
H1C	1.1270	0.4277	0.1783	0.103*
C2	0.8073 (4)	0.5114 (2)	0.25205 (17)	0.0524 (7)
H2	0.8783	0.4586	0.2602	0.063*
C3	0.6713 (4)	0.4978 (3)	0.28448 (19)	0.0599 (9)
H3	0.6513	0.4370	0.3146	0.072*
C4	0.5676 (4)	0.5746 (3)	0.27171 (17)	0.0589 (8)
H4A	0.4761	0.5675	0.2938	0.071*
C5	0.5985 (3)	0.6628 (2)	0.22600 (16)	0.0490 (7)
H5	0.5271	0.7146	0.2157	0.059*
C6	0.7362 (3)	0.6743 (2)	0.19527 (14)	0.0372 (6)
C7	0.7802 (3)	0.7679 (2)	0.14765 (14)	0.0363 (5)
C8	0.6739 (3)	0.8530 (3)	0.12494 (18)	0.0517 (7)
H8A	0.7240	0.9121	0.1002	0.077*
H8B	0.6249	0.8799	0.1697	0.077*
H8C	0.6042	0.8222	0.0902	0.077*
C9	1.1021 (3)	0.8555 (2)	0.06741 (14)	0.0400 (6)
C10	1.0719 (4)	1.0403 (2)	0.0096 (2)	0.0595 (8)
H10	0.9725	1.0328	0.0290	0.071*
C11	1.1403 (6)	1.1369 (3)	0.0467 (2)	0.0941 (16)
H11A	1.2421	1.1404	0.0324	0.113*
H11B	1.1351	1.1288	0.1019	0.113*
C12	1.0651 (7)	1.2431 (3)	0.0232 (2)	0.0936 (15)
H12A	0.9671	1.2436	0.0437	0.112*
H12B	1.1171	1.3044	0.0452	0.112*
C13	1.0586 (6)	1.2563 (3)	-0.0593 (2)	0.0750 (11)
H13A	1.1561	1.2688	-0.0786	0.090*
H13B	1.0007	1.3203	-0.0712	0.090*
C14	0.9968 (8)	1.1623 (3)	-0.0982 (3)	0.123 (2)
H14A	1.0086	1.1717	-0.1531	0.147*

H14B	0.8935	1.1592	-0.0875	0.147*
C15	1.0660 (8)	1.0544 (3)	-0.0745 (2)	0.115 (2)
H15A	1.0109	0.9947	-0.0967	0.138*
H15B	1.1639	1.0509	-0.0950	0.138*

*Atomic displacement parameters (Å<sup>2</sup>)*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Sn1	0.03555 (9)	0.03065 (8)	0.04623 (9)	0.00216 (7)	-0.00566 (7)	0.00345 (7)
C11	0.0458 (4)	0.0469 (4)	0.0550 (4)	0.0002 (3)	-0.0045 (3)	-0.0088 (3)
C12	0.0986 (7)	0.0686 (5)	0.0500 (4)	-0.0184 (5)	-0.0149 (4)	-0.0065 (4)
S1	0.0341 (3)	0.0362 (3)	0.0656 (4)	0.0016 (3)	0.0043 (3)	0.0044 (3)
N1	0.0436 (13)	0.0366 (11)	0.0431 (11)	-0.0039 (10)	-0.0010 (9)	0.0059 (9)
N2	0.0395 (12)	0.0274 (10)	0.0383 (10)	0.0034 (9)	0.0030 (9)	0.0019 (8)
N3	0.0371 (11)	0.0332 (10)	0.0491 (11)	0.0028 (10)	0.0050 (9)	0.0071 (10)
N4	0.0442 (15)	0.0431 (14)	0.093 (2)	0.0067 (11)	0.0179 (14)	0.0239 (14)
C1	0.060 (2)	0.0515 (19)	0.095 (3)	0.0150 (17)	-0.021 (2)	0.0203 (18)
C2	0.061 (2)	0.0445 (16)	0.0518 (16)	-0.0074 (15)	-0.0060 (15)	0.0125 (13)
C3	0.067 (2)	0.058 (2)	0.0554 (17)	-0.0250 (18)	-0.0023 (16)	0.0152 (15)
C4	0.0554 (18)	0.067 (2)	0.0543 (16)	-0.0200 (19)	0.0087 (16)	0.0020 (14)
C5	0.0451 (17)	0.0505 (17)	0.0513 (15)	-0.0044 (12)	0.0106 (13)	-0.0029 (12)
C6	0.0389 (14)	0.0353 (13)	0.0375 (12)	-0.0017 (11)	0.0023 (11)	-0.0067 (10)
C7	0.0371 (13)	0.0348 (12)	0.0371 (13)	0.0036 (10)	0.0032 (10)	-0.0028 (10)
C8	0.0415 (15)	0.0502 (17)	0.0632 (16)	0.0141 (14)	0.0045 (13)	0.0060 (14)
C9	0.0426 (14)	0.0317 (13)	0.0458 (13)	-0.0002 (12)	0.0011 (10)	0.0017 (11)
C10	0.0435 (15)	0.0419 (15)	0.093 (2)	0.0065 (15)	0.0162 (18)	0.0256 (15)
C11	0.159 (5)	0.059 (2)	0.064 (2)	0.046 (3)	-0.036 (3)	-0.0167 (19)
C12	0.154 (4)	0.052 (2)	0.075 (2)	0.032 (3)	-0.021 (3)	-0.0103 (19)
C13	0.092 (3)	0.0485 (18)	0.085 (2)	0.013 (2)	0.013 (2)	0.0252 (17)
C14	0.214 (7)	0.071 (3)	0.083 (3)	0.057 (3)	-0.064 (4)	-0.008 (2)
C15	0.197 (6)	0.058 (2)	0.090 (3)	0.049 (3)	-0.067 (4)	-0.022 (2)

*Geometric parameters (Å, °)*

Sn1—C1	2.136 (3)	C5—C6	1.383 (4)
Sn1—N1	2.269 (2)	C5—H5	0.9300
Sn1—N2	2.224 (2)	C6—C7	1.473 (4)
Sn1—S1	2.4814 (7)	C7—C8	1.483 (4)
Sn1—C11	2.4960 (7)	C8—H8A	0.9600
Sn1—C12	2.4701 (8)	C8—H8B	0.9600
S1—C9	1.753 (3)	C8—H8C	0.9600
N1—C2	1.332 (4)	C10—C15	1.479 (5)
N1—C6	1.348 (3)	C10—C11	1.488 (6)
N2—C7	1.292 (3)	C10—H10	0.9800
N2—N3	1.350 (3)	C11—C12	1.529 (5)
N3—C9	1.326 (3)	C11—H11A	0.9700
N4—C9	1.330 (4)	C11—H11B	0.9700
N4—C10	1.468 (4)	C12—C13	1.450 (5)

N4—H4	0.861 (10)	C12—H12A	0.9700
C1—H1A	0.9600	C12—H12B	0.9700
C1—H1B	0.9600	C13—C14	1.453 (6)
C1—H1C	0.9600	C13—H13A	0.9700
C2—C3	1.383 (5)	C13—H13B	0.9700
C2—H2	0.9300	C14—C15	1.523 (5)
C3—C4	1.359 (5)	C14—H14A	0.9700
C3—H3	0.9300	C14—H14B	0.9700
C4—C5	1.372 (4)	C15—H15A	0.9700
C4—H4A	0.9300	C15—H15B	0.9700
C1—Sn1—N2	171.66 (12)	N2—C7—C8	122.8 (2)
C1—Sn1—N1	99.56 (12)	C6—C7—C8	121.1 (2)
N2—Sn1—N1	72.12 (8)	C7—C8—H8A	109.5
C1—Sn1—C12	93.43 (11)	C7—C8—H8B	109.5
N2—Sn1—C12	86.65 (6)	H8A—C8—H8B	109.5
N1—Sn1—C12	85.57 (6)	C7—C8—H8C	109.5
C1—Sn1—S1	109.56 (11)	H8A—C8—H8C	109.5
N2—Sn1—S1	78.74 (6)	H8B—C8—H8C	109.5
N1—Sn1—S1	150.85 (6)	N3—C9—N4	117.1 (3)
C12—Sn1—S1	93.69 (3)	N3—C9—S1	127.9 (2)
C1—Sn1—C11	92.01 (11)	N4—C9—S1	115.0 (2)
N2—Sn1—C11	86.94 (6)	N4—C10—C15	110.0 (3)
N1—Sn1—C11	86.53 (6)	N4—C10—C11	110.1 (3)
C12—Sn1—C11	171.05 (3)	C15—C10—C11	110.8 (3)
S1—Sn1—C11	91.18 (3)	N4—C10—H10	108.6
C9—S1—Sn1	95.70 (9)	C15—C10—H10	108.6
C2—N1—C6	120.1 (3)	C11—C10—H10	108.6
C2—N1—Sn1	124.3 (2)	C10—C11—C12	111.5 (3)
C6—N1—Sn1	115.58 (17)	C10—C11—H11A	109.3
C7—N2—N3	118.5 (2)	C12—C11—H11A	109.3
C7—N2—Sn1	119.51 (17)	C10—C11—H11B	109.3
N3—N2—Sn1	121.94 (16)	C12—C11—H11B	109.3
C9—N3—N2	115.7 (2)	H11A—C11—H11B	108.0
C9—N4—C10	126.0 (3)	C13—C12—C11	112.3 (3)
C9—N4—H4	123 (2)	C13—C12—H12A	109.1
C10—N4—H4	111 (2)	C11—C12—H12A	109.1
Sn1—C1—H1A	109.5	C13—C12—H12B	109.1
Sn1—C1—H1B	109.5	C11—C12—H12B	109.1
H1A—C1—H1B	109.5	H12A—C12—H12B	107.9
Sn1—C1—H1C	109.5	C12—C13—C14	113.1 (4)
H1A—C1—H1C	109.5	C12—C13—H13A	109.0
H1B—C1—H1C	109.5	C14—C13—H13A	109.0
N1—C2—C3	121.5 (3)	C12—C13—H13B	109.0
N1—C2—H2	119.3	C14—C13—H13B	109.0
C3—C2—H2	119.3	H13A—C13—H13B	107.8
C4—C3—C2	119.0 (3)	C13—C14—C15	113.3 (4)
C4—C3—H3	120.5	C13—C14—H14A	108.9

C2—C3—H3	120.5	C15—C14—H14A	108.9
C3—C4—C5	119.6 (3)	C13—C14—H14B	108.9
C3—C4—H4A	120.2	C15—C14—H14B	108.9
C5—C4—H4A	120.2	H14A—C14—H14B	107.7
C4—C5—C6	119.7 (3)	C10—C15—C14	112.7 (4)
C4—C5—H5	120.1	C10—C15—H15A	109.0
C6—C5—H5	120.1	C14—C15—H15A	109.0
N1—C6—C5	120.1 (2)	C10—C15—H15B	109.0
N1—C6—C7	116.6 (2)	C14—C15—H15B	109.0
C5—C6—C7	123.4 (2)	H15A—C15—H15B	107.8
N2—C7—C6	116.1 (2)		
C1—Sn1—S1—C9	-178.73 (15)	C3—C4—C5—C6	-1.9 (5)
N2—Sn1—S1—C9	2.07 (10)	C2—N1—C6—C5	1.0 (4)
N1—Sn1—S1—C9	3.74 (16)	Sn1—N1—C6—C5	-176.8 (2)
Cl2—Sn1—S1—C9	-83.77 (9)	C2—N1—C6—C7	180.0 (2)
Cl1—Sn1—S1—C9	88.72 (9)	Sn1—N1—C6—C7	2.1 (3)
C1—Sn1—N1—C2	2.9 (3)	C4—C5—C6—N1	0.8 (4)
N2—Sn1—N1—C2	-177.8 (2)	C4—C5—C6—C7	-178.0 (3)
Cl2—Sn1—N1—C2	-89.9 (2)	N3—N2—C7—C6	-176.5 (2)
S1—Sn1—N1—C2	-179.50 (18)	Sn1—N2—C7—C6	4.4 (3)
Cl1—Sn1—N1—C2	94.3 (2)	N3—N2—C7—C8	2.6 (4)
C1—Sn1—N1—C6	-179.4 (2)	Sn1—N2—C7—C8	-176.5 (2)
N2—Sn1—N1—C6	-0.01 (17)	N1—C6—C7—N2	-4.3 (3)
Cl2—Sn1—N1—C6	87.89 (18)	C5—C6—C7—N2	174.6 (2)
S1—Sn1—N1—C6	-1.7 (3)	N1—C6—C7—C8	176.6 (2)
Cl1—Sn1—N1—C6	-87.92 (18)	C5—C6—C7—C8	-4.5 (4)
C1—Sn1—N2—C7	1.9 (9)	N2—N3—C9—N4	-179.0 (3)
N1—Sn1—N2—C7	-2.46 (18)	N2—N3—C9—S1	1.1 (3)
Cl2—Sn1—N2—C7	-88.88 (19)	C10—N4—C9—N3	2.6 (5)
S1—Sn1—N2—C7	176.68 (19)	C10—N4—C9—S1	-177.5 (3)
Cl1—Sn1—N2—C7	84.87 (19)	Sn1—S1—C9—N3	-2.6 (3)
C1—Sn1—N2—N3	-177.3 (8)	Sn1—S1—C9—N4	177.5 (2)
N1—Sn1—N2—N3	178.4 (2)	C9—N4—C10—C15	-120.0 (5)
Cl2—Sn1—N2—N3	91.99 (17)	C9—N4—C10—C11	117.6 (4)
S1—Sn1—N2—N3	-2.45 (17)	N4—C10—C11—C12	175.8 (4)
Cl1—Sn1—N2—N3	-94.26 (17)	C15—C10—C11—C12	53.9 (6)
C7—N2—N3—C9	-177.5 (2)	C10—C11—C12—C13	-54.4 (7)
Sn1—N2—N3—C9	1.6 (3)	C11—C12—C13—C14	52.4 (7)
C6—N1—C2—C3	-1.8 (4)	C12—C13—C14—C15	-50.4 (8)
Sn1—N1—C2—C3	175.8 (2)	N4—C10—C15—C14	-174.0 (5)
N1—C2—C3—C4	0.7 (5)	C11—C10—C15—C14	-52.0 (7)
C2—C3—C4—C5	1.2 (5)	C13—C14—C15—C10	50.5 (8)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
---------------	-------	-------------	-------------	---------------



---

N4—H4 $\cdots$ C11 <sup>i</sup>	0.86 (1)	2.36 (1)	3.219 (3)	177 (3)
---------------------------------	----------	----------	-----------	---------

---

Symmetry code: (i)  $x+1/2, -y+3/2, -z$ .