

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

9-Benzyl-9H-carbazole

 Nesimi Uludağ,^a Murat Ateş,^a Barış Tercan,^b Emel Ermiş^c
 and Tuncer Hökelek^{d*}
^aNamik Kemal University, Faculty of Arts and Sciences, Department of Chemistry, 59100 Tekirdağ, Turkey, ^bKarabük University, Department of Physics, 78050 Karabük, Turkey, ^cAnadolu University, Faculty of Science, Department of Chemistry, 26470 Yenibağlar, Eskişehir, Turkey, and ^dHacettepe University, Department of Physics, 06800 Beytepe, Ankara, Turkey

Correspondence e-mail: merzifon@hacettepe.edu.tr


Received 26 March 2010; accepted 2 April 2010

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.076; wR factor = 0.210; data-to-parameter ratio = 14.4.

The asymmetric unit of the title compound, $\text{C}_{19}\text{H}_{15}\text{N}$, contains two crystallographically independent molecules. In both molecules, the planar carbazole moieties [maximum deviations = 0.037 (4) and 0.042 (3) Å] are oriented with respect to the adjacent benzene rings, at dihedral angles of 85.29 (8) and 89.89 (7)°, respectively. In the crystal structure, weak C—H... π interactions are observed involving the carbazole rings.

Related literature

For tetrahydrocarbazole systems present in the framework of a number of indole-type alkaloids of biological interest, see: Phillipson & Zenk (1980); Saxton (1983); Abraham (1975). For related structures, see: Hökelek *et al.* (1994, 1998, 1999, 2004, 2006); Patır *et al.* (1997); Hökelek & Patır (1999, 2002); Çaylak *et al.* (2007). For bond-length data, see: Allen *et al.* (1987).



Experimental

Crystal data

 $\text{C}_{19}\text{H}_{15}\text{N}$
 $M_r = 257.32$
 Monoclinic, $P2_1/c$
 $a = 14.9305$ (4) Å
 $b = 5.5612$ (2) Å
 $c = 32.7916$ (8) Å
 $\beta = 94.518$ (3)°

Data collection

 $V = 2714.27$ (14) Å³
 $Z = 8$
 Mo $K\alpha$ radiation
 $\mu = 0.07$ mm⁻¹
 $T = 100$ K
 $0.27 \times 0.15 \times 0.14$ mm

 Bruker Kappa APEXII CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Bruker, 2005)
 $T_{\min} = 0.981$, $T_{\max} = 0.990$

 24870 measured reflections
 6816 independent reflections
 3384 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.103$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.076$
 $wR(F^2) = 0.210$
 $S = 1.03$
 6816 reflections
 474 parameters

 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.27$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.26$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C6}-\text{H6}\cdots\text{Cg1}^{\text{i}}$	0.97 (4)	2.940 (4)	3.636 (5)	129.46 (5)
$\text{C10}'-\text{H10C}\cdots\text{Cg1}^{\text{ii}}$	0.98 (3)	2.787 (4)	3.700 (5)	154.92 (4)
$\text{C4}'-\text{H4}'\cdots\text{Cg3}^{\text{i}}$	0.99 (4)	2.706 (4)	3.554 (4)	144.36 (5)

 Symmetry codes: (i) $-x, y - \frac{1}{2}, -z + \frac{1}{2}$; (ii) $x, y + 1, z$. $\text{Cg1}'$ and Cg3 are the centroids of the $\text{C1}'-\text{C4}'/\text{C4A}'/\text{C9A}'$ and $\text{C5A}/\text{C5}-\text{C8}/\text{C8A}$ rings, respectively.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997); software used to prepare material for publication: WinGX (Farrugia, 1999) and PLATON (Spek, 2009).

The authors are indebted to Anadolu University and the Medicinal Plants and Medicine Research Centre of Anadolu University, Eskişehir, Turkey, for the use of the X-ray diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2745).

References

- Abraham, D. J. (1975). *The Catharanthus Alkaloids*, edited by W. I. Taylor & N. R. Fransworth, chs. 7 and 8. New York: Marcel Dekker.
- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bruker (2005). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Çaylak, N., Hökelek, T., Uludağ, N. & Patır, S. (2007). *Acta Cryst. E63*, o3913–o3914.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Hökelek, T., Gündüz, H., Patır, S. & Uludağ, N. (1998). *Acta Cryst. C54*, 1297–1299.
- Hökelek, T. & Patır, S. (1999). *Acta Cryst. C55*, 675–677.
- Hökelek, T. & Patır, S. (2002). *Acta Cryst. E58*, o374–o376.
- Hökelek, T., Patır, S., Gülce, A. & Okay, G. (1994). *Acta Cryst. C50*, 450–453.
- Hökelek, T., Patır, S. & Uludağ, N. (1999). *Acta Cryst. C55*, 114–116.
- Hökelek, T., Uludağ, N. & Patır, S. (2004). *Acta Cryst. E60*, o25–o27.
- Hökelek, T., Uludağ, N. & Patır, S. (2006). *Acta Cryst. E62*, o791–o793.
- Patır, S., Okay, G., Gülce, A., Salih, B. & Hökelek, T. (1997). *J. Heterocycl. Chem.* **34**, 1239–1242.
- Phillipson, J. D. & Zenk, M. H. (1980). *Indole and Biogenetically Related Alkaloids*, ch. 3. New York: Academic Press.
- Saxton, J. E. (1983). Editor. *Heterocyclic Compounds*, Vol. 25, *The Monoterpenoid Indole Alkaloids*, chs. 8 and 11. New York: Wiley.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

supporting information

Acta Cryst. (2010). E66, o1077 [https://doi.org/10.1107/S1600536810012444]

9-Benzyl-9H-carbazole

Nesimi Uludağ, Murat Ateş, Barış Tercan, Emel Ermiş and Tuncer Hökelek

S1. Comment

Tetrahydrocarbazole systems are present in the framework of a number of indole-type alkaloids of biological interest (Phillipson & Zenk, 1980; Saxton, 1983; Abraham, 1975). The structures of tricyclic, tetracyclic and pentacyclic ring systems with dithiolane and other substituents of the tetrahydrocarbazole core, have been the subject of much interest in our laboratory. These include 1,2,3,4-tetrahydrocarbazole-1-spiro-2'-[1,3]dithiolane, (II) (Hökelek *et al.*, 1994), *N*-(2-methoxyethyl)-*N*-{2,3,4,9-tetrahydrospiro[1*H*-carbazole-1, 2-(1,3)dithiolane]-4-yl}benzene-sulfonamide, (III) (Patır *et al.*, 1997), spiro[carbazole-1(2*H*),2'-[1,3]-dithiolan]-4(3*H*)-one, (IV) (Hökelek *et al.*, 1998), 9-acetonyl-3-ethylidene-1,2,3,4-tetrahydrospiro[carbazole-1,2'-[1,3] dithiolan]-4-one, (V) (Hökelek *et al.*, 1999), *N*-(2,2-dimethoxyethyl)-*N*-{9-methoxymethyl-1,2,3,4-tetrahydrospiro[carbazole-1,2'-[1,3]dithiolan] -4-yl}benzamide, (VI) (Hökelek & Patır, 1999), 3a,4,10,10 b-tetrahydro-2*H* -furo[2,3-*a*]carbazol-5(3*H*)-one, (VII) (Çaylak *et al.*, 2007); also the pentacyclic compounds 6-ethyl-4-(2-methoxyethyl)-2,6-methano-5-oxo-hexahydro- pyrrolo(2,3-*d*)carbazole-1-spiro-2'-(1,3)dithiolane, (VIII) (Hökelek & Patır, 2002), *N*-(2-benzyloxyethyl)-4,7-dimethyl-6-(1,3-dithiolan-2-yl)-1,2, 3,4,5,6-hexahydro-1,5-methano-2-azocino[4,3-*b*]indol-2-one, (IX) (Hökelek *et al.*, 2004) and 4-ethyl-6,6-ethylenedithio-2-(2-methoxyethyl)-7-methoxy- methylene-2,3,4,5,6,7-hexahydro-1,5-methano-1*H*-azocino[4,3-*b*]indol-3-one, (X) (Hökelek *et al.*, 2006). The title compound, (I), may be considered as a synthetic precursor of tetracyclic indole alkaloids of biological interests. The present study was undertaken to ascertain its crystal structure.

The title compound consists of a carbazole skeleton with a benzyl group. Its asymmetric unit, (Fig. 1), contains two crystallographically independent molecules, where the bond lengths (Allen *et al.*, 1987) and angles are within normal ranges, and generally agree with those in compounds (II)-(X). In all structures atom N9 is substituted.

An examination of the deviations from the least-squares planes through individual rings shows that rings A (C1—C4/C4a/C9a), B (C4a/C5a/C8a/N9/C9a), C (C5a/C5—C8/C8a), D (C11—C16) and A' (C1'-C4'/C4a'/C9a'), B' (C4a'/C5a'/ C8a'/N9'/C9a'), C' (C5a'/C5'-C8'/C8a'), D' (C11'-C16') are planar. The carbazole skeletons, containing the rings A, B, C and A', B', C' are also nearly coplanar [with a maximum deviations of 0.037 (4) and 0.042 (3) Å for atoms C2 and C7', respectively] with dihedral angles of A/B = 1.28 (10), A/C = 1.57 (9), B/C = 0.32 (7) ° and A'/B' = 0.94 (10), A'/C' = 2.37 (10), B'/C' = 1.72 (11) °. Rings D and D' are oriented with respect to the planar carbazole skeletons at dihedral angles of 85.29 (8) and 89.89 (7) °, respectively. Atoms C10 and C10' displaced by -0.109 (3), -0.005 (4) Å and -0.016 (3), -0.098 (3) Å from the planes of the corresponding carbazole skeletons and benzene rings, respectively.

In the crystal structure, three weak C—H... π interactions (Table 1) involving the carbazole rings are observed.

S2. Experimental

For the preparation of the title compound, (I), sodium hydride (2.38 g, 59.85 mmol) was added to a solution of carbazole (5.00 g, 29.92 mmol) in dry tetrahydrofuran (200 ml) in several portions, and stirred at room temperature for 1 h under argon atmosphere. Then, benzylchloride (5.68 g, 44.88 mmol) was added and stirred at 343 K for 6 h. The reaction

mixture was cooled in an ice bath, and hydrochloric acid (10%, 200 ml) was added. After the extraction with dichloromethane (300 ml), the organic layer was dried over anhydrous magnesium sulfate and the solvent was evaporated under reduced pressure. The residue was purified by column chromatography using silica gel and dichloromethane-petroleum ether (1:1), and the product was recrystallized from diethyl ether and cyclohexane mixture (1:1) (yield; 4.00 g, 80%, m.p. 388 K).

S3. Refinement

H3 and H7' atoms were positioned geometrically, with C—H = 0.93 Å for aromatic H atoms and constrained to ride on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. The remaining H atoms were located in difference synthesis and refined isotropically.




Figure 1

The molecular structure of the title molecule with the atom-numbering scheme. The displacement ellipsoids are drawn at the 50% probability level.

9-Benzyl-9H-carbazole

Crystal data

$\text{C}_{19}\text{H}_{15}\text{N}$
 $M_r = 257.32$
 Monoclinic, $P2_1/c$
 Hall symbol: $-P 2_1/c$
 $a = 14.9305 (4) \text{ \AA}$
 $b = 5.5612 (2) \text{ \AA}$
 $c = 32.7916 (8) \text{ \AA}$
 $\beta = 94.518 (3)^\circ$
 $V = 2714.27 (14) \text{ \AA}^3$
 $Z = 8$

$F(000) = 1088$
 $D_x = 1.259 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 1669 reflections
 $\theta = 2.5\text{--}22.9^\circ$
 $\mu = 0.07 \text{ mm}^{-1}$
 $T = 100 \text{ K}$
 Block, colorless
 $0.27 \times 0.15 \times 0.14 \text{ mm}$

Data collection

Bruker Kappa APEXII CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Bruker, 2005)

$T_{\min} = 0.981$, $T_{\max} = 0.990$

24870 measured reflections

6816 independent reflections

3384 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.103$

$\theta_{\max} = 28.4^\circ$, $\theta_{\min} = 1.3^\circ$

$h = -19 \rightarrow 18$

$k = -7 \rightarrow 7$

$l = -42 \rightarrow 43$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.076$

$wR(F^2) = 0.210$

$S = 1.03$

6816 reflections

474 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0918P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.27 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$

Extinction correction: *SHELXL97* (Sheldrick,
2008), $F_c^* = kFc^*[1 + 0.001x\text{Fc}^2\lambda^3/\sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0068 (11)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.5696 (2)	0.6160 (7)	0.83884 (11)	0.0399 (9)
H1	0.536 (2)	0.451 (6)	0.8446 (9)	0.042 (9)*
C2	0.5654 (2)	0.7192 (8)	0.80036 (11)	0.0495 (10)
H2	0.524 (2)	0.643 (6)	0.7769 (10)	0.040 (9)*
C3	0.6133 (2)	0.9241 (8)	0.79216 (10)	0.0486 (10)
H3	0.6077	0.9890	0.7659	0.058*
C4	0.6692 (2)	1.0341 (7)	0.82209 (10)	0.0364 (8)
H4	0.702 (2)	1.169 (6)	0.8155 (9)	0.030 (9)*
C4A	0.67613 (19)	0.9336 (6)	0.86118 (9)	0.0286 (7)
C5	0.7898 (2)	1.1740 (6)	0.91002 (10)	0.0286 (7)
H5	0.807 (2)	1.303 (6)	0.8898 (10)	0.052 (10)*
C5A	0.72814 (19)	0.9919 (5)	0.89881 (8)	0.0244 (6)
C6	0.8284 (2)	1.1795 (6)	0.94938 (10)	0.0308 (7)
H6	0.870 (2)	1.308 (6)	0.9577 (9)	0.045 (10)*
C7	0.8064 (2)	1.0064 (6)	0.97798 (10)	0.0294 (7)

H7	0.836 (2)	1.010 (6)	1.0055 (10)	0.037 (9)*
C8	0.7450 (2)	0.8268 (6)	0.96804 (9)	0.0284 (7)
H8	0.727 (2)	0.696 (6)	0.9883 (10)	0.045 (9)*
C8A	0.70640 (19)	0.8208 (5)	0.92781 (9)	0.0243 (6)
N9	0.64413 (16)	0.6597 (4)	0.90964 (7)	0.0263 (6)
C9A	0.62622 (19)	0.7261 (5)	0.86924 (9)	0.0274 (7)
C10	0.5991 (2)	0.4654 (6)	0.92981 (11)	0.0314 (7)
H10A	0.604 (2)	0.313 (6)	0.9142 (9)	0.036 (9)*
H10B	0.632 (2)	0.433 (5)	0.9553 (9)	0.031 (9)*
C11	0.50251 (19)	0.5224 (5)	0.93644 (9)	0.0260 (7)
C12	0.4798 (2)	0.7304 (6)	0.95693 (10)	0.0335 (8)
H12	0.527 (2)	0.852 (6)	0.9672 (10)	0.048 (10)*
C13	0.3908 (2)	0.7795 (6)	0.96297 (10)	0.0348 (8)
H13	0.375 (2)	0.931 (6)	0.9762 (10)	0.052 (10)*
C14	0.3239 (2)	0.6231 (6)	0.94863 (9)	0.0311 (7)
H14	0.263 (3)	0.675 (6)	0.9533 (10)	0.054 (11)*
C15	0.3458 (2)	0.4166 (6)	0.92833 (10)	0.0337 (8)
H15	0.296 (2)	0.295 (6)	0.9180 (9)	0.038 (9)*
C16	0.4348 (2)	0.3666 (6)	0.92232 (10)	0.0305 (7)
H16	0.4532 (19)	0.217 (5)	0.9088 (8)	0.027 (8)*
C1'	0.9099 (2)	0.8115 (6)	0.57706 (9)	0.0291 (7)
H1'	0.865 (2)	0.673 (5)	0.5719 (8)	0.032 (8)*
C2'	0.9229 (2)	0.9787 (6)	0.54711 (10)	0.0333 (8)
H2'	0.883 (2)	0.951 (5)	0.5217 (9)	0.029 (8)*
C3'	0.9879 (2)	1.1600 (6)	0.55305 (10)	0.0345 (8)
H3'	0.998 (2)	1.271 (6)	0.5318 (9)	0.035 (9)*
C4'	1.0415 (2)	1.1763 (6)	0.58923 (10)	0.0294 (7)
H4'	1.085 (3)	1.310 (7)	0.5918 (11)	0.063 (12)*
C4A'	1.02988 (19)	1.0096 (5)	0.61992 (9)	0.0258 (7)
C5'	1.1437 (2)	1.0841 (6)	0.68366 (10)	0.0325 (8)
H5'	1.177 (2)	1.222 (6)	0.6724 (9)	0.041 (9)*
C5A'	1.07331 (19)	0.9688 (5)	0.66042 (9)	0.0258 (7)
C6'	1.1713 (2)	0.9928 (7)	0.72151 (10)	0.0399 (8)
H6'	1.223 (3)	1.067 (7)	0.7361 (11)	0.061 (11)*
C7'	1.1302 (2)	0.7899 (6)	0.73662 (9)	0.0374 (8)
H7'	1.1509	0.7306	0.7622	0.045*
C8'	1.0594 (2)	0.6735 (6)	0.71486 (9)	0.0305 (7)
H8'	1.028 (2)	0.519 (6)	0.7270 (9)	0.043 (9)*
C8A'	1.03215 (19)	0.7642 (5)	0.67661 (8)	0.0249 (7)
N9'	0.96646 (16)	0.6806 (4)	0.64747 (7)	0.0255 (6)
C9A'	0.96460 (19)	0.8272 (5)	0.61347 (9)	0.0258 (7)
C10'	0.9071 (2)	0.4776 (5)	0.65181 (10)	0.0267 (7)
H10C	0.910 (2)	0.373 (6)	0.6277 (10)	0.042 (10)*
H10D	0.932 (2)	0.372 (5)	0.6747 (9)	0.033 (9)*
C11'	0.81229 (19)	0.5454 (5)	0.66032 (8)	0.0241 (6)
C12'	0.7931 (2)	0.7589 (6)	0.67972 (9)	0.0314 (7)
H12'	0.847 (2)	0.884 (6)	0.6895 (9)	0.047 (10)*
C13'	0.7052 (2)	0.8047 (7)	0.68913 (11)	0.0401 (9)

H13'	0.690 (2)	0.951 (6)	0.7018 (9)	0.042 (10)*
C14'	0.6376 (2)	0.6429 (7)	0.67937 (12)	0.0481 (10)
H14'	0.576 (3)	0.680 (7)	0.6901 (12)	0.075 (13)*
C15'	0.6564 (2)	0.4339 (7)	0.65933 (12)	0.0462 (10)
H15'	0.605 (3)	0.327 (6)	0.6506 (10)	0.054 (11)*
C16'	0.7434 (2)	0.3849 (6)	0.64961 (10)	0.0350 (8)
H16'	0.758 (2)	0.242 (6)	0.6364 (10)	0.041 (10)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0220 (17)	0.053 (2)	0.045 (2)	-0.0002 (16)	0.0017 (14)	-0.0173 (18)
C2	0.029 (2)	0.085 (3)	0.034 (2)	0.009 (2)	-0.0012 (16)	-0.015 (2)
C3	0.031 (2)	0.086 (3)	0.0284 (18)	0.015 (2)	0.0033 (15)	0.0019 (19)
C4	0.0239 (18)	0.055 (2)	0.0316 (19)	0.0087 (17)	0.0072 (14)	0.0043 (17)
C4A	0.0183 (15)	0.0393 (18)	0.0288 (17)	0.0061 (14)	0.0056 (12)	-0.0019 (14)
C5	0.0219 (16)	0.0278 (16)	0.0372 (19)	0.0003 (13)	0.0090 (13)	-0.0011 (14)
C5A	0.0196 (15)	0.0290 (15)	0.0254 (15)	0.0057 (13)	0.0075 (12)	0.0015 (13)
C6	0.0225 (17)	0.0326 (17)	0.0379 (19)	-0.0054 (14)	0.0062 (14)	-0.0049 (15)
C7	0.0226 (16)	0.0366 (17)	0.0291 (17)	-0.0006 (14)	0.0024 (13)	-0.0058 (15)
C8	0.0247 (17)	0.0331 (16)	0.0280 (17)	0.0018 (14)	0.0059 (13)	0.0017 (14)
C8A	0.0172 (15)	0.0259 (15)	0.0308 (16)	0.0018 (12)	0.0074 (12)	-0.0026 (13)
N9	0.0188 (13)	0.0243 (12)	0.0358 (15)	-0.0004 (10)	0.0034 (10)	-0.0009 (11)
C9A	0.0153 (15)	0.0365 (17)	0.0305 (17)	0.0066 (13)	0.0030 (12)	-0.0057 (14)
C10	0.0203 (16)	0.0262 (17)	0.048 (2)	-0.0022 (13)	0.0043 (15)	0.0017 (16)
C11	0.0223 (16)	0.0238 (14)	0.0321 (16)	0.0021 (13)	0.0025 (12)	0.0043 (13)
C12	0.0273 (18)	0.0325 (17)	0.0408 (19)	-0.0022 (15)	0.0035 (14)	-0.0034 (15)
C13	0.0306 (19)	0.0331 (18)	0.042 (2)	0.0030 (15)	0.0097 (14)	-0.0013 (16)
C14	0.0210 (17)	0.0367 (18)	0.0360 (18)	0.0040 (15)	0.0046 (13)	0.0088 (15)
C15	0.0236 (17)	0.0416 (19)	0.0357 (18)	-0.0096 (15)	0.0011 (14)	0.0044 (16)
C16	0.0252 (17)	0.0284 (17)	0.0378 (19)	-0.0031 (14)	0.0021 (13)	-0.0021 (14)
C1'	0.0205 (16)	0.0371 (18)	0.0298 (17)	0.0043 (14)	0.0024 (13)	0.0036 (15)
C2'	0.0202 (16)	0.048 (2)	0.0318 (18)	0.0100 (15)	0.0054 (13)	0.0058 (16)
C3'	0.0312 (19)	0.0403 (19)	0.0336 (19)	0.0107 (15)	0.0126 (15)	0.0119 (16)
C4'	0.0245 (17)	0.0282 (16)	0.0376 (19)	0.0042 (14)	0.0154 (14)	0.0013 (14)
C4A'	0.0193 (15)	0.0261 (15)	0.0333 (16)	0.0024 (13)	0.0099 (12)	-0.0013 (13)
C5'	0.0248 (17)	0.0365 (18)	0.0378 (19)	-0.0033 (15)	0.0115 (14)	-0.0101 (15)
C5A'	0.0202 (15)	0.0294 (15)	0.0287 (16)	0.0016 (13)	0.0081 (12)	-0.0049 (13)
C6'	0.0304 (19)	0.054 (2)	0.0357 (19)	-0.0029 (18)	0.0048 (15)	-0.0164 (18)
C7'	0.0321 (19)	0.055 (2)	0.0249 (17)	0.0100 (17)	0.0017 (13)	-0.0072 (16)
C8'	0.0268 (17)	0.0378 (18)	0.0274 (17)	0.0045 (14)	0.0042 (13)	0.0007 (15)
C8A'	0.0199 (15)	0.0291 (16)	0.0263 (16)	0.0024 (13)	0.0060 (12)	-0.0036 (13)
N9'	0.0202 (13)	0.0284 (13)	0.0280 (13)	-0.0010 (11)	0.0030 (10)	0.0035 (11)
C9A'	0.0197 (15)	0.0291 (15)	0.0293 (16)	0.0026 (13)	0.0070 (12)	0.0011 (13)
C10'	0.0214 (16)	0.0252 (15)	0.0344 (18)	-0.0027 (13)	0.0075 (13)	-0.0020 (15)
C11'	0.0229 (15)	0.0278 (15)	0.0219 (15)	0.0012 (13)	0.0041 (12)	0.0057 (13)
C12'	0.0312 (18)	0.0352 (18)	0.0288 (17)	0.0031 (15)	0.0092 (13)	0.0028 (14)
C13'	0.039 (2)	0.043 (2)	0.042 (2)	0.0130 (18)	0.0205 (16)	0.0135 (18)

C14'	0.027 (2)	0.063 (3)	0.056 (2)	0.0101 (19)	0.0134 (17)	0.031 (2)
C15'	0.0245 (19)	0.051 (2)	0.062 (2)	-0.0046 (18)	-0.0030 (17)	0.022 (2)
C16'	0.0276 (18)	0.0325 (18)	0.044 (2)	-0.0049 (15)	-0.0042 (15)	0.0083 (16)

Geometric parameters (Å, °)

C1—C2	1.383 (5)	C1'—C2'	1.377 (4)
C1—H1	1.07 (3)	C1'—H1'	1.03 (3)
C2—H2	1.04 (3)	C2'—H2'	0.99 (3)
C3—C2	1.383 (6)	C3'—C2'	1.403 (5)
C3—H3	0.9300	C3'—H3'	0.95 (3)
C4—C3	1.380 (5)	C4'—C3'	1.381 (5)
C4—H4	0.93 (3)	C4'—H4'	0.99 (4)
C4A—C4	1.395 (4)	C4A'—C4'	1.389 (4)
C5—H5	1.02 (4)	C4A'—C5A'	1.449 (4)
C5A—C4A	1.442 (4)	C5'—C6'	1.374 (5)
C5A—C5	1.398 (4)	C5'—H5'	1.00 (3)
C6—C5	1.372 (4)	C5A'—C5'	1.404 (4)
C6—C7	1.401 (4)	C6'—H6'	0.97 (4)
C6—H6	0.97 (4)	C7'—C6'	1.394 (5)
C7—H7	0.97 (3)	C7'—H7'	0.9300
C8—C7	1.377 (4)	C8'—C7'	1.387 (4)
C8—H8	1.04 (3)	C8'—H8'	1.07 (3)
C8A—C5A	1.401 (4)	C8A'—C5A'	1.416 (4)
C8A—C8	1.398 (4)	C8A'—C8'	1.383 (4)
N9—C8A	1.391 (4)	N9'—C8A'	1.394 (4)
N9—C9A	1.381 (4)	N9'—C9A'	1.379 (4)
N9—C10	1.458 (4)	N9'—C10'	1.450 (4)
C9A—C1	1.396 (4)	C9A'—C1'	1.395 (4)
C9A—C4A	1.410 (4)	C9A'—C4A'	1.411 (4)
C10—C11	1.509 (4)	C10'—C11'	1.511 (4)
C10—H10A	1.00 (3)	C10'—H10C	0.98 (3)
C10—H10B	0.95 (3)	C10'—H10D	1.00 (3)
C11—C12	1.393 (4)	C11'—C12'	1.388 (4)
C11—C16	1.384 (4)	C11'—C16'	1.386 (4)
C12—H12	1.02 (4)	C12'—C13'	1.395 (4)
C13—C12	1.386 (4)	C12'—H12'	1.09 (3)
C13—H13	0.98 (3)	C13'—H13'	0.95 (3)
C14—C13	1.379 (5)	C14'—C13'	1.371 (5)
C14—C15	1.379 (5)	C14'—C15'	1.375 (6)
C14—H14	0.99 (4)	C14'—H14'	1.03 (4)
C15—C16	1.387 (4)	C15'—H15'	1.00 (4)
C15—H15	1.04 (3)	C16'—C15'	1.388 (5)
C16—H16	0.99 (3)	C16'—H16'	0.94 (3)
C2—C1—C9A	116.8 (4)	C2'—C1'—C9A'	117.5 (3)
C2—C1—H1	122.0 (17)	C2'—C1'—H1'	121.1 (16)
C9A—C1—H1	121.0 (17)	C9A'—C1'—H1'	121.2 (16)

C1—C2—H2	119.3 (18)	C1'—C2'—C3'	121.4 (3)
C3—C2—C1	122.1 (3)	C1'—C2'—H2'	112.6 (17)
C3—C2—H2	118.6 (18)	C3'—C2'—H2'	126.0 (17)
C2—C3—H3	119.3	C2'—C3'—H3'	120.5 (19)
C4—C3—C2	121.3 (3)	C4'—C3'—C2'	121.0 (3)
C4—C3—H3	119.3	C4'—C3'—H3'	118.5 (19)
C3—C4—C4A	118.2 (4)	C3'—C4'—C4A'	118.8 (3)
C3—C4—H4	119.5 (19)	C3'—C4'—H4'	117 (2)
C4A—C4—H4	122.2 (19)	C4A'—C4'—H4'	124 (2)
C4—C4A—C9A	119.9 (3)	C4'—C4A'—C5A'	134.1 (3)
C4—C4A—C5A	133.8 (3)	C4'—C4A'—C9A'	119.7 (3)
C9A—C4A—C5A	106.3 (3)	C9A'—C4A'—C5A'	106.2 (2)
C5A—C5—H5	122 (2)	C5A'—C5'—H5'	121.5 (18)
C6—C5—C5A	118.8 (3)	C6'—C5'—C5A'	118.7 (3)
C6—C5—H5	119 (2)	C6'—C5'—H5'	119.6 (19)
C5—C5A—C4A	133.4 (3)	C5'—C5A'—C4A'	133.5 (3)
C5—C5A—C8A	119.8 (3)	C5'—C5A'—C8A'	119.5 (3)
C8A—C5A—C4A	106.8 (3)	C8A'—C5A'—C4A'	107.1 (2)
C5—C6—C7	120.7 (3)	C5'—C6'—C7'	120.7 (3)
C5—C6—H6	119.5 (19)	C5'—C6'—H6'	117 (2)
C7—C6—H6	119.8 (19)	C7'—C6'—H6'	122 (2)
C6—C7—H7	120 (2)	C6'—C7'—H7'	118.9
C8—C7—C6	121.8 (3)	C8'—C7'—C6'	122.3 (3)
C8—C7—H7	118.5 (19)	C8'—C7'—H7'	118.9
C7—C8—C8A	117.2 (3)	C7'—C8'—H8'	121.4 (17)
C7—C8—H8	124.1 (18)	C8A'—C8'—C7'	117.0 (3)
C8A—C8—H8	118.6 (19)	C8A'—C8'—H8'	121.6 (17)
N9—C8A—C5A	109.2 (2)	N9'—C8A'—C5A'	108.3 (2)
N9—C8A—C8	129.2 (3)	C8'—C8A'—C5A'	121.8 (3)
C8—C8A—C5A	121.6 (3)	C8'—C8A'—N9'	129.9 (3)
C8A—N9—C10	126.8 (3)	C8A'—N9'—C10'	126.5 (2)
C9A—N9—C8A	108.1 (2)	C9A'—N9'—C8A'	109.0 (2)
C9A—N9—C10	124.9 (3)	C9A'—N9'—C10'	124.5 (2)
N9—C9A—C1	128.9 (3)	C1'—C9A'—C4A'	121.6 (3)
N9—C9A—C4A	109.5 (3)	N9'—C9A'—C1'	128.9 (3)
C1—C9A—C4A	121.6 (3)	N9'—C9A'—C4A'	109.5 (2)
N9—C10—C11	112.9 (2)	N9'—C10'—C11'	114.4 (2)
N9—C10—H10B	108.5 (19)	N9'—C10'—H10C	108.6 (19)
N9—C10—H10A	109.9 (18)	N9'—C10'—H10D	109.6 (18)
C11—C10—H10B	110.0 (18)	C11'—C10'—H10C	113.1 (19)
C11—C10—H10A	111.5 (18)	C11'—C10'—H10D	108.0 (18)
H10B—C10—H10A	104 (3)	H10C—C10'—H10D	103 (2)
C12—C11—C10	121.0 (3)	C12'—C11'—C10'	121.9 (3)
C16—C11—C10	120.1 (3)	C16'—C11'—C10'	118.7 (3)
C16—C11—C12	118.9 (3)	C16'—C11'—C12'	119.4 (3)
C11—C12—H12	121.7 (19)	C11'—C12'—C13'	119.3 (3)
C13—C12—C11	120.3 (3)	C11'—C12'—H12'	120.4 (18)
C13—C12—H12	118.0 (19)	C13'—C12'—H12'	120.1 (18)

C12—C13—H13	120 (2)	C12'—C13'—H13'	121 (2)
C14—C13—C12	120.3 (3)	C14'—C13'—C12'	121.1 (4)
C14—C13—H13	120 (2)	C14'—C13'—H13'	118 (2)
C13—C14—C15	119.8 (3)	C13'—C14'—C15'	119.4 (3)
C13—C14—H14	115 (2)	C13'—C14'—H14'	117 (2)
C15—C14—H14	125 (2)	C15'—C14'—H14'	124 (2)
C14—C15—C16	120.1 (3)	C14'—C15'—C16'	120.4 (4)
C14—C15—H15	120.5 (18)	C14'—C15'—H15'	117 (2)
C16—C15—H15	119.3 (18)	C16'—C15'—H15'	122 (2)
C11—C16—C15	120.7 (3)	C11'—C16'—C15'	120.3 (3)
C11—C16—H16	117.0 (17)	C11'—C16'—H16'	118 (2)
C15—C16—H16	122.3 (18)	C15'—C16'—H16'	122 (2)
C9A—C1—C2—C3	-1.1 (5)	C9A'—C1'—C2'—C3'	-0.8 (5)
C4—C3—C2—C1	1.0 (6)	C4'—C3'—C2'—C1'	0.3 (5)
C4A—C4—C3—C2	-0.2 (5)	C4A'—C4'—C3'—C2'	-0.2 (4)
C5A—C4A—C4—C3	177.8 (3)	C5A'—C4A'—C4'—C3'	178.8 (3)
C9A—C4A—C4—C3	-0.5 (5)	C9A'—C4A'—C4'—C3'	0.5 (4)
C5—C5A—C4A—C4	1.2 (6)	C4'—C4A'—C5A'—C5'	0.0 (6)
C5—C5A—C4A—C9A	179.7 (3)	C4'—C4A'—C5A'—C8A'	-178.6 (3)
C8A—C5A—C4A—C4	-179.4 (3)	C9A'—C4A'—C5A'—C5'	178.5 (3)
C8A—C5A—C4A—C9A	-0.9 (3)	C9A'—C4A'—C5A'—C8A'	-0.2 (3)
C8A—C5A—C5—C6	0.6 (4)	C5A'—C5'—C6'—C7'	-0.1 (5)
C4A—C5A—C5—C6	179.9 (3)	C4A'—C5A'—C5'—C6'	-177.8 (3)
C7—C6—C5—C5A	-0.2 (4)	C8A'—C5A'—C5'—C6'	0.7 (4)
C5—C6—C7—C8	-0.7 (5)	C8'—C7'—C6'—C5'	-1.0 (5)
C8A—C8—C7—C6	1.2 (4)	C8A'—C8'—C7'—C6'	1.5 (5)
C8—C8A—C5A—C4A	-179.5 (3)	N9'—C8A'—C5A'—C4A'	0.3 (3)
C8—C8A—C5A—C5	0.0 (4)	N9'—C8A'—C5A'—C5'	-178.5 (3)
N9—C8A—C5A—C4A	0.4 (3)	C8'—C8A'—C5A'—C4A'	178.7 (3)
N9—C8A—C5A—C5	179.9 (2)	C8'—C8A'—C5A'—C5'	-0.1 (4)
N9—C8A—C8—C7	179.2 (3)	N9'—C8A'—C8'—C7'	177.1 (3)
C5A—C8A—C8—C7	-0.8 (4)	C5A'—C8A'—C8'—C7'	-0.9 (4)
C9A—N9—C8A—C5A	0.2 (3)	C9A'—N9'—C8A'—C5A'	-0.3 (3)
C9A—N9—C8A—C8	-179.8 (3)	C9A'—N9'—C8A'—C8'	-178.6 (3)
C10—N9—C8A—C5A	-175.1 (3)	C10'—N9'—C8A'—C8'	3.0 (5)
C10—N9—C8A—C8	4.8 (5)	C10'—N9'—C8A'—C5A'	-178.7 (3)
C8A—N9—C9A—C1	178.6 (3)	C8A'—N9'—C9A'—C1'	179.9 (3)
C8A—N9—C9A—C4A	-0.8 (3)	C8A'—N9'—C9A'—C4A'	0.2 (3)
C10—N9—C9A—C1	-5.9 (5)	C10'—N9'—C9A'—C1'	-1.7 (5)
C10—N9—C9A—C4A	174.7 (3)	C10'—N9'—C9A'—C4A'	178.6 (3)
C8A—N9—C10—C11	104.4 (3)	C8A'—N9'—C10'—C11'	102.9 (3)
C9A—N9—C10—C11	-70.2 (4)	C9A'—N9'—C10'—C11'	-75.2 (4)
N9—C9A—C1—C2	-178.9 (3)	N9'—C9A'—C1'—C2'	-178.5 (3)
C4A—C9A—C1—C2	0.5 (5)	C4A'—C9A'—C1'—C2'	1.1 (4)
N9—C9A—C4A—C4	179.8 (3)	N9'—C9A'—C4A'—C4'	178.7 (2)
N9—C9A—C4A—C5A	1.1 (3)	N9'—C9A'—C4A'—C5A'	0.0 (3)
C1—C9A—C4A—C4	0.3 (4)	C1'—C9A'—C4A'—C4'	-1.0 (4)

C1—C9A—C4A—C5A	-178.4 (3)	C1'—C9A'—C4A'—C5A'	-179.7 (3)
N9—C10—C11—C12	-54.4 (4)	N9'—C10'—C11'—C16'	155.3 (3)
N9—C10—C11—C16	125.9 (3)	N9'—C10'—C11'—C12'	-26.9 (4)
C10—C11—C12—C13	-179.6 (3)	C10'—C11'—C12'—C13'	-175.9 (3)
C16—C11—C12—C13	0.0 (5)	C16'—C11'—C12'—C13'	1.9 (4)
C10—C11—C16—C15	179.8 (3)	C10'—C11'—C16'—C15'	175.8 (3)
C12—C11—C16—C15	0.2 (5)	C12'—C11'—C16'—C15'	-2.0 (5)
C14—C13—C12—C11	-0.2 (5)	C11'—C12'—C13'—C14'	-0.3 (5)
C15—C14—C13—C12	0.2 (5)	C15'—C14'—C13'—C12'	-1.1 (5)
C13—C14—C15—C16	-0.1 (5)	C13'—C14'—C15'—C16'	1.0 (5)
C14—C15—C16—C11	-0.1 (5)	C11'—C16'—C15'—C14'	0.5 (5)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C6—H6...Cg1 ⁱ	0.97 (4)	2.940 (4)	3.636 (5)	129.46 (5)
C10'—H10C...Cg1 ⁱⁱⁱ	0.98 (3)	2.787 (4)	3.700 (5)	154.92 (4)
C4'—H4'...Cg3 ⁱ	0.99 (4)	2.706 (4)	3.554 (4)	144.36 (5)

Symmetry codes: (i) $-x, y-1/2, -z+1/2$; (ii) $x, y+1, z$.