

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

rac-4-(2-Methoxyphenyl)-2,6-dimethylcyclohex-3-enecarboxylic acid

Songwen Xie,^{a*} Brian P. Fowler,^a Sara M. Deyo^a and Maren Pink^b

^aDepartment of Natural, Information, and Mathematical Sciences, Indiana University Kokomo, Kokomo, IN 46904-9003, USA, and ^bIndiana University Molecular Structure Center, Indiana University, Bloomington, IN 47405-7102, USA

Correspondence e-mail: soxie@iuk.edu


Received 17 May 2010; accepted 25 May 2010

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.048; wR factor = 0.138; data-to-parameter ratio = 16.6.

The title compound, $\text{C}_{16}\text{H}_{20}\text{O}_3$, was synthesized to study the hydrogen-bonding interactions of the two enantiomers in the solid state. Intermolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds produce centrosymmetric $R_2^2(8)$ rings which dimerize the two chiral enantiomers together through their carboxyl groups.

Related literature

In similar compounds previously reported (Xie *et al.*, 2002, 2007*a*, 2008*a,b*), the racemates also consist of carboxylic acid *RS* dimers. For the structure of the precursor, see: Xie *et al.* (2007*b*). The chirality of the title compound is solely generated by the presence of the double bond in the cyclohexene ring, see: Xie *et al.* (2004). For hydrogen-bond motifs, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{20}\text{O}_3$
 $M_r = 260.32$
 Monoclinic, $P2_1/c$
 $a = 14.2283$ (9) Å

$b = 7.1202$ (5) Å
 $c = 14.9517$ (10) Å
 $\beta = 106.069$ (2)°
 $V = 1455.55$ (17) Å³

$Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.08$ mm⁻¹

$T = 150$ K
 $0.25 \times 0.23 \times 0.07$ mm

Data collection

Bruker APEXII Kappa Duo diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.980$, $T_{\max} = 0.994$

11165 measured reflections
 2964 independent reflections
 2296 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.138$
 $S = 1.04$
 2964 reflections
 179 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.56$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.34$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{O}2-\text{H}2\cdots\text{O}1^i$	0.88 (3)	1.79 (3)	2.6640 (18)	177 (3)

Symmetry code: (i) $-x + 2, -y + 1, -z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SIR2004 (Burla *et al.*, 2005); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

SX, BF, and SD are grateful for the Grant-in-aid for Faculty Research from Indiana University Kokomo, as well as the Senior Research Grant from Indiana Academy of Science.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: OM2342).

References

- Bernstein, J., Davis, R., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
- Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Xie, S., Hou, Y., Meyers, C. Y. & Robinson, P. D. (2002). *Acta Cryst.* **E58**, o1460–o1462.
- Xie, S., Kenny, C. & Robinson, P. D. (2007*a*). *Acta Cryst.* **E63**, o3897.
- Xie, S., Kenny, C. & Robinson, P. D. (2007*b*). *Acta Cryst.* **E63**, o1660–o1662.
- Xie, S., Meyers, C. Y. & Robinson, P. D. (2004). *Acta Cryst.* **E60**, o1362–o1364.
- Xie, S., O'Hearn, C. R. & Robinson, P. D. (2008*a*). *Acta Cryst.* **E64**, o554.
- Xie, S., Stein, H. J. & Pink, M. (2008*b*). *Acta Cryst.* **E64**, o1869.

supporting information

Acta Cryst. (2010). E66, o1516 [https://doi.org/10.1107/S1600536810019732]

rac*-4-(2-Methoxyphenyl)-2,6-dimethylcyclohex-3-enecarboxylic acid*Songwen Xie, Brian P. Fowler, Sara M. Deyo and Maren Pink****S1. Comment**

The title carboxylic acid was prepared to study the interaction of the two enantiomers in the solid state. We have previously reported the structure of its precursor, which is achiral and forms hydrogen-bonded dimers (Xie *et al.*, 2007b). The chirality of the title compound is solely generated by the presence of the double bond in the cyclohexene ring (Xie *et al.*, 2004). The resultant racemate is made up of carboxylic acid RS dimers (Xie *et al.*, 2002, 2007a, 2008a,b). The structure and atom numbering are shown in Fig. 1, which illustrates the half-chair conformation of the cyclohexene ring. The torsion angles involving atoms C4, C5, C6, C1, and C2 are near 0°. The carboxyl group is almost perpendicular to the cyclohexene ring with an angle of 82.2° between the O1—C14—O2—C3 plane and the C1—C6 ring. The double bond between C5—C6 is not fully conjugated with the aromatic ring as shown by the C1—C6—C5 plane to benzene ring angle of 52.6°. Unlike other previously reported *para* substituted analogs and like other previously reported *meta* substituted analogs (Xie *et al.*, 2008b), the molecule also has a chiral axis due to the *ortho* methoxy substituent on the aromatic ring.

Fig. 2 shows the hydrogen bonding scheme. Atom O2 acts as a donor in an intermolecular hydrogen bond to atom O1, producing an R22(8) ring (Bernstein *et al.*, 1995), thus creating a hydrogen-bonded dimer. There is no evidence to suggest that weak directional interactions interconnect the dimers. Hydrogen bond geometry is given in Table 1.

S2. Experimental

The title carboxylic acid was synthesized following a similar method reported by Xie *et al.*, 2002. Purified compound was recrystallized from hexane-dichloromethane as colorless plates (m.p. 417–418 K).

S3. Refinement

All non-hydrogen atoms were refined with anisotropic displacement parameters. The hydrogen atoms not involved in hydrogen bonding were placed in ideal positions and refined as riding atoms with relative isotropic displacement parameters. H1 was freely refined.




Figure 1

The molecular structure showing thermal ellipsoids at the 50% probability level and the atom numbering scheme.




Figure 2

Hydrogen bonded dimer. Dashed lines represent hydrogen bonds (symmetry code: #1 $-x+2, -y+3, -z$).

4-(2-Methoxyphenyl)-2,6-dimethylcyclohex-3-enecarboxylic acid

Crystal data

$C_{16}H_{20}O_3$

$M_r = 260.32$

Monoclinic, $P2_1/c$

Hall symbol: $-P 2_1/c$

$a = 14.2283 (9) \text{ \AA}$

$b = 7.1202 (5) \text{ \AA}$

$c = 14.9517 (10) \text{ \AA}$

$\beta = 106.069 (2)^\circ$

$V = 1455.55 (17) \text{ \AA}^3$

$Z = 4$

$F(000) = 560$

$D_x = 1.188 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 4026 reflections

$\theta = 2.8\text{--}26.3^\circ$

$\mu = 0.08 \text{ mm}^{-1}$

$T = 150 \text{ K}$

Plate, colorless

$0.25 \times 0.23 \times 0.07 \text{ mm}$

Data collection

Bruker APEXII Kappa Duo
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
Detector resolution: 83.33 pixels mm⁻¹
 ω and φ scans
Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)
 $T_{\min} = 0.980$, $T_{\max} = 0.994$

11165 measured reflections
2964 independent reflections
2296 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.022$
 $\theta_{\max} = 26.4^\circ$, $\theta_{\min} = 1.5^\circ$
 $h = -16 \rightarrow 17$
 $k = -7 \rightarrow 8$
 $l = -13 \rightarrow 18$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.048$
 $wR(F^2) = 0.138$
 $S = 1.04$
2964 reflections
179 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H atoms treated by a mixture of independent
and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0653P)^2 + 0.6215P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.56 \text{ e } \text{Å}^{-3}$
 $\Delta\rho_{\min} = -0.34 \text{ e } \text{Å}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.88885 (9)	0.4779 (2)	0.00968 (10)	0.0602 (5)
O2	1.02302 (9)	0.3518 (2)	0.10376 (10)	0.0531 (4)
H2O	1.051 (2)	0.411 (4)	0.0670 (18)	0.082 (8)*
O3	0.55725 (8)	-0.14999 (17)	0.09567 (7)	0.0342 (3)
C1	0.76051 (16)	-0.0149 (3)	0.11315 (14)	0.0478 (5)
H1A	0.7060	-0.0803	0.0683	0.057*
H1B	0.8079	-0.1122	0.1446	0.057*
C2	0.81042 (14)	0.1106 (3)	0.05858 (12)	0.0418 (5)
H2	0.7585	0.1771	0.0100	0.050*
C3	0.87036 (12)	0.2591 (3)	0.12493 (12)	0.0366 (4)
H3	0.9175	0.1927	0.1775	0.044*
C4	0.80346 (14)	0.3809 (3)	0.16515 (13)	0.0409 (4)
H4	0.7610	0.4559	0.1129	0.049*
C5	0.73751 (16)	0.2583 (3)	0.20376 (14)	0.0479 (5)
H5	0.7053	0.3137	0.2450	0.057*

C6	0.72161 (11)	0.0741 (2)	0.18303 (11)	0.0300 (4)
C7	0.66838 (11)	-0.0451 (2)	0.23497 (10)	0.0279 (4)
C8	0.69907 (12)	-0.0459 (2)	0.33172 (11)	0.0322 (4)
H8	0.7515	0.0337	0.3629	0.039*
C9	0.65520 (14)	-0.1596 (2)	0.38384 (11)	0.0377 (4)
H9	0.6772	-0.1570	0.4499	0.045*
C10	0.57966 (14)	-0.2762 (3)	0.33929 (12)	0.0392 (4)
H10	0.5501	-0.3557	0.3748	0.047*
C11	0.54636 (13)	-0.2784 (2)	0.24267 (12)	0.0343 (4)
H11	0.4942	-0.3592	0.2122	0.041*
C12	0.58967 (11)	-0.1624 (2)	0.19099 (11)	0.0279 (4)
C13	0.87078 (18)	-0.0056 (4)	0.00957 (17)	0.0680 (7)
H13A	0.8984	0.0766	-0.0292	0.102*
H13B	0.8290	-0.1004	-0.0297	0.102*
H13C	0.9239	-0.0680	0.0560	0.102*
C14	0.92794 (13)	0.3744 (3)	0.07423 (12)	0.0409 (5)
C15	0.86046 (19)	0.5186 (3)	0.23835 (16)	0.0615 (6)
H15A	0.8146	0.5943	0.2615	0.092*
H15B	0.8997	0.6013	0.2105	0.092*
H15C	0.9037	0.4488	0.2901	0.092*
C16	0.47037 (14)	-0.2506 (3)	0.05064 (12)	0.0441 (5)
H16A	0.4516	-0.2223	-0.0161	0.066*
H16B	0.4175	-0.2129	0.0772	0.066*
H16C	0.4822	-0.3857	0.0599	0.066*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0321 (7)	0.0899 (12)	0.0542 (8)	-0.0105 (7)	0.0048 (6)	0.0408 (8)
O2	0.0299 (7)	0.0756 (11)	0.0510 (8)	-0.0106 (7)	0.0066 (6)	0.0317 (7)
O3	0.0347 (6)	0.0422 (7)	0.0275 (6)	-0.0110 (5)	0.0115 (5)	-0.0011 (5)
C1	0.0570 (12)	0.0433 (11)	0.0542 (11)	-0.0197 (9)	0.0341 (10)	-0.0117 (9)
C2	0.0365 (9)	0.0554 (12)	0.0387 (9)	-0.0140 (9)	0.0190 (8)	-0.0047 (8)
C3	0.0303 (9)	0.0444 (10)	0.0336 (8)	-0.0108 (8)	0.0066 (7)	0.0099 (7)
C4	0.0461 (11)	0.0374 (10)	0.0397 (9)	-0.0125 (8)	0.0124 (8)	0.0025 (8)
C5	0.0604 (13)	0.0390 (11)	0.0554 (11)	-0.0104 (9)	0.0346 (10)	-0.0058 (9)
C6	0.0270 (8)	0.0360 (9)	0.0285 (8)	-0.0060 (7)	0.0100 (6)	-0.0011 (7)
C7	0.0277 (8)	0.0292 (8)	0.0294 (8)	0.0019 (7)	0.0121 (6)	0.0008 (6)
C8	0.0318 (9)	0.0335 (9)	0.0315 (8)	0.0020 (7)	0.0091 (7)	-0.0001 (7)
C9	0.0475 (10)	0.0398 (10)	0.0274 (8)	0.0071 (8)	0.0129 (7)	0.0056 (7)
C10	0.0515 (11)	0.0355 (9)	0.0374 (9)	-0.0013 (8)	0.0236 (8)	0.0088 (8)
C11	0.0382 (9)	0.0314 (9)	0.0374 (9)	-0.0046 (7)	0.0173 (7)	0.0010 (7)
C12	0.0301 (8)	0.0279 (8)	0.0291 (8)	0.0011 (7)	0.0138 (6)	0.0007 (6)
C13	0.0605 (14)	0.0884 (18)	0.0704 (15)	-0.0219 (13)	0.0437 (12)	-0.0283 (13)
C14	0.0299 (9)	0.0529 (11)	0.0365 (9)	-0.0122 (8)	0.0036 (7)	0.0126 (8)
C15	0.0778 (16)	0.0467 (12)	0.0576 (13)	-0.0251 (12)	0.0148 (12)	-0.0074 (10)
C16	0.0473 (11)	0.0508 (11)	0.0334 (9)	-0.0185 (9)	0.0098 (8)	-0.0066 (8)

Geometric parameters (Å, °)

O1—C14	1.219 (2)	C6—C7	1.491 (2)
O2—C14	1.312 (2)	C7—C8	1.391 (2)
O2—H2O	0.88 (3)	C7—C12	1.406 (2)
O3—C12	1.3740 (19)	C8—C9	1.386 (2)
O3—C16	1.426 (2)	C8—H8	0.9500
C1—C6	1.456 (2)	C9—C10	1.376 (3)
C1—C2	1.513 (2)	C9—H9	0.9500
C1—H1A	0.9900	C10—C11	1.390 (2)
C1—H1B	0.9900	C10—H10	0.9500
C2—C13	1.519 (3)	C11—C12	1.387 (2)
C2—C3	1.537 (2)	C11—H11	0.9500
C2—H2	1.0000	C13—H13A	0.9800
C3—C14	1.504 (2)	C13—H13B	0.9800
C3—C4	1.528 (3)	C13—H13C	0.9800
C3—H3	1.0000	C15—H15A	0.9800
C4—C5	1.508 (2)	C15—H15B	0.9800
C4—C15	1.525 (3)	C15—H15C	0.9800
C4—H4	1.0000	C16—H16A	0.9800
C5—C6	1.352 (3)	C16—H16B	0.9800
C5—H5	0.9500	C16—H16C	0.9800
C14—O2—H2O	110.0 (18)	C9—C8—H8	119.1
C12—O3—C16	117.10 (12)	C7—C8—H8	119.1
C6—C1—C2	117.26 (16)	C10—C9—C8	119.56 (15)
C6—C1—H1A	108.0	C10—C9—H9	120.2
C2—C1—H1A	108.0	C8—C9—H9	120.2
C6—C1—H1B	108.0	C9—C10—C11	120.40 (16)
C2—C1—H1B	108.0	C9—C10—H10	119.8
H1A—C1—H1B	107.2	C11—C10—H10	119.8
C1—C2—C13	110.54 (18)	C12—C11—C10	119.74 (16)
C1—C2—C3	108.53 (14)	C12—C11—H11	120.1
C13—C2—C3	113.54 (16)	C10—C11—H11	120.1
C1—C2—H2	108.0	O3—C12—C11	122.89 (15)
C13—C2—H2	108.0	O3—C12—C7	116.24 (13)
C3—C2—H2	108.0	C11—C12—C7	120.84 (15)
C14—C3—C4	111.98 (15)	C2—C13—H13A	109.5
C14—C3—C2	109.42 (14)	C2—C13—H13B	109.5
C4—C3—C2	110.45 (14)	H13A—C13—H13B	109.5
C14—C3—H3	108.3	C2—C13—H13C	109.5
C4—C3—H3	108.3	H13A—C13—H13C	109.5
C2—C3—H3	108.3	H13B—C13—H13C	109.5
C5—C4—C15	111.17 (16)	O1—C14—O2	122.86 (16)
C5—C4—C3	110.10 (15)	O1—C14—C3	122.40 (16)
C15—C4—C3	112.43 (17)	O2—C14—C3	114.73 (15)
C5—C4—H4	107.6	C4—C15—H15A	109.5
C15—C4—H4	107.6	C4—C15—H15B	109.5

C3—C4—H4	107.6	H15A—C15—H15B	109.5
C6—C5—C4	123.77 (17)	C4—C15—H15C	109.5
C6—C5—H5	118.1	H15A—C15—H15C	109.5
C4—C5—H5	118.1	H15B—C15—H15C	109.5
C5—C6—C1	120.94 (16)	O3—C16—H16A	109.5
C5—C6—C7	120.64 (15)	O3—C16—H16B	109.5
C1—C6—C7	118.33 (15)	H16A—C16—H16B	109.5
C8—C7—C12	117.67 (14)	O3—C16—H16C	109.5
C8—C7—C6	119.05 (14)	H16A—C16—H16C	109.5
C12—C7—C6	123.25 (14)	H16B—C16—H16C	109.5
C9—C8—C7	121.77 (16)		
C6—C1—C2—C13	164.10 (19)	C1—C6—C7—C12	-53.0 (2)
C6—C1—C2—C3	39.0 (2)	C12—C7—C8—C9	0.9 (2)
C1—C2—C3—C14	174.85 (16)	C6—C7—C8—C9	-177.28 (15)
C13—C2—C3—C14	51.5 (2)	C7—C8—C9—C10	0.4 (3)
C1—C2—C3—C4	-61.4 (2)	C8—C9—C10—C11	-0.9 (3)
C13—C2—C3—C4	175.22 (18)	C9—C10—C11—C12	0.0 (3)
C14—C3—C4—C5	172.34 (15)	C16—O3—C12—C11	5.2 (2)
C2—C3—C4—C5	50.1 (2)	C16—O3—C12—C7	-172.68 (15)
C14—C3—C4—C15	-63.1 (2)	C10—C11—C12—O3	-176.46 (16)
C2—C3—C4—C15	174.67 (15)	C10—C11—C12—C7	1.4 (2)
C15—C4—C5—C6	-141.8 (2)	C8—C7—C12—O3	176.16 (14)
C3—C4—C5—C6	-16.6 (3)	C6—C7—C12—O3	-5.7 (2)
C4—C5—C6—C1	-6.3 (3)	C8—C7—C12—C11	-1.8 (2)
C4—C5—C6—C7	170.29 (17)	C6—C7—C12—C11	176.29 (15)
C2—C1—C6—C5	-5.8 (3)	C4—C3—C14—O1	-56.8 (3)
C2—C1—C6—C7	177.46 (16)	C2—C3—C14—O1	66.0 (3)
C5—C6—C7—C8	-51.6 (2)	C4—C3—C14—O2	124.50 (18)
C1—C6—C7—C8	125.06 (18)	C2—C3—C14—O2	-112.68 (19)
C5—C6—C7—C12	130.28 (19)		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2O \cdots O1 ⁱ	0.88 (3)	1.79 (3)	2.6640 (18)	177 (3)

Symmetry code: (i) $-x+2, -y+1, -z$.